



Induction of Antiproliferative Effect by Diosgenin Through P53 Activation, Bcl-2 Down Regulation and Caspase-3 Modulation Activity in Human Colon Cancer (Caco-2) Cells

Entissar S. Al-Suhaibani

Associate professor of genetics. King Saud University, Riyadh, KSA. Riyadh 11342

Nermin A. El-Morshedi

Assistant professor of Histology, cell biology and genetics.

ABSTRACT

Previously, a plant steroid, diosgenin, altered cell cycle distribution and induced apoptosis in human osteosarcoma cell line. The study objective was to investigate if antiproliferative effect of diosgenin was similar for human colon cancer (Caco-2) cell. Moreover, this work focused on the mitochondrial pathway. Cells were incubated with diosgenin concentrations, 60µM/L, 80µM/L, 100µM/L and 120µM/L for 72h at 37°C and 5%CO₂. Diosgenin induced cells death in a dose-dependent manner. Diosgenin-treated cells had an important antiproliferative effect, additionally, a typical characteristics of apoptosis including DNA fragmentation which is caspase-3 dependent with caspase-3 over expression and fall of mitochondrial membrane potential which appeared by Bcl-2 expression down regulating, p53 activation as signal for cell cycle arrest and cytological alterations. Future study will may deal with further investigations of diosgenin possible usages as a new alternative or complementary chemotherapeutic agent for human cancer types specially colon cancer type.

KEYWORDS : diosgenin, Caco-2 Cells, p53, Bcl-2, caspase-3.

INTRODUCTION

Programmed cell death or apoptosis plays an important role in normal development and is impaired in many types of cancer. Apoptosis occurs under a variety of physiological and pathological conditions that control the development and homeostasis of multicellular organisms¹. The major apoptotic pathways can be divided into caspase- and mitochondria- dependent pathways, according to caspase-3 activation which is generally considered to be a key hallmark of apoptosis². Apoptosis is characterized by chromatin condensation and DNA fragmentation, and is mediated by the cysteine protease family called caspase-3³.

The tumor suppressor protein p53 is a principal factor in regulation of growth arrest as well as apoptosis. Many apoptotic signals are mediated to the cell death machinery via p53. It interacts with other proteins or functions as a transcription factor⁴. Indeed, in response to various types of stress, p53 becomes activated and, as a consequence, cells can undergo marked phenotype changes, ranging from increased DNA repair to senescence and apoptosis⁵. P53 has been shown to be involved in the induction of apoptosis, cell-cycle arrest and differentiation responses that prevent further proliferation of stressed or damaged cells and so protect from the outgrowth of cells harboring malignant alterations. P53 role in the repair of DNA damage has also been described and the ability of p53 to induce reversible cell-cycle arrest may contribute to the ability of cells to repair and recover from damage before reentering a normal proliferative state⁶.

Mitochondria are involved in a variety of key events, including release of caspase-3 activators, changes in electron transport, loss of mitochondrial membrane potential, and participation of both pro- and anti-apoptotic Bcl-2 protein⁷. Alterations in mitochondrial structure and function have been shown to play a crucial role in caspase-3-dependent apoptosis and Bcl-2 expression⁸. Bcl-2 is the founding member of family of genes that either prevents or promotes cellular apoptosis. Bcl-2 itself is an anti-apoptotic gene that prevents initiation steps of apoptosis and programmed cell death⁹.

Over the past decade, there is continuous increase in colorectal carcinoma in the world as the most common malignant diseases. Colorectal cancer is the cause of more than 1/2 million deaths worldwide, and it was ranked as the third leading cause of cancer-related death after lung cancer and stomach cancer¹⁰. Epidemiological studies have shown strong evidence that diet and lifestyle play an important role in preventing cancer. In particular, an increased consumption of fruits and vegetables is associated with decreasing in cancer onset and mortality¹¹. Previously, we showed that diosgenin, a plant steroid, altered cell cycle and induced apoptosis in the human osteosarcoma

1547 cell line¹². In order to know if results obtained with diosgenin on 1547 cells could be extended to other types of cancer cells, we tested this molecule on laryngocarcinoma HEP-2 cells and melanoma M4Beu cells¹³. Moreover, to gain further insight into the mechanisms by which diosgenin induces apoptosis in human cancer cells, we examined in this study the functional status of caspase-3 on apoptosis rate and the effect of diosgenin on mitochondrial integrity in human colon cancer cell lines (Caco-2 cells).

MATERIALS AND METHODS

Chemical reagents: Diosgenin, MTT salt or 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl tetrazolium bromide, dimethylsulfoxide (DMSO), commercial methanol, commercial ethanol, commercial acetone, Tris-HCl, edetic acid, Triton-X100, RNase A, proteinase K, NaCl, 2-propanol, phosphate-buffered saline (PBS), ethidium bromide, agarose gel, Peroxidase, trypsin, Hematoxylin and eosin (Hx & E) stain, rabbit polyclonal antibodies against cleaved caspase-3, primary monoclonal antibody against Bcl-2 and against p53, AB reagent, biotinylated immunoglobulin secondary antibody and Tween 20 were purchased from Sigma-Aldrich, Egypt.

Cell line and cell culture: Caco-2 cell line, was obtained from American Type Culture Collection (ATCC, USA). They were sub-cultured as mono-layer according to the instructions provided by ATCC in Dulbecco modified Eagle medium (DMEM) supplemented with 10% heat inactivated (56°C, 30min) fetal bovine serum, 2mmol/L L-glutamine, 100U/mL Penicillin-Streptomycin and 100U/mL Amphotericin B at 37°C in a humidified atmosphere of 5% CO₂. Cells were used when monolayer reached 80% confluence in all experiments. Cell propagation media was purchased from Invitrogen (Carlsbad, CA).

Methods: 1. Cell Viability Assay: In vitro evaluation of antiproliferative effect: growth inhibition was evaluated by MTT assay. MTT salt or 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl tetrazolium bromide was reduced by mitochondrial dehydrogenases to water blue insoluble formazans¹⁴. Viable cell number/well is directly proportional to formazans production. 8.25×10³ cells were seeded into each well of 96-well plate, incubated with culture medium overnight (12h), replaced with fresh medium containing diosgenin at concentrations: 60µM/L, 80µM/L, 100µM/L and 120µM/L for 72h at 37°C in an incubator with 5%CO₂. After incubation, diosgenin modified medium was replaced by 100µL of MTT (0.5mg/mL) medium for incubation (3h at 37°C and 5%CO₂). MTT medium was then replaced with 100µL of DMSO and left for 10min on a platform shaker to solubilize converted formazan. The absorbance values were determined at 570nm test wavelength and 630nm reference wavelength (Spekol 1200 spectrophotometer). Untreated cells were as a positive control cells and all values were

correlated with this set of data. The experiment was performed in triplicates. Inhibition Percentage=[1-(net Absorbance of treated well/net Absorbance of control well)]x100%, then was plotted against diosgenin concentrations.

2.Determination of DNA fragmentation by DNA laddering assay:

cells were seeded in 60-mm petri dishes at density 4×10^5 cells/plate (treated cells by IC_{50} concentration of diosgenin or positive control cells). Adherent and floating cells were collected by centrifugation at $1000 \times g/5$ min. Cell pellet was suspended in cell lysis buffer (Tris-HCl 10mmol/L pH7.4, edetic acid 10mmol/L pH8.0, Triton-X100 0.5%) and kept at $4^\circ C/10$ min then, lysate was centrifuged at $25,000 \times g/20$ min. Supernatant was incubated with RNase A $40 \mu g/L/1h$ ($37^\circ C$), incubated with proteinase K $40 \mu g/L/1h$ ($37^\circ C$), mixed with NaCl 0.5mol/L and 50% 2-propanol overnight ($-20^\circ C$), then centrifuged at $25,000 \times g/15$ min. After drying, DNA was dissolved in buffer (Tris-HCl 10mmol/L pH7.4, edetic acid 1mmol/L pH 8.0) and separated by 2% agarose gel electrophoresis at 100V for 50min. DNA was visualized under ultraviolet light after staining with ethidium bromide¹⁵.

3.Cytological changes investigation: detached and trypsinized cells (IC_{50} concentration of diosgenin treated cells and positive control cells) were collected and centrifuged at 2000 rpm for 5min. Cell pellet was re-suspended with $100 \mu L$ of PBS (pH7.3). $10 \mu L$ of the suspension were smeared on a glass slide, allowed to air-dry, fixed with cool methanol for 5min before proceeding by Hx&E stain and examined under light microscope¹⁶.

4.Immunocytochemical investigations: by detection of Bcl-2, p53 and Caspase-3 by immunocytochemistry staining kits. The procedure was done according to the manufacturer's instructions, simplified as follows: 1-2 drops of Peroxidase was applied to cells (IC_{50} concentration of diosgenin treated cells and positive control cells) on the slide (10min), followed by blocking solution (10min). Cells were fixed in ethanol:acetone (9:1) for 30min at $-20^\circ C$ and then rinsed again with cold PBS at room temperature. Cells were incubated overnight with rabbit polyclonal antibodies against cleaved caspase-3 at $4^\circ C$, then AB reagent and substrate-chromogen mixture (30min). Between each step, the slide was washed with washing buffer (PBS) with 0.1% Tween 20). Cells were incubated overnight with primary monoclonal antibody against Bcl-2 and against p53 at dilution of 1:75 at $4^\circ C$, then in Tris buffer and biotinylated immunoglobulin secondary antibody was used¹⁷. The slides were then mounted and examined under light microscope.

5.Statistical analysis: results were presented as mean \pm standard deviations (SD). Analysis of variance (ANOVA) for two variables (Two Way-ANOVA) was used together with student t-test. Significant analysis of variance results were subjected to post hoc. Statistical significance was set at $P < 0.05$ and high significance was set at $P \leq 0.01$ ¹⁸.

RESULTS

1.Cell viability assay: In vitro evaluation of antiproliferation effect.

Cytotoxic effect of diosgenin concentrations ($60 \mu M$, $80 \mu M$, $100 \mu M$ and $120 \mu M$)/72h on Caco-2 cells was determined by MTT assay (Figure 1). Cells number started to reduce immediately after treatment with diosgenin concentrations in a dose dependent manner. All concentrations were found to be high significantly different ($P \leq 0.01$) in respect to their antiproliferative and apoptotic effects when compared with positive control cells. Cell inhibition percentage was gradually increased with diosgenin concentration increasing and 90% of cell inhibition was observed in treated cells with $120 \mu M/72h$. Cell proliferation reduced about 20% and 32% when cells were treated with $60 \mu M$ and $80 \mu M$ for 72h, respectively. Cells proliferation decreased to 51% as treated with concentration $100 \mu M/72h$.

2.Determination of DNA fragmentation by DNA laddering assay.

DNA degradation into multiple internucleosomal fragments is a distinct biochemical hallmark for apoptosis. Nuclear DNA isolated from Caco-2 cancer cells was separated by agarose gel electrophoresis and stained with ethidium bromide, and a typical ladder formation was observed upon 72h when treated with diosgenin concentration at $100 \mu M$ whereas untreated cells did not show typical ladder (Figure 2). Results indicated that diosgenin induced DNA fragmentation which

was caused by apoptosis.

3.Cytological changes investigation.

Positive control cells group had round nuclei, distinct small nucleoli and homogeneous chromatin with an accentuated nuclear membrane (Figure 3a). After Caco-2 cells treatment by diosgenin concentration at $100 \mu M/72h$, apoptotic cells were identified by a series morphological changes as an important experimental proof of underlying processes alterations appeared as: bleb plasma membrane, cellular shrinkage, chromatin condensation granules, vacuolated cytoplasm, degrading nucleus and apoptotic bodies formation were observed (Figure 3b, 3c and 3d).

4.Immunocytochemical investigation.

After Caco-2 cells treatment by diosgenin concentration at $100 \mu M/72h$, the reaction of caspase-3 protein was considered positive (over expression of caspase-3 protein) when over 50% of treated tumor cells had a clear brown cytoplasm staining, with slight degrading in the intensity in the same field (Figures 3f). Specially those fields that had necrotic or apoptotic nucleus as sign for diosgenin treatment effect, but fields of positive control cells have negative reaction of caspase-3 (cytoplasm did not show the brownish reaction stain) (Figure 3e). On the other hand regarding to the positive control Caco-2 cells, Bcl-2 protein reaction was considered positive (over expression of Bcl-2 protein) when over 55% of cells had nuclear membrane, mitochondrial outer membrane and endoplasmic reticulum membrane brown staining, with slight intensity degrading in the same field (Figures 3g). After Caco-2 cells treatment by diosgenin concentration at $100 \mu M/72h$, those fields that had necrotic or apoptotic nucleus as sign for diosgenin apoptotic effect with Bcl-2 negative reaction (faint to non-brown stain) (Figure 3h). Also, when applying p53 stain, p53 protein reaction in the positive control Caco-2 cells, was showed negative reaction (no brown stain) (Figure 3j). Treated Caco-2 cells, those fields had necrotic or apoptotic nucleus for diosgenin effect showed p53 positive reaction (over expression of p53 protein) when over 55% of cells had nuclear brown staining, with slight intensity degrading in the same field (Figures 3i).




Figure 1: Effect of diosgenin with different concentrations on the cells viability of Caco-2 cells. The experiment was performed in triplicates and values means were calculated [mean \pm SD, n (for each concentration)=4].




Figure 2: DNA fragmentation by DNA laddering assay of extracted DNA from diosgenin treated cells and positive control cells. DNA laddering, typical for apoptotic cells, which were visible in treated Coca-2 cells (T), and there was no any apoptotic features in the positive untreated cells (C) where M indicating marker.




Figure 3: Cells in different stages of apoptosis in treated cells are easily distinguishable. Cell with normal morphology (a). Complete apoptotic cell (b). Nuclear condensation is evident in cells (dark, condensed and irregular rounded nucleus), bleb membrane and cell shrinkage (c). Degradation of nucleus, vacuolated cytoplasm with ap-

optotic bodies (d). Immunocytochemistry of caspase-3 protein. Control positive cell showing cytoplasm negative reaction for caspase-3 protein (e). Treated cell showing cytoplasm positive reaction for caspase-3 protein (f). Control positive cell showing Bcl-2 protein nuclear membrane, mitochondrial outer membrane and endoplasmic reticulum membrane showing brownish positive reaction (g). Treated cell showing negative reaction of apoptotic cell apoptosis (h). Treated cell showing p53 protein nuclear positive reaction (i). Control positive cell showing nuclear negative reaction (j).

DISCUSSION

It is of great importance to understand the mechanisms of apoptosis in cancer cells, as apoptosis is believed to be one of the major consequences of anticancer drug treatment against malignancies¹⁹. Recently, we showed that three plant steroids had pro-apoptotic properties on osteosarcoma cells, as diosgenin is the most effective in induction of apoptosis²⁰. The aim of this study is to verify if its action was also similar on other human colon cancer cell lines such as Caco-2 cells. Recently, we compared diosgenin with two other plant steroids (hecogenin and tigogenin) in the same cell lines and among these three plant steroids, diosgenin appeared to be the most effective in inducing cell death¹². Our results showed that diosgenin strongly inhibited proliferation of Caco-2 cells and blocked the cell cycle as previously described¹³.

It is now established that the tumor suppressor p53 inhibits cell growth through activation of cell cycle arrest and apoptosis²¹. Regulation of p53 activity is through multiple mechanisms, one of which is phosphorylation. Indeed, it was established that phosphorylation of p53 protein may play a critical role in its stabilization, up-regulation, and functional activation²².

p53 has been described as 'The Guardian of the genome', referring to its role in conserving stability by preventing genome mutation²³. Upon genotoxic and other stress, p53 protein levels increase. Activated p53 releases signal to cells to undergo growth arrest, cell differentiation or apoptosis²⁴. Caco-2 cells which were treated with diosgenin exhibited increased levels of p53 expression at concentration of 100 μ M/72h, which suggested that p53 involved in diosgenin-induced Caco-2 cell death. Bcl-2 family of proteins serves as critical regulators of pathways involved in apoptosis. The main protagonists are suggested to be anti-apoptotic and known as Bcl-2⁸. Caco-2 cells which was treated with diosgenin exhibited reduced levels of Bcl-2 expression. These results suggested that the mitochondrial pathway was involved in diosgenin-induced Caco-2 cell death.

Apoptosis, as programmed cell death, is a highly organized cell death process characterized by an early obvious condensation of nuclear chromatin, loss of plasma membrane phospholipid asymmetry, activation of nucleases, enzymatic cleavage of DNA into oligonucleosomal fragments and segmentation of the cells into membrane-bound apoptotic bodies²⁵. DNA fragmentation, a hallmark of apoptosis, is regulated by a specific nuclease called caspase-activated DNase and its inhibitor²⁶. Apoptosis has specific signals instructing the cells with specific morphological change as plasma and nuclear membrane blebbings, chromatin condensation, proteases activation and DNA fragmentation that are considered as landmarks of the apoptotic process²⁷. That was agreed with the results of recent study after treatment by diosgenin. Diosgenin decreased the viable percentage of cell number (dose dependent effect) and induced apoptosis of Caco-2 cells. Therefore, we may presume that as primary mechanism involved in diosgenin growth-inhibitory effects as it considered main apoptotic signals. In response to apoptogenic stimuli, the mitochondrial protein apoptosis-inducing factor (AIF) translocates through the outer mitochondrial membrane to the cytosol and to the nucleus, resulting in the induction of nuclear chromatin condensation and large DNA fragmentation in a caspase-independent manner²⁸. It is well known that a family of cysteinyl proteases, caspases, is involved in the apoptotic cell death. Caspase-3, one of the active executioners, promotes apoptosis by cleaving cellular substrates such as ICAD²⁹. That was agreed in the recent study by caspase-3 expression inhibition. We found that diosgenin induced caspase-3 activity in Caco-2 cancer cell line.

CONCLUSION

In summary, we demonstrated that diosgenin caused an inhibition of cell growth with apoptosis induction by DNA fragmentation and p53

activation in human colon (Caco-2) cancer cell line. Moreover, a large part of our study essentially focused on the mitochondrial pathway and we investigated that diosgenin's action was caspase-3 dependent according to its inhibitory effect on Bcl-2 expression. There was also noticeable cytological alterations. These new findings suggest that diosgenin-induced effects may have novel therapeutic applications for the treatment of different cancer type as previously described for osteosarcoma. Future in vitro and in vivo study will may deal with further investigations of the possible usages of diosgenin as a new alternative chemotherapeutic agent.

ACKNOWLEDGMENTS

We are grateful for all of Nile center for experimental researches team specially Miss. Noha T. Badawy (Department of cell culture) for her kindly support in the cell culture and drug induction stage.

REFERENCES

- Gamet-Payastre, L. and Lumeau, S. (2000). Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in HT29 human colon cancer cells. *Cancer Res.*, 60:1426-1429. | 2.Kidd, V. (1998). Proteolytic activities that mediate apoptosis. *Annu. Rev. Physiol.*, 60:533-538. | 3.Sarin, A.; Elias, K. and Haddad, P. (1998). Caspase dependence of target cell damage induced by cytotoxic lymphocytes. *J. Immunol.*, 161:2810-2813. | 4.Jeffers, J.; Parganas, E.; Wang, J.; Brennan, J.; Chittenden, T. and Ihle, J. (2003). Puma is an essential mediator of p53-dependent and -independent apoptotic pathways. *Cancer Cell*, 4:321-328. | 5.Polyak, K.; Xia, Y.; Kinzler, K. and Vogelstein, B. (1997). A model for p53-induced apoptosis. *Nature.*, 389:300-305. | 6.Polyak, K.; Waldman, T.; He, C.; Kinzler, W. and Vogelstein, B. (1996). Genetic determinants of p53-induced apoptosis and growth arrest. *Genes Dev.*, 10:1945-1952. | 7.Adams, J. and Cory, S. (1998). Bcl-2 protein family: arbiters of cell survival. *Science.*, 281:1322-1325. | 8.Wolvetang, E.; Perez, J.; Roig, T.; Manzano, A.; Larm, J. and Moutsoulas, P. (1996). Apoptosis induced by inhibitors of plasma membrane NADH-oxidase involves Bcl-2 and calcineurin. *Cell Growth Differ.*, 7:1315-1318. | 9.An, W.; Bermudez, J.; Gamez, A.; Wang, M. and Tashiro, S. (2004). Norcantharidin induces human melanoma A375-S2 cell apoptosis through mitochondrial and caspase pathways. *J. Korean. Med. Sci.*, 19:560-563. | 10.Mayer, R. (2009). Targeted therapy for advanced colorectal cancer-more is not better. *J. Med.*, 360:623-625. | 11.Murillo, G.; Hoard, M.; Naithani, R. and Mehta, R. (2008). Efficacy of herbal products in colorectal cancer prevention. *Current Colorectal Cancer Reports.*, 4:34-42. | 12.Corbiere, C.; Liagre, B. and Bianchi, A. (2003). Different contribution of apoptosis to the antiproliferative effects of diosgenin and other plant steroids, hecogenin and tigogenin, on human 1547 osteosarcoma cells. *Int. J. Oncol.*, 22:899-905. | 13.Moalic, S.; Liagre, B. and Corbiere, C. (2001). A plant steroid, diosgenin, induces apoptosis, cell cycle arrest and COX activity in osteosarcoma cells. *FEBS. Lett.*, 506:225-30. | 14.Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods.*, 65:55-63. | 15.Zhang, Y.; Chen, X.; Liu, J. and Wong, T. (2003). DNA excision repair system of highly radioresistant bacterium *Deinococcus radiodurans* is facilitated by pentose phosphate pathway. *Mol. Microbiol.*, 48:1317-1323. | 16.John, A. and Abraham, S. (1991). Cytological changes produced by red pepper in mitotic cells of *Vicia faba* L. *Caryologia.*, 44:325-331. | 17.Yoon, K.; Nakamura, Y.; Doubek, D. and Arakawa, H. (2004). Identification of ALDH4 as p53, caspase-3 and Bcl-2-inhibitor gene and its protective role in cellular stresses. *J. Hum. Genet.*, 49:134-140. | 18.Snedecor, G. and Cochran, W. (1980). *Statistical Methods*, 7th ed Iowa State univ. Press, Iowa, U.S.A. | 19.Vousden K. (2002). Activation of the p53 tumor suppressor protein. *Biochim. Biophys. Acta.*, 1602:47-59. | 20.Moalic, S.; Liagre, B. and Corbiere, C. (2001). A plant steroid, diosgenin, induces apoptosis, cell cycle arrest and COX activity in osteosarcoma cells. *FEBS. Lett.*, 2506:225-30. | 21.Moll, U. and Zaika, A. (2001). Nuclear and mitochondrial apoptotic pathways of p53. *FEBS. Lett.*, 493:65-9. | 22.el-Deiry, W.; Tokino, T.; Velculescu, V. et al., (1993). WAF1, a potential mediator of p53 tumor suppression. *Cell.*, 75:817-825. | 23.Vogelstein, B.; Zweier, L.; Lane, D. and Levine, A. (2000). Surfing the p53 network. *Nature.*, 408:307-310. | 24.Contente, A.; Dittmer, A.; Koch, M.; Roth, J. and Dobbelstein, M. (2002). A polymorphic microsatellite that mediates induction of PIG3 by p53. *Nat. Genet.*, 30:315-320. | 25.Boada, J.; Roig, T.; Perez, X.; Bartrons, R. and Cascante, M. (2000). Cells overexpressing fructose-2,6-bisphosphatase showed enhanced pentose phosphate pathway flux and resistance to oxidative stress. *FEBS. Lett.*, 480:261-264. | 26.Hengartner, M. (2000). The biochemistry of apoptosis. *Nature.*, 407:770-776. | 27.Zamzami, N.; Susin, S.; Marchetti, P. et al. (1996). Mitochondrial control of nuclear apoptosis. *J. Exp. Med.*, 183:1533-1544. | 28.Daugas, E.; Susin, S. and Zamzami, N. (2000). Mitochondrio-nuclear translocation of AIF in apoptosis and necrosis. *FASEB. J.*, 14:729-739. | 29.Enari, M.; Jen, K.; Cheung, V.; Sakahira, H. and Yokoyama, H. (1998). A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature.*, 391:43-46. |