



Clinical Profile of Patients with Organophosphorus Poisoning in an Intensive Care Unit in a Tertiary Care Hospital

Dr Tilottama Parate

Associate Professor, Department of Medicine Indira Gandhi Government Medical College & Hospital, Nagpur

Dr Ramesh Parate

Associate Professor, Department of Radio-diagnosis, Government Medical College & Hospital, Nagpur

Dr Narayan Bhalerao

Resident, Department of Medicine, Government Medical College & Hospital, Nagpur

*** Dr Nilesh Agrawal**

Assistant Superintendent Government Medical College & Hospital, Nagpur * Corresponding Author

ABSTRACT

Background: POP scale & Serum cholinesterase levels may be effective indicators for mortality in OP poisoning.

Objective: To study the POP scale score and serum cholinesterase levels at initial presentation in OP poisoning and its correlation with need for ventilator support and mortality.

Methodology: 150 eligible participants were recruited and POP scale & Serum cholinesterase levels were assessed.

Observations: Total 32.7% patients died; of which 30 out of 75 belonged to moderate and 19 out of 22 patients belonged to severe group; as per POP scale score. 49 out of 75 in moderate poisoning & all patients in severe poisoning required ventilator support. W.r.t. Serum Cholinesterase level; 20 out of 23 and 29 out of 77 patients died in the severe & moderate poisoning categories respectively. 48 out of 77 in moderate poisoning & all patients in severe poisoning required ventilator support.

Conclusion: Assessment by POP scale and serum Cholinesterase levels is useful in assessing the need for ventilator support and are indicator of mortality in OP poisoning cases.

KEYWORDS : POP scale, serum Cholinesterase level, OP poisoning

INTRODUCTION

Organophosphorus (OP) compounds are commonly used as pesticides¹. Easy availability & widespread use has increased the likelihood of poisoning with these compounds. WHO estimates three million pesticide poisoning episodes annually; of which a minimum of 300,000 die.² OP poisoning is an important indication for emergency admission in hospitals throughout India.³

Mortality in OP poisoning is reported 4-30% in various Indian studies. Respiratory Failure is the commonest complication leading to death: prompt ventilator support may improve survival. With limited availability of resources, all OP poisoning patients are not managed in ICUs in India. Hence clinical features and criteria are important to be identified early, to predict the need for ventilator support.³

RBC cholinesterase level is a sensitive indicator¹, but its estimation is difficult and not commonly available. Serum cholinesterase level, which can be routinely estimated locally, is decreased after OP poisoning.⁴ Peradeniya OP scale (POP scale) hasn't been studied much in Indian scenario. It could be a simple and effective system to determine early the need for ventilation, as has been speculated internationally.⁵

The present study aims to correlate POP scale score and serum cholinesterase levels and the need for ventilation and mortality.

METHODOLOGY

Study design: Prospective observational study.

Study setting: ICU at Tertiary care Hospital.

Study duration: December 1st 2013 to November 30th 2015.

Sample size estimation: Based on Murat study⁶, anticipated mortality in patients of OP poisoning was 27.65%. At absolute preci-

sion-10% & confidence level-99%, minimum sample size required for study was 133. Total 150 participants were studied.

Outcome measures: Death/discharge from hospital.

Participant Selection:

Inclusion criteria –

History of OP poisoning within 24hr of admission.

Presence of signs/symptoms of OP poisoning with decreased serum cholinesterase level.

Exclusion criteria –

- Patient taken treatment elsewhere for current episode.
- Poisoning with other compounds along with organophosphates
- History of chronic liver/pancreatic disease.
- Pregnancy.
- Diabetes.
- History of alcohol consumption and/or drug abuse.

The diagnostic criteria of intermediate syndrome (IMS) included:

- History of acute OP poisoning.
- The presence of clinical manifestation of intermediate syndrome.
- Recovery from the acute cholinergic crisis of intermediate syndrome.⁷

All patients with history of OP poisoning who were admitted to ICU in Department of Medicine were put through the selection criteria. Eligible patients underwent history, clinical examination, biochemical examination. Information was collected through pre-tested proforma. Clinical score (POP scale) was noted on admission and accordingly patient were grouped into mild (POP score 0-3), moderate (POP score 4-7) and severe poisoning (POP score 8-11). Blood samples were tak-

en immediately and sent for estimation of plasma cholinesterase level before doing any intervention. According to cholinesterase activity (Pseudocholinesterase) the OP poisoning were graded as Normal (>50%), Mild (20-50%), Moderate (10-20%) & Severe (<10%). Patients were followed up to death/discharge.

OBSERVATIONS

Majority of cases (56.7%) were between the age group 20 to 40 years, with male predominance (71.3%). Main motive for majority of poison consumption was suicidal (86%), the major route of poisoning being oral ingestion (96.7%). Chi-square analysis for trends showed increasing mortality rate amongst participants with higher quantity of compound consumed, more time lag between consumption of poison and starting of treatment & in those developing IMS; all of the results being statistically significant.

According to POP scale assessment, 53 (35.33%) patients had mild poisoning while 75 (50%) had moderate and 22 (14.67%) patient had severe poisoning. 19 out of 22 severely poisoned patients died, while in moderate category 30 out of 75 patients died. When analysed statistically it was found to be statistically significant (P=0.034) suggestive of more severe POP scale score more is mortality. 49 out of 75 in moderate poisoning & all patients in severe poisoning required ventilator support. Need of ventilation was more in people with high POP scale. Chi-square test for trend revealed highly significant correlation (P<0.001) between POP scale score & the need for mechanical ventilation.


POP Scale	Total Number (%)	Ventilator Support		No Ventilator Support	
		Survival	Mortality	Survival	Mortality
0-3 (Mild)	53 (35.33)	-	-	53	0
4 - 7 (Moderate)	75 (50.0)	19	30	26	0
8 - 11 (Severe)	22 (14.67)	3	19	0	0
Total	150 (100)	22	49	79	0

Categorisation as per Cholinesterase levels revealed 45 (30%) patients in mild, 77 (51.33%) in moderate and 23 (15.33) patients in severe poisoning category. 20 out of 23 patients and 29 out of 77 patients died in the severe & moderate poisoning categories respectively; suggestive of lower Cholinesterase level being directly correlated to mortality, which was statistically significant. 48 out of 77 in moderate poisoning & all patients in severe poisoning required ventilation. Chi-square test for trend applied to compare Cholinesterase level & ventilator support depicted high statistical significance (P<0.001).

Pseudo-cholinesterase Level	Total Number (%)	Ventilator Support		No Ventilator Support	
		Survival	Mortality	Survival	Mortality
< 10 % (Severe)	23 (15.33)	3	20	-	-
10 - 20% (Moderate)	77 (51.33)	19	29	29	-
20 - 50% (Mild)	45 (30)	-	-	45	-
> 50 % (Normal)	5 (3.33)	-	-	5	-
TOTAL	150 (100)	22	49	79	-

Negative correlation was observed between the POP score and Pseudo-cholinesterase level, with statistically significant r-value of 0.7588.

(Fig. 1 comes here)



DISCUSSION

The incidence of OP poisoning was higher in young adults, with almost two third participants being males, as reported in previous studies with similar hypothesis.^{3,8,9}

Suicide being the main motive (86%) for poison consumption has been corroborated by Mood et al¹⁰ who reported it to be 94.3%. It was reported at 67% by Saadeh et al¹¹ & at 68% by Murat et al⁶.

In study carried out by Laudari et al¹¹ increase amount of OP poison intake (>40ml) had significant correlation with increase in the mortality rate (p=0.02). Suliman et al¹² reported higher mortalities (6.3%) in those consuming >30ml than those <30ml (mortality being 2.7%). Findings of both the studies are in-line with our study, in which mean quantity consumed in survival was 17.30 ± 10.73ml & in mortality group was 28.89 ± 9.37ml.

In present study, we observed higher mortality rate amongst participants reporting late to the hospital after OP consumption; which was refuted by Laudari et al¹¹, but supported by statistically significant findings of Kavya et al³ & Patil et al¹³.

As for severity of poisoning, in a study carried out by Kavya et al³, 27% were in mild category, 50.8% were in moderate category & 22% were severely poisoned. In study conducted by Sen et al¹⁴, 29% patients had mild, 45% moderate and 26% had severe poisoning. In study conducted by Nermeen et al¹⁵, 51.7% had mild, 33.3% moderate and 15% had severe poisoning.

In present study, according to POP scale, out of 150 patients, 35.33% were in mild category, 50% in moderate category and 14.67% patients fell in severe category. A total of 32.7% of the patients died, of which 30 out of 75 patients belonged to moderate and 19 out of 22 patients belonged to severe group. So, POP scale correlated, directly & significantly, with mortality. 49 out of 75 in moderate poisoning & all patients in severe poisoning required ventilator support. Need of ventilation was reported to be more in people with high POP scale.

In present study, according to assessment of Pseudo-cholinesterase levels, 30% patients had mild poisoning, 51.3% had moderate and 15.33% patients were severely poisoned. 20 out of 23 patients who were severely poisoned died, while in moderate category 29 out of 77 patients died. Thus, a direct correlation was also observed between decreased serum cholinesterase level and mortality, which was statistically significant.

The current study observed significant negative correlation of POP score and Pseudo-cholinesterase level, with an r-value of -0.7588. This is suggestive of more severe POP scale with lower Pseudo-cholinesterase level.

According to Pseudo-cholinesterase levels, 48 out of 77 in moderate poisoning & all patients in severe poisoning required ventilator support & trend was statistically significant. Need of ventilation was reported to be more in people with low Pseudo-cholinesterase level. In study by Kavya et al³, there was significant correlation between severity of poisoning and the serum cholinesterase at the time of initial presentation of the patients (P<0.001). There was also positive rela-

tionship between POP scoring and lower Pseudocholinesterase level to that of need for ventilation. Incidence of mortality was significantly associated with lower Pseudocholinesterase level and POP Scoring. Similar findings were reported by Shah et al.⁸

CONCLUSIONS

Assessment by POP scale and by Pseudocholinesterase levels at initial presentation in OP poisoning cases is useful in assessing the severity & need for ventilation; at the same time being important indicators of mortality.

REFERENCES

1. Taylor P. Anticholinesterase agents. Goodman and Gilman's The Pharmacological basis of Therapeutics. 11th ed.2006.Pg-176-82.
2. Mancini F, Janice LS, O'Malley M. Reducing incidence of acute pesticide poisoning by educating farmers on integrated pest management in South India. *International journal of occupational and environmental health* 15.2(2009):143-151.
3. Kavya ST, Srinivas V, Chandana, Madhumati R. Clinical Profile of Patients with Organophosphorus poisoning in intensive care unit in tertiary hospital. *International Journal of Clinical cases and investigations*.2012:Vol-4(issue 3),24:31.
4. Rehimani S, Lohani SP, Bhattarai MP. Correlation of serum cholinesterase level, clinical score at presentation and severity of OP poisoning. *J Nepal Med Assoc*.2008;(170):47-52.
5. Senanayake N, de Silva HJ, Karalliedde L. A scale to assess severity in organophosphorus intoxication: POP scale. *Hum ExpToxicol*.1993;12:297-9.
6. Murat S, Muhammed G. Intensive care management of organophosphate insecticide poisoning. *Crit Care*.2001;5:211-5.
7. De Bleecker JL. Intermediate syndrome: Prolonged cholinesterase inhibition. *J Toxicol Clin Toxicol*.1993;31(1):197-199.
8. Shah H et al. "Acute organophosphorus poisoning and clinical admission Score association among patients admitted in emergency ward of tertiary hospital of medical college." *Journal of Pharmaceutical and Biomedical Sciences*.17.17(2012).
9. Weissmann-Brenner A, Aviv-Vidan A and Hourvitz A. Organophosphate poisoning: A multi-hospital survey. *I.M.A.J.* 4:573-576 (2002).
10. M Balali-Mood; K Balali-Mood; F Hosseini Shirazi. *Iranian J Pharmaceutical Res*. 2006,2,79-87.
11. Laudari S, Patowary BS. Analysis of Organophosphorus compound poisoning patients attending CMS-TH, Bharatpur, Nepal. *J Coll Med Sci-Nepal* 2011;7:9-19.
12. Thungs G, Sam KG, Khera K, Pandey S, Sagar SV. Evaluation of organophosphorus poisoning cases in a tertiary care hospital. *JToxEnv Health Sci*;Oct 2010;2(5):73-76.
13. Patil, Virendra C. Clinical Profile and Outcome of Organophosphorus Poisoning at Tertiary Care Centre in Western Maharashtra. *Indian Journal of Forensic Medicine & Toxicology* 6.2 (2012):239.
14. Sen R, Nayak J, Khadanga S. Study of serum cholinesterase, CPK and LDH as prognostic biomarkers in organophosphorus poisoning. *Int J Med Res Rev*, May-June 2014; 2(3):185-189.
15. Nermeen AM, Abdelmonem G. Madbolyet al. Correlation between Serum Creatine Phosphokinase and Severity of Acute Organophosphorus Poisoning. *IOSR Journal of Environmental Science, Toxicology and Food Technology*:May -Jun 2013;4(5):18-29.