



## A Clinico-Pathological Study on Cervical Lymphadenopathy

**Dr.K.LAXMANA  
MURTHY M.S.**

ASSOCIATE PROFESSOR, DEPARTMENT OF GENERAL SURGERY  
KAKATIYA MEDICAL COLLEGE, WARANGAL-506002

**DR K . RAGHU.MS**

ASSISSTANT PROFESSOR, DEPARTMENT OF GENERAL  
SURGERY,KAKATIYA MEDICAL COLLEGE, WARANGAL-506002

**Dr. RAJESHWARI  
KONDABATHINI.,  
M.B.B.S.**

PG IN M.S GENERAL SURGERY,DEPARTMENT OF GENERAL SURGERY  
KAKATIYA MEDICAL COLLEGE, WARANGAL-506002

**Dr. RAJESHWARI  
KONDABATHINI.,  
M.B.B.S**

HNO 8-3-41/A/6/1 1ST FLOOR LALCHI DODDI , OLD BEAT BAZAAR  
WARANGAL TELANGANA -506002

### ABSTRACT

**BACKGROUND AND OBJECTIVE:** The analysis of lymph node enlargement in the neck is not a easy task. It is a challenge for surgeon to assess its clinical behaviour and come to a final diagnosis. These diseases which can be neoplastic also demands correct diagnosis for further management. The study intends to find out systematically the various pathological conditions presenting with enlarged lymph nodes in the neck, also the various modes of clinical presentation and behaviour of these conditions. Relevant investigations have also been studied.

**METHODS:** The study population consisted of patients above 12 years presenting with cervical lymph node enlargement. The proforma which was drafted is used. Patient was examined systematically giving utmost importance to local examination. After making a clinical diagnosis, further relevant investigations were done to confirm the diagnosis. Treatment was instituted appropriately and followed up the patients.

**RESULTS:** Majority of the cases in this study had non-neoplastic causes for cervical lymphadenopathy in which tuberculosis is most common. Male to female ratio of 1.38:1 is noted with most cases between 12 and 30 years. Posterior triangle group of lymph nodes was most commonly affected in tuberculosis. In lymphomas level 2 group of lymphnodes is most commonly involved. Variable results were noted among the groups of lesions, with regard to local characteristics like number, laterality, mobility and involvement of other group of lymph nodes, etc. FNAC by virtue of it being inexpensive,

quick in getting the results and easy to perform, is one of the important and essential diagnostic procedures.

**INTERPRETATION AND CONCLUSION:** Clinical symptoms in cervical lymphadenopathy is of limited significance because clinical behaviour can be highly variable. As cervical lymphadenopathy is an important disease, it always calls for meticulous attention, analysis and treatment. FNAC is found to be a frontline investigation of choice with biopsy and histopathological examination done for confirmation. Most of the non-neoplastic lesions are medically curable with limited role for surgery.

**KEYWORDS :** Cervical lymphadenopathy; FNAC; Histopathological examination

### AIMS & OBJECTIVES

1. To study about the various clinical presentations of cervical lymphadenopathy.
2. To correlate pathological findings with the clinical diagnosis.
3. To study the role of FNAC by correlating with confirmed biopsy report.
4. To study the management, outcome and clinical behaviour of cervical lymph nodes on follow up.

### MATERIALS AND METHODS

#### MATERIALS

The clinical material consists of all inpatients and outpatients of Mahatma Gandhi Memorial Hospital attached to Kakatiya Medical College Warangal. The material consists of patients during the period of December 2013 to October 2015. This study consists of 100 consecutive cases. Diagnosis is made on the basis of histopathological findings.

#### Inclusion Criteria

1. Patients presenting with cervical lymph node enlargement more than 3 weeks duration .
2. Patients more than 12 years of age.

#### Exclusion Criteria

- 1.Acute cervical lymphadenitis

2. Patients less than 12 years of age.

3. Patients where FNAC and/or biopsy of node could not be carried out were excluded.

### METHODS

In this study the data was taken from Mahatma Gandhi Memorial Hospital attached to Kakatiya Medical College Warangal According to proforma detailed history was taken, thorough examination was carried and basic relevant investigations was done in all patients to arrive at a provisional diagnosis.

#### Investigations carried out include

##### a. Routine Investigations

##### Blood

Hb% ,TC,DC , ESR

Liver function tests

##### B. Radiography

a. X-ray neck ,b. X-5ray chest

##### C. Imaging

a. Ultrasonography

b. CT scan , Chest And Abdomen

##### D. Endoscopy

Direct laryngoscopy, Nasopharyngoscopy, Esophagoscopy (triple

endoscopy) Indirect laryngoscopy, bronchoscopy, etc.

E. Skin tests

- in cat scratch disease/tularaemia
- Mantoux for tuberculosis

F. FINE NEEDLE ASPIRATION CYTOLOGY (FNAC)

G. EXCISION BIOPSY OF CERVICAL LYMPH NODE

H. Sputum for AFB

I. Thyroid profile

Investigations like Fine Needle Aspiration Cytology and blood examination were done as a routine. Biopsy was done for all patients. Radiological examination of the chest were done to find primary lesion of lung. Lymph node biopsy specimen was sent to pathologist for expert opinion.

Also ENT opinion, contrast radiological investigation, X-ray endoscopy was carried out in relevant cases.




Figure 1 : Excision biopsy

OBSERVATION AND RESULTS

In the present study 100 cases were selected in the surgery outpatient department and inpatient in surgical wards of Mahatma Gandhi Memorial Hospital attached to Kakatiya Medical College Warangal. From period of December 2013 to October 2015.

Table 1: The number and percentage of non-neoplastic and neoplastic lesions

	Number of cases	Percentage
Non-neoplastic	80	80
Neoplastic	20	20
Total	100	100

Figure 2: The number and percentage of non-neoplastic and neoplastic lesions




Table 2: Histopathological diagnosis in 100 cases

Histopathological diagnosis	Number of cases	Percentage
Tuberculosis	53	53

Reactive lymphadenopathy	27	27
Secondaries	14	14
Hodgkin's lymphoma	1	1
Non-Hodgkin's lymphoma	5	5
Total	100	100

Figure 3: Histopathological diagnosis in 100 cases




Table 3: Sex distribution

Sex	Number of cases	Percentage
Male	58	58
Female	42	42
Total	100	100

Figure 4: Sex distribution




Table 4: Age distribution in both sexes

Age group (years)	Male		Female		Total	
	No.	%	No.	%	No.	%
12 to 20	6	6	8	8	14	14
21 to 30	19	19	13	13	32	32
31 to 40	14	14	8	8	22	22
41 to 50	7	7	6	6	13	13
51 to 60	7	7	3	3	10	10
>60	5	5	4	4	9	9
Total	58	58	42	42	100	100

Figure 5: Age distribution in both sexes




Table 5: Incidence of presenting symptoms

Symptoms	Number of cases
Neck swelling	100
Pain	15
Fever	19
Cough	13
Loss of appetite	12
Loss of weight	17
Difficulty in swallowing	2
Change in voice	1

Figure 6: Incidence of presenting symptoms




Table 6: History of contact with tuberculosis in tubercu-  
losis lymphadenitis cases

Contact with tuberculosis	Number of cases	Percentage
Positive	12	22.6
Negative	41	77.4

Figure 7: History of contact with tuberculosis in tubercu-  
losis lymphadenitis cases




Table 7: Site distribution of tubercular cervical lymphadenitis, reactive lymphadenitis, lymphomas, secondaries

Site	Tubercular cervical	Reactive lymphadenitis	Lymphomas	Secondaries	Total
Level I (submental and submandibular group)	9	10	0	0	19
Level II (upper jugular group)	14	6	3	7	30

Level III (middle jugular group)	7	1	0	4	12
Level IV (lower jugular group)	5	3	1	1	10
Level V (posterior triangle group)	18	7	2	2	29
Level VI (anterior compartment group)	0	0	0	0	0
Total	53	27	6	14	100

Figure 8: Site distribution of tubercular cervical lymphadenitis, reactive lymphadenitis, lymphomas, secondaries




Table 8: Discrete/Matted presentations of lymph nodes in tubercular cervical lymphadenitis

Presentations	Number of cases	Percentage
Matted	20	37.74
Discrete	33	62.26
Total	53	100

Figure 9: Discrete/Matted presentations of lymph nodes in tubercular cervical lymphadenitis




Table 9: Chest X-ray positivity in tubercular cervical lymphadenitis

Chest X-ray	Number of cases	Percentage
Positive	5	9.43
Negative	48	90.57
Total	53	100

Figure 10: Chest X-ray positivity in tubercular cervical lymphadenitis




Table 10: Involvement of other lymph nodes in cervical lymphadenopathy

Lymph node group	Tubercular cervical lymphadenitis		Reactive lymph node		Lymphomas	
	No.	%	No.	%	No.	%
Cervical+Inguinal	2	3.7	1	3.7	1	16.7
Cervical+Axillary	2	3.7	4	14.8	0	0
Cervical+Axillary+Inguinal	0	0	0	0	2	33.3

Figure 11: Involvement of other lymph nodes in cervical lymphadenopathy




Table 11: Main types of lymphomas

Types of lymphomas	Number of cases	Percentage
Non-Hodgkin's lymphoma	5	83.3
Hodgkin's lymphoma	1	16.7
Total	6	100

Figure 12: Main types of lymphomas




Table 12: Distribution of primary in malignant secondaries in neck

Primary site of malignancy	Histopathological pattern	Number of cases
Esophagus	SCC	4
Larynx	SCC	2
Stomach	Adenocarcinoma	2
Thyroid	Papillary carcinoma	2
Unknown	SCC	3
	Adenocarcinoma	1

Figure 13: Distribution of primary in malignant secondaries in neck




Table 13: Sensitivity and specificity of FNAC in diagnosing tuberculous cervical lymphadenitis

FNAC	Number of cases
True positive	40
False positive	0
False negative	13
True negative	47
Total	100

Table 14: Sensitivity and specificity of FNAC in diagnosing reactive lymphadenitis

FNAC	Number of cases
True positive	27
False positive	5
False negative	0
True negative	68

Table 15: Sensitivity and specificity of FNAC in diagnosing secondaries in cervical lymph nodes

FNAC	Number of cases
True positive	13
False positive	0
False negative	1
True negative	86
Total	100

Table 16: Sensitivity and specificity of FNAC in diagnosing lymphoma in cervical lymph node

FNAC	Number of cases
True positive	5
False positive	0
False negative	1
True negative	94
Total	100

Table 17: Treatment and outcome

Diagnosis	No. of cases	Treatment	No. of cases	Outcome		
				Recovered	Not followed	Expired
Reactive lymphadenitis	27	Swelling antibiotics	27	27	0	0
Tubercular cervical lymphadenitis	53	Swelling	48	46	2	0
		Swelling with cold abscess or sinus	5	5	0	0
Secondaries	14	Chemotherapy	0	0	0	0
		Operated	2	2	0	0
		Referred	12	0	11	1
Hodgkin's lymphoma	1	Chemotherapy	1	1	0	0
Non-Hodgkin's lymphoma	5	Referred	0	0	0	0
		Chemotherapy	4	4	0	0
		Referred	1	0	1	0

DISCUSSION

The discussion is mainly based on analysis and observations made regarding presenting symptoms, clinical behaviour, signs, investigations, management and postoperative events in 100 cases of cervical lymph node enlargement attending to Mahathma Gandhi Memorial Hospital attached to Kakatiya Medical College Warangal. during the period of December 2013 to October 2015.

In the present study, which studies 100 cases of cervical lymphadenopathy, 80 were non-neoplastic lesions and 20 were neoplastic lesions.

In the study made by Shafullah and Syed Humayun Shah et al.,<sup>77</sup> the incidence of non-neoplastic and neoplastic lesions were 90.6% and 9.4% respectively.

In the present study, tuberculosis accounted for 53% of cases, 27% turned out to be reactive lymphadenitis. Among the neoplastic lesions, malignant secondaries accounted for 14% while non-Hodgkin's lymphoma and Hodgkin's lymphoma accounted for 5% and 1% respectively. The observation made by Jha BC et al.<sup>78</sup> who studied 94 cases, of which tuberculosis was confirmed in

63.8% cases.

Table 18: Comparison of distribution of different lesions

	Tuberculosis	Reactive lymphadenitis	Sec-ondar-ies	Non-Hodg-kin's lympho-ma	Hodgkin's lympho-ma
Shafullah et al.	69%	17.8%	2.9%	3.4%	3.1%
Jha BC et al. (2001)	63.8%	9.6%	20.7%		
Present study	53%	27%	14%	5%	1%

Sex distribution in cervical lymphadenopathy

Of the 100 cases, 58 cases were males and 42 females. The sex ratio in the present study was 1.38:1 (M:F).

Table 19: Comparative analysis of sex distribution

	Bedi RS et al.	Ammari FF et al.	Dworski	Dandap- at MC et al. <sup>81</sup>	Purohit SD et al.	Present study
M:F ratio	01:01.7	1:02	01:01.4	01:01.2	1.4:1	1.22:1

Most of these studies show female predilection. Few studies like Purohit SD et al. and Tripathy SN et al. are comparable with the present study

History of constitutional symptoms

In the present study, 15% of patients presented with pain, 19% with fever, 15% with cough, 12% with loss of appetite, 17% with loss of weight, 2 patients presented with dysphagia and 1 with change in voice.

Site distribution in cervical lymphadenopathy

This study utilised the Memorial Kettering Hospital Classification of neck lymph nodes from Level I to Level VII.

It was observed that in tuberculosis, level V was most commonly affected (33.9%) followed by Level II (26.4%). In contrast, in secondaries Level II group was most commonly involved (50%) and similarly in lymphomas Level II group was involved.

In the Jha BC et al. series, Level II group was most involved in tuberculosis. The result of this study is comparable to the study made by Baskota DK et al. study, wherein tuberculosis Level V lymph nodes is most commonly involved.

In this present study, 20 cases (37.7%) showed multiple matted lymph nodes in tuberculous lymphadenopathy. Thirty-three cases (72.3%) showed single discrete lymph nodes. Jha BC et al. study showed matted lymph nodes in 38.3% of cases which is comparable with the present study.

Chest X-ray positivity was seen in 9.43% of cases of present study. The studies made by Aggarwal P et al. series showed 28.3% positivity and Jha BC et al. series showed 16% positivity.

In the present study, non-Hodgkin's lesion: Hodgkin's lesion ratio is 5:1. While findings by Peh SC and Shamie et al. had a ratio of 9:1. Raymond Alexandrian study had a ratio of 5.02:1, which has similar results as this present study.

The commonest site of primary in a case of malignant secondary was lungs and pancreas in the studies by Linderman et al. In the present study it was esophagus followed by larynx. In the study by Osama Gaber et al., it was possible to establish primary in 86.7% whereas in the present study it was only 71.5%. In reset of the cases, primary could not be diagnosed because of limited resources of our hospital.

Role of FNAC in cervical lymphadenopathy

In the present study, the sensitivity and specificity of FNAC in detecting various lesions of cervical lymph node are shown in the following table..

Table 20: The sensitivity and specificity of FNAC

Histopathological diagnosis	Sensitivity	Specificity
Tubercular lymphadenitis	75.8%	100%
Reactive lymphadenitis	100%	93.1%
Malignant secondaries	92.8%	100%
Lymphomas	83.3%	100%

The study by Jha BC et al. reported a sensitivity of 92.8% in diagnosing tubercular lymphadenitis. Dandapat MC et al. reported a sensitivity of 83% for tuberculosis. The study by Chao SS, Loh KS et al. showed sensitivity of 88% and specificity of 96% for the same. Similarly Das-



gupta A et al. reported a sensitivity of 84.4% for tuberculosis and 89% for malignant secondary deposits.

Prasad RR et al. studied 2216 cases and noted sensitivity and specificity of 84% and 95% respectively for tubercular lymphadenitis, 97% and 99% for metastatic deposits, 80% and 98% for Hodgkin's disease. 81% and 96% for non-Hodgkin's lymphoma.

In the present study, FNAC sensitivity for tubercular lymphadenitis is low as compared to above studies.

#### Treatment

In the treatment of tubercular lymphadenitis similar findings as in present study was obtained from Jha BC et al., where short course chemotherapy was given with no recurrence.

Another study made by Kaulikama M et al. shows that all patients recovered by combination of anti tuberculous therapy and surgical treatment.

The reactive lymphadenitis were adequately managed with antibiotics and local treatment. The malignant secondaries and lymphomas were staged and treated as per accepted protocols and were referred to higher oncologic centres.

For reactive lymphadenitis, tubercular lymphadenitis medical treatment was instituted.

For secondaries and lymphoma, which needs radiotherapy, chemotherapy and expert oncologic surgeries, patients were referred to MNJ Cancer Hospital Hyderabad.

For all patients, necessary advice given and were asked to attend the surgical outpatient department for follow-up.

#### CONCLUSION

The clinical material consists of patient consecutively selected with history of cervical lymphadenopathy, who came to surgical OPD and who were admitted in wards of Mahathma Gandhi Memorial Hospital attached to Kakatiya Medical College Warangal during the period of December 2013 to October 2015

One hundred consecutive cases were selected and they were personally studied in the present study.

Of the 100 cases, tuberculous lymphadenopathy had the maximum incidence of 53% followed in reactive lymphadenitis (27%), secondaries (14%) and lymphomas (6%).

Overall age at presentation was maximum between 12 years and 30 years followed by 31-40 years.

In investigations, Fine Needle Aspiration Cytology was found to be accurate with 75.5% accuracy for the diagnosis of tuberculosis. Few patients were diagnosed as non-specific lymphadenopathy which were later confirmed by biopsy to have either tuberculosis or reactive lymphadenitis.

In metastatic lymph node, method of diagnosis was Fine Needle Aspiration Cytology and two patients were treated with surgery. One patient expired before referral. Rest 11 cases were referred to oncologic centre and they did not come for follow-up.

Lymphomas were diagnosed by Fine Needle Aspiration Cytology and confirmed with excision biopsy.

Hodgkin's lymphomas was treated with chemotherapy and was followed up regularly till the study concluded. No mortality noted.

Among 5 non-Hodgkin's lymphoma cases, 4 cases were treated with chemotherapy and they were followed up regularly all the study concluded. No mortality noted during the study 1 case got referred to oncologic centre.

In this present study, fine needle aspiration cytology was found

to be reliable and cheapest method of diagnosis without any significant morbidity and with good patient compliance.

#### REFERENCES

1. Shafullah Syed H. Tuberculous lymphadenitis on Afghan refugees. *J Pathol* 1999;187:28-38.
2. Jha BCA, Nagarkar NM, Gupta R, Sighal S. Cervical tubercular lymphadenopathy changing clinical patterns and concepts in management. *Postgraduate Med J* 2001 Mar;77(905):185-7.
3. Ammari FF, Bani Hani AH, Gharibeh KI. Tuberculosis of lymph glands of neck; a limited role for surgery. *Otolaryngeal. Head Neck Surgery* 2003 Apr; 128(4):576-80.
4. Dworski I. Tuberculosis of cervical lymph nodes. *Plunice Bolesti* 1989 Jul-Dec; 41(3-4):169-71.
5. Dandapat MC, Mishra BM, Dash SP, Kar PK. Peripheral lymph node tuberculosis: a review of 80 cases. *Br J Surg* 1990 Aug;77(8):911-2.
6. Baskosa DK, Prasad R Kumar, Sinha B, Amatya RC. Distribution of lymph nodes in the neck in cases of cervical lymphadenitis. *Acta Otolaryngeal* 2004 Nov;124(9):1095-8.
7. Osama G, Peter R, Charles E, Joseph J. Metastatic malignant disease of unknown origin. *Am J Surg Pathol* 145:493-7.
8. Dasgupta A, Ghosh RN, Poddar AK. FNAC of cervical lymph nodes with special reference to tuberculosis. *J Indian Medical Assoc* 1994 Feb;92(2):44-
9. Prasad RR, Narasimha R, Sankaran V. FNAC in diagnosis of superficial lymphadenopathy: an analysis of 2418 cases. *Diag Cytopathol* 1996 Dec; 15(5):382-6