



Study The Effect of Socio-Economic Status and Water Sanitation on Common Water Born Diseases

Dr Satendra Singh Rajput

MD Pediatrics , PGMO(SNCU) KRH Hospital Gwalior

ABSTRACT

Introduction; Water born diseases are caused by pathogenic microorganism that most commonly are transmitted in contaminated fresh water. Various forms of waterborne diarrheal diseases, probably are the most prominent examples, and affect mainly children in developing countries. **Method;** This study was designed to see the effect of water sanitation and socio-economic status on common feco-oral transmitted diseases. A total of 116 children, fulfilling inclusion criteria were recruited in the study after taking written informed consent from parents or guardians of children. Detailed data including general information, age, sex, environment, socio-economic status, source of water supply, sewage disposal and any method of water treatment at home were taken. History of diarrheal episode in last month of enrollment were also taken to find out incidence of diarrhea. **Results;** Results showed significant higher incidence of diarrhea and hepatitis A among children with lower socio-economic classes and poor sanitation practices. **Conclusion:** Improving access to safe drinking water and adequate sanitation, as well as promoting good hygiene, are key components in preventing diarrhea and hepatitis A.

KEYWORDS : Diarrhea, Socio-economic class , water born diseases

INTRODUCTION

Water born diseases are caused by pathogenic microorganism that most commonly are transmitted in contaminated fresh water. Various forms of waterborne diarrheal diseases probably are the most prominent examples, and affect mainly children in developing countries¹.

Diarrheal diseases and hepatitis are a major cause of hospitalization & child's death globally. Together they account for approximately one in six deaths among children younger than five year's (WHO). Major risk for diarrhea and hepatitis includes environment contamination & increased exposure to enter pathogens. Viral hepatitis continues to be a major public health problem in India as well as in other part of the world.² The hepatitis A virus is an enterically transmitted virus and is the major cause of acute viral hepatitis in children.³

This study was designed to see the effect of water sanitation on common feco-oral transmitted diseases that is diarrhea and hepatitis.

MATERIAL AND METHOD:

This was a cross- sectional and observational study conducted in Out-patient department (OPD), Department of pediatrics, at a Medical College hospital, Gwalior with due approval from ethical committee for a period of one year (August 2012 to September 2013). Participants were children belonging to Gwalior district from both urban and rural areas, between 1 to 5 years, attending pediatric outpatient department with minor problems. Children who received HAV vaccination were excluded from study

A total of 116 children, fulfilling inclusion criteria were recruited in the study, detailed data including age, sex, environment, source of water supply, sewage disposal and any method of water treatment at home were taken. History of diarrheal episode in last month of enrollment were also taken to find out incidence of diarrhea.

RESULTS

Table-1 ; Socioeconomic status distribution in study population and its relation to HAV seropositivity and incidence of diarrhea

Socioeconomic class	Total no.	Anti-HAV positive		Incidence of Diarrhea	
		No.(%)	P value	No.(%)	P value
Upper [I & II] Lower [III,IV & V]	22 94	12(54.5%) 82(87.2%)	0.00075	3(13.6%) 32(34.04%)	<.005

Among a total of 22 children belonging to upper socioeconomic class 12 (54.50%) were seropositive for anti-HAV antibody compared to 82 (87.2%) out of 94 children belonging to lower socioeconomic class. In this study difference of positivity for anti-HAV antibody with children having a lower (class III,IV,V) SEC and upper (class I,II) SEC was sta-

tistically significant with P value 0.00075.

Among a total of 22 children belonging to upper socioeconomic class 3 (13.6%) were positive for diarrheal episode compared to 32 (34.04%) out of 94 children belonging to lower socioeconomic class, and this difference was significant.

Table -2; Correlation of method of water treatment at home with seropositivity of HAV ab and incidence of diarrhea

Method of water treatment	No. of cases 116 (100%)	Anti HAV Ab +ve No. (%)	p value	Incidence of diarrhoea in last month (n=35)	P value
No treatment at home	98 (84.48%)	84 (85.71%)	0.0036	33(33.6%)	<0.005
Boiling	04 (3.44%)	02(50.00%)		0	
Chlorine treatment	00 (00%)	00 (00%)		0	
Mechanical filtration	06 (5.17%)	05 (83.33%)		1(16.6%)	
Filtration + UV treatment	8 (6.89%)	03 (37.50%)		1(12.5%)	

Among total 116 children, 98 were not having any treatment of water at home out of them 84 (85.71%) were seropositive for anti-HAV antibody. In our study difference of seropositivity for anti-HAV antibody with children having any method of water treatment at home and without any treatment was statistically significant with p value 0.0036.

We also take history of diarrheal episode in last month of enrollment among study cases and we found that out of 98 study subject, those were not treating drinking water, 33 had history of diarrhea while only 2 study case had diarrheal episode in last month of enrollment among 18 children those were using any method of water treatment at home. This difference was statistically significant.

Table - 3 ; Correlation of source of water supply with seropositivity of anti-HAV antibodies and incidence of diarrhea

Source of water supply	Total no.	Anti-HAV positive			Incidence of Diarrhea		
		No.	%	P value	No.	%	P value
Municipal bore well	54	49	88.88	0.026	19	(35.1%)	0.0034
Municipal Tighra dam supply	24	19	79.16		11	(45.8%)	
Personal bore well in home	38	26	68.16		5	(13.15%)	

Out of 54 children with municipal bore well water supply, 49 (88.88%) were seropositive for total anti-HAV antibody and out of 24 children with municipal Tighra dam supply 19 (79.16%) were seropositive compared to 26 (68.16%) out of 38 children with an inside house source of water. The difference of positivity for total anti-HAV antibody with children having outside house water supply either from municipal bore well or dam and inside house water supply was statistically significant with P value 0.026.

Similarly incidence of diarrhea is more with children having outside house water supply either from municipal bore well or dam then inside house water supply, which was statistically significant with P value 0.0034.

DISCUSSION:

Several previous studies showed a clear inverse correlation between exposure to HAV and socioeconomic level. It is well known that HAV infection is strongly correlated with poverty and inadequate sanitation. In present study difference of positivity for hepatitis A IgM/IgG antibody with children having a lower [class I, II, III] class and higher [class IV, V] class was statistically significant with p value 0.00075. This result was in the line of study by C P Rath et al⁵, Dhawan et al⁴. showing a statistically significant difference. Similar to this study, Some studies have shown that the association between socio-economic factors, such as poor housing, crowded condition, low income and higher rate of diarrhea was statistically significant. The results are the reflection of the fact that higher socio-economic groups have better access and affordability to food hygiene and water sanitation.

In this study seropositivity was significantly lower in children living in family having source of water in their own house by personal bore well (68.16) in compared to those children with family having pipe line supply either from public bore well (88.88%) or dam supply(79.16%) suggesting probably a contamination while carrying water from outside in to the house. Similar results were observed by C P Rath et al⁵. I .J. Salama et al⁷ concluded that the risk of infection with HAV was 3 times higher among children using a public water supply compared to those with piped water inside the home.

As diarrhea is acquired via contaminated water and foods, water-related factors are very important determinants of diarrhea occurrence. Increasing distance from water sources, poor storage of drinking water^{10, 12, 14, 15} (e.g. obtaining water from storage containers by dipping, no drinking water storage facility), use of unsafe water sources (such as rivers, pools, dams, lakes, streams, wells and other surface water sources)^{11, 13, 16, 17} water storage in wide mouthed containers; low per capita water used¹⁷; have been found to be risk factors for more diarrhea occurrence among children less than five. The present study also showed higher incidence of diarrhea among children using a public water supply compared to those with piped water inside the home.

In this study difference of seropositivity for anti-HAV antibody with children having any method of water treatment at home and without any treatment was statistically significant. This is the only study of its type in which impact of water treatment at home on seropositivity of anti HAV IgM/IgG antibody and incidence of diarrhea was observed.

CONCLUSIONS

Improving access to safe drinking water and adequate sanitation, as well as promoting good hygiene, are key components in preventing diarrhea and hepatitis A and other waterborn diseases as well.

REFERENCES:

1. "Burden of disease and cost-effectiveness estimates". World Health Organization. Retrieved April 5, 2014.
2. Jensen PK, Ensink JH, Jayasinghe G, van der Hoek W, Cairncross S, Dalsgaard A. Domestic transmission routes of pathogens: the problem of in-house contamination of drinking water during storage in developing countries. *Trop Med Int Health*; 2002; 7(7): 604–9.
3. Gust ID, Feinstone SM, Hepatitis A. In: Popper H, Schaffner F, eds. *Progress in liver diseases*. Philadelphia, WB Saunders Co., 1998:371-8.
4. Dhawan PS,Shah SS,Avares JF,Kher A,Shankaran,Kandoth PW,Seth PN,Kamath H,Kamath A,KOppikar GV,Karo RH.Seroprevalence of hepatitis A virus in Mumbai and immunogenicity and safety of hepatitis A vaccine. *Indian J Gastroenterology*. 1998 Jan; 17(1):16-8.
5. CP Rath, Arvind Akki, SV Patil and SS Kalyanshettar. 'Seroprevalence of hepatitis A virus antibody in Bijapur, Karnataka' *Journal of Indian Pediatrics* 2011; 48:71-73.
6. Jindal M, Rana SS, Gupta RK, Das K, Kar Pserological study of hepatitis A virus infection amongst the students of a medical college in Delhi and evaluation of the need of vaccination. *Indian J Med Res* 2002; 115: 1-4.
7. I.I.Salama, S.M. Samy, F.A. Shaaban, A.I. Hassanin and L.A.Abou Ismail. 'Seroprevalence of hepatitis A among children of different socioeconomic status in Cairo. *La Revue de Sante de la Mediterranee orientale*.20007;13:1256-64.
8. International Institute for Population Sciences (IIPS) and Macro International. 2007. *National Family Health Survey (NFHS-3), 2005–06: India: Volume I. Mumbai: IIPS.*
9. Yassin K. Morbidity and risk factors of diarrheal disease among under-five children in rural upper Egypt. *J Trop Pediatr*; 2000; 46 (5): 282-7.
10. Teklemariam S, Getaneh T, Bekele F. Environmental determinants of diarrheal morbidity in under-five children, Keffa-Sheka zone, south west Ethiopia. *Ethiop Med J*; 2000; 38(1):27-34.
11. Karim AS, Akhter S, Rahman MA, Nazir MF. Risk factors of persistent diarrhea in children below five years of age. *Indian J Gastroenterol*. 2001; 20(2):59-61.
12. Jinadu MK, Olusi SO, Agun JI, Fabiyo AK. Childhood diarrhoea in rural Nigeria. I. Studies on prevalence, mortality and socio-environmental factors. *J Diarrhoeal Dis Res*.1991;(4):323-7.
13. Marjatta BS. Water supply and diarrhea in East African community. A casecontrol study on the quality of water supplies and the occurrence of diarrhea among small children in a rural area of Western Kenya. University of Oulu Printing Center, 1994: 37-57.
14. Molbak K, Jensen H, Ingholt L, Aaby P. Risk factors for diarrheal disease incidence in early childhood: a community cohort study from Guinea-Bissau. *Am J Epidemiol*. 1997;146(3):273-82.
15. Wijewardene K, Fonseka P, Wijayasiri WA. Risk factors contributing to acute diarrhoeal disease in children below five years. *Ceylon Med J*, 1992; 37(4):116-9.
16. Tumwine JK, Thompson J, Katua-Katua M, Mujuwajuzi M, Johnstone N, Porras I. Diarrhoea and effects of different water sources, sanitation and hygiene behaviour in East Africa. *Trop Med Int Health*. 2002; 7(9):750-6.
17. Brooks JT et al. Epidemiology of sporadic bloody diarrhea in rural western Kenya. *Am J Trop Med Hyg*. 2003; 68 (6): 671–7.