



HISTOMORPHOMETRIC ANALYSIS OF UMBILICAL CORD IN GESTATIONAL DIABETES MELLITUS

Amit Jain	Associate Professor, Department of Anatomy, R.D.Gardi Medical College, Ujjain
Rajiv Ranjan	Assistant Professor, Department of Anatomy, RIMS, Ranchi
Rashmi Bhujade	Assistant Professor, Department of community medicine, R.D.Gardi Medical College, Ujjain

ABSTRACT **Aim:** The aim of present study was to determine the adverse effects of gestational diabetes on the histomorphological features of the umbilical cord.

Place of study: Department of Anatomy, R.D.Gardi Medical College, Ujjain

Materials and Methods: Total 100 umbilical cords with the placenta were examined. The relevant data were collected from patients delivered in C.R. Gardi hospital, Ujjain within 36th to 40th weeks of gestation. Control cords (n = 50) were from healthy women whose pregnancies were not complicated by any major disease. GDM cords (n = 50) were from mothers with gestational diabetes mellitus (GDM group). All the placentas with cords were treated identically in a standard manner. All placentas were fixed in 10% buffered formalin for 2 weeks. The diameter of the umbilical cord was measured in millimeters and the Circumference, the Cross-Sectional Area of the umbilical cord and Cross-Sectional Area of the Wharton's jelly were measured in millimeters and sq. millimeters respectively.

Result: The study demonstrated that in GDM group the mean diameter, the mean circumference, the mean CSA of the umbilical cord and the mean CSA of the Wharton's jelly were larger in the GDM group than the Control group. The difference was significant in case of the mean diameter while it did not reach to the significant level for the other parameters.

KEYWORDS : Umbilical cord, Diabetes, Histomorphometry

Introduction:

The umbilical cord normally contains three vessels, two arteries and one vein, surrounded by a connective tissue known as "Wharton's jelly" 1. At term, the mean length of the umbilical cord is 55 cm. A cord is considered short if it is less than 35 cm in length (lower 6th percentile), and long if measures more than 80 cm (upper 6th percentile). The mean umbilical cord circumference at 40 weeks is 3.6 cm (range 2.6 to 6.0 cm). The 90th percentile for the area of the umbilical cord at term is 1.3 cm square. It is two centimeters in diameter. The umbilical cord has an organ-like property. For the better interpretation of feto-maternal functional relationship, understanding the morphology of the umbilical cord is important. The umbilical cord diameter, circumference and cross-sectional area (CSA) are important measurements. Due to the increase in its Wharton jelly content, the umbilical cord is larger in diameter in the fetuses of mothers with gestational diabetes than in the normal population. Also the Wharton's jelly shows large empty spaces 2,3.

Materials and Method:

The study was conducted in the Department of Anatomy, R.D.Gardi Medical College, Ujjain from Dec'2015 to Jan'2017. The study included 100 subjects. They were divided into two groups. Group I consisted of umbilical cords obtained from normal non-diabetic pregnant women (n=50). Group II included the umbilical cord obtained from mothers with gestational diabetes mellitus; the GDM group.(n=50) The umbilical cord with the placenta was collected from the C. R. Gardi hospital within 36 to 40th weeks of gestation. The vernier calipers were used to measure the mean diameter of the umbilical cords. [(Maximum diameter + Minimum diameter)/2] in millimeters. Measurement of the cross-sectional area of umbilical cord was taken on a plane adjacent to the insertion of the cord into fetal abdomen. Circumference and Cross-Sectional Area of the umbilical cord were measured in mm and mm² respectively using Adobe photoshop CC software. The cross-sectional areas of umbilical cord vessels were measured in mm² and the values were subtracted from the total cord cross-sectional area to assess the cross-sectional area of the Wharton jelly 4. The results were analyzed by applying Student t-test and Pearson's correlation.

Results:

S. no.	Variable	Control n = 50	GDM n = 50	P value
1	Diameter of the umbilical cord (mm) Range Mean ± SD	8.6 – 13.7 11.5 ± 1.93	9.4 – 17.10 12.9 ± 2.95	(P= 0.007) Significant
2	Circumference of the umbilical cord (mm) Range Mean ± SD	27.0 – 43.01 37.3 ± 6.91	29.51 – 52.59 40.03 ± 8.03	(P= 0.118) Not Significant
3	CSA of the umbilical cord (sq. mm) Range Mean ± SD	58.05 – 147.33 86.54 ± 27.26	69.36 – 229.54 103.07 ± 40.44	(P= 0.083) Not Significant
4	CSA of the Wharton's jelly (sq. mm) Range Mean ± SD	40.06 – 136.66 74.03 ± 29.33	45.38 – 170.39 90.66 ± 39.31	(P= 0.117) Not Significant

Results with P < 0.05 were considered significant.

Correlation Co-efficient:

It is a measure of the strength of linear association between two variables. Correlation will always between -1.0 and +1.0. If the correlation is positive, we have a positive relationship. If it is negative, the relationship is negative. Results with P < 0.05 are significant.

Table - 2

Strength of the linear relationship between the mean diameter and mean circumference of the umbilical cord in the study Groups

Correlation variables	Group	Correlation coefficient (r)	Significance
The mean diameter and the mean circumference of the umbilical cord	GDM	0.663	Significant
	Control	0.814	Significant

Table - 3

Strength of the linear relationship between the mean cross-sectional area of the umbilical cord and the Wharton's jelly in the study groups

Correlation variables	Group	Correlation coefficient (r)	Significance
The mean cross-sectional area of the umbilical cord and the Wharton's jelly	GDM	0.986	Significant
	Control	0.976	Significant

Discussion:

In the present study, the GDM group showed significantly higher values of mean diameter of the umbilical cord than the Control group ($P < 0.05$). Weissman et al. also reported the same that the umbilical cord was also significantly larger in the fetus of mothers with gestational diabetes than that of normal mothers.² Increase in the width was because of increase in the Wharton's jelly. The mean umbilical cord circumference at 40 weeks is 3.6 cm (range 2.6 to 6.0 cm)⁵. In our study, the mean circumference of the umbilical cord was larger in the GDM group than in the Control group but statistically it was not significant ($P > 0.05$).

The 90th percentile for the area of the umbilical cord at term is 1.3 cm square⁶. In our study, the mean cross-sectional area of the umbilical cord was larger in the GDM group than the Control group but statistically it was non-significant ($P > 0.05$). In our study, within the GDM and in the Control group, we found a significant positive correlation between the mean diameter ($r = 0.663$, $P < 0.001$) and mean circumference of the umbilical cord ($r = 0.814$, $P < 0.001$). We also found a significant positive correlation between mean cross-sectional area of the umbilical cord ($r = 0.986$, $P < 0.001$) and the mean of the Wharton's jelly ($r = 0.976$, $P < 0.001$).

In this study we found that in the GDM group, the mean cross-sectional area of the Wharton's jelly was much greater than in the Control group but the difference was not significant statistically. The reason for the increase in cross sectional area of the Wharton's jelly was due to presence of many empty spaces within it which could be due to degeneration process involving the connective tissue⁹.

References:

1. Singh I, Pal GP. Human Embryology. 7th ed. Chennai: Macmillan India Limited; 2001. 54.
2. Weissman A, Jakobi P. Sonographic measurements of the umbilical cord in pregnancies complicated by gestational diabetes. *J Ultrasound Med* 1997; 16:691-4.
3. Gill P, Jarjoura D. Wharton's jelly in the umbilical cord. A study of its quantitative variations and clinical correlates. *J Reprod Med* 1993; 38:611-4.
4. Togni FA, Araujo Júnior E, Vasques FA, Moron AF, Torloni MR, Nardoza LM. The cross-sectional area of umbilical cord components in normal pregnancy. *Int J Gynaecol Obstet* 2007; 96:156-61.
5. Dooley SL, Lamb R, Helseth DL Jr. Umbilical cord circumference as a measure of Wharton's jelly: Clinical correlates. Abstract 32, Sixth Annual Meeting of the Society of Perinatal Obstetricians. San Antonio, Texas 1986.
6. Ghezzi F, Raio L, Di Naro E, Franchi M, Balestreri D, D'Addario V. Nomogram of Wharton's jelly as depicted in the sonographic cross section of the umbilical cord. *Ultrasound Obstet Gynecol* 2001; 18: 121-5.
7. Patel D, Dawson M, Kalyanam P, Lungus E, Weiss H, Flaherty E, Nora EG Jr. Umbilical cord circumference at birth. *Am J Dis Child* 1989; 143:638-639.
8. Manuel Vázquez Blanco, Hilda Ruda Vegab, Roberto A. Histopathology and histomorphometry of umbilical cord blood vessels. Findings in normal and high risk pregnancies: Volume 5, Issue 2, June 2011, Pages 50-57
9. A. Cromi, F. Ghezzi, E. Di Naro, G. Siesto, V. Bergamini, L. Raio. Large cross-sectional area of the umbilical cord as a predictor of fetal macrosomia. *Ultrasound Obstet Gynecol* 2007; 30:861-866
10. Predanic M. Sonographic Assessment of the Umbilical Cord. *Donald School J Ultrasound Obstet Gynecol*, 2009;3(2):48-57