

Dr Priyanka Singh

Assistant Professor, Department of Pathology, Narayan Medical College and Hospital, Sasaram, Jamuhar, Bihar-(Corresponding Author)

ABSTRACT There has been increasing interest in the evaluation of essential trace elements present in different concentrations in the human body fluids and their correlation to the human health. Trace elements play important roles in a number of body functions. Our aim was the relationship between concentration of zinc in seminal plasma and semen quality. The results revealed that serum zinc levels in infertile men group were lower than in the fertile men group, the zinc has positive effect on sperm count and morphology. The fact that zinc plays a positive role in influencing the process of spermatogenesis cannot be ruled out. Varicocele decreases the concentration of zinc and decreases the availability of oxygen for live sperm.

**KEYWORDS** : Male Infertility, Seminal Plasma and Sperm Count.

# Introduction:

Zinc in human semen seems to play an important role in the physiology of spermatozoa (processes of fertility, reproduction and sexual maturation). It is secreted by the prostate in small vesicles called protasomes. Zinc (Zn) in seminal plasma stabilizes the cell membrane and nuclear chromatin of spermatozoa. It may also have an antibacterial function.<sup>1</sup> Many recent studies have indicated an increasing prevalence of various abnormalities of the reproductive system in human males.<sup>2</sup> Male infertility is as the inability to conceive a baby after one year of unprotected sex in the absence of female causes. Most causes of male infertility reflect an abnormal sperm count or quality. Although it only takes one sperm to fertilize an egg, in an average ejaculate a man will eject nearly 200 million sperm. However, because of the natural barriers in the female reproductive tract only about 40 sperm will ever reach the vicinity of an egg. There is a strong correlation between the number of sperm in an ejaculate and fertility. In about 90% of the cases of a low sperm count, the reason is deficient sperm production. Unfortunately, in about 90% of cases, the cause for the decreased sperm formation cannot be identified. Other causes of male infertility may include ductal obstruction, ejaculatory dysfunctions, and infections or disorders of the accessory glands<sup>3</sup>. Prostate gland is the largest accessory gland of the male reproductive tract and there is a good deal of research being conducted on the prostate. Yet it remains one of the least understood structures in the body. One area of study concentrates on the major function of the gland, the production and the secretion of minerals that play an important role in prostate and the reproductive system functions. Zinc is a component of semen and it is though that at least 1 mg of zinc is excreted by an ejaculum. Zinc plays an important role in the processes of fertility, reproduction and sexual maturation.<sup>4</sup> At a biochemical level, zinc's primary role as a component of zinc metalloenzymes, many enzymes of the cell nucleus involved with genetic information transfer and cellular replication are metallonezymes of zinc. Zinc also participates in RNA metabolism, prostatic fluid and ocular tissues contain the highest levels of zinc in the body.<sup>5</sup>Our aim was to be clarification of the interrelationship of serum and semen zinc levels, in both men with a history of infertility and men with no history of infertility. Also, investigate the possible role of seminal zinc in the regulation of human spermatozoa functions.

## **Material and Methods:**

This present study was conducted in Department of Pathology, Narayan Medical College and Hospital, Sasaram, Jamuhar, Bihar, India during the period from July 2016 to January 2017. Bihari men in the age range of 20 to 45 years with no apparent chronic or acute disease were selected for the study, which were categorized in two groups:

1. Group A: 35 men of known fertility and

2. Group B: 35 infertile men were studied.

The infertile men were unable to be fathers of children for at least two and half (2.5) year; their wives were considered fertile based on extensive gynaecologic evaluations. At the first interview for the infertile men, a complete history was obtained, especially with respect to any history of sexual dysfunction. Physical examination was performed to exclude patients with chromosomal abnormalities. None of the subjects were on drug treatment or on especial diet and none of the subjects used vitamin supplements.

Zinc estimation: Seminal plasma was diluted 100 times with triple distilled water. Estimation of zinc in diluted seminal plasma was carried out using flame atomic absorption spectrophotometer. (Perkin Elmer Double beam, USA). Lamp used for Zn: Electrode Less Discharge Lamp was used and 213.9 mm wavelength was used for Zn.

Statistical analysis was done using SPSS-17. The differences between means of more than two groups were tested by performing ANOVA. The Pearson's correlations (r, p) and regression equation were calculated between each of the measured parameters to assess the shape of relationship based on the highest (r) value obtained.

## Results and Discussion:

This present study shows the demographic characteristics of the fertile and infertile men. The mean±standard deviation and range of semen quality counts, motility, and morphology, serum reproductive hormones LH, FSH, Testosterone and serum zinc and semen zinc are listed in table-I. The results revealed that serum zinc levels in infertile men group were lower than in the fertile men group (P<0.05). In addition, seminal plasma zinc levels were lower in the infertile men group than in the fertile men group (P<0.05). Table-II shows, there was no significant correlation between serum zinc and the parameters of semen quality in both groups. Moreover, no significant correlation between semen zinc and the parameters of semen quality in both groups.

Table-I: Evaluation of semen quality, hormones, and Zinc concentrations in fertile and infertile men groups:

Parameters	Group A (N=35)		Group B (	P-	
	Mean±SD	Range	Mean±SD	Range	value
Count (60- 150x106/ml)	94.05±16. 02	57-154	38.3±24.0 3	0-98	<0.05
Motility (>60%)	86.13±4.8 2	55-95	16.5±10.2	0-60	<0.05

#### VOLUME-6, ISSUE-9, SEPTEMBER-2017 • ISSN No 2277 - 8160

Morphology	84.21±9.3	70-90	65.04±20.	0-90	< 0.05
(>70%)			6		
LH (1-5mlu/ml)	4.02±1.24	1.5-5.0	2.3±1.02	0.9-23	*
FSH (1-	4.5±2.7	2.8-9.8	5.8±3.0	0.9-34.5	*
9mlu/ml)					
Testosterone	26.02±4.7	15.8-33.0	14.0±3.21	7.0-44.0	< 0.05
(8.2-34.6					
nmol/L)					
Serum Zn (70-	117.02±1.	90.6-162	86.7±21.5	36-147.4	< 0.05
120µg/dl)	32				
Seminal	3.2±0.21	2.5-3.8	1.5±0.2	0.39-2.7	<0.05
plasma Zn(1.2-					
3.9nmol/L)					

NOTE: (Statistically Significant (P<0.05) \* NS (Not Significant)

The present data are in agreement with those of Bonde et al<sup>6</sup>, who reported that not sperm concentration but sperm quality determines fertilizing capacity of spermatozoa. Although, the most cases of male infertility are nonendocrine in origin. However, routine evaluation of hormonal parameter is not warranted unless sperm density is extremely low or there is clinical suspicion of an endocrinopathy. A scrotal varicocele is the most common causative finding in infertile men.<sup>7</sup> To explain the abnormalities in spermatogenesis with varicocele, the most point have been proposed was the abnormal blood flow can interfere with testosterone production, which in turn can interfere with sperm production and this in agreement with our results that about 25% of infertile men had varicocele. We did not find any difference in the our data and in those previously published in the literature for that the serum zinc and semen zinc levels were significantly lower in infertile patients than fertile males<sup>8</sup>, in contrast to other reports, that unable to find a significant difference in serum and semen zinc levels between fertile and infertile men<sup>9,1</sup>

## Table-II: The correlation coefficient (r) between Zinc concentration and parameters of semen quality for both groups:

Groups	variables	Count	Motility	Morphology
Group A	Serum Zn	-0.089	-0.271	-0.062
(N=35)	Seminal Plasma Zn	0.049	0.12	-0.349
Group B	Serum Zn	-0.005	-0.023	-0.460
(N=35)	Seminal Plasma Zn	-0.032	-0.168	0.080

The lack of correlation between zinc concentration and semen quality found in our study suggests that biochemical complexity of seminal fluid attempts to perform such simple correlations between seminal plasma component and andrological parameters are likely to produce inconsistent results. These effects include no significant correlation between the total amount of zinc per ejaculate and sperm quality<sup>11</sup> with no statistically significant correlation between zinc concentration and the motile sperm concentration.<sup>12</sup> However, our results are in contrast to other studies that showed zinc-related decrease in human semen quality.<sup>13,14</sup> Most previous results of colorimetric methods and the present method found that the results obtained using the proposed method were not statistically different from those obtained by atomic absorption spectrophotometry.<sup>15,16</sup>

The high level of zinc found in semen is due primarily to the secretions of the prostate gland and reflects prostatic stores. Serum zinc may be a reasonable indicator of zinc status. The lack of correlation between serum zinc and semen zinc found in our study suggests that mild zinc deficiency may lower serum zinc while the larger prostate zinc stores remain unaffected. However, we obtained 25% had varicocele, which are enlargement of the internal spermatic veins that drain the testes. The significant decrease of the zinc in seminal plasma of varicocele men were significantly lower than in the normal subjects. This decrease indicated an impairment of the prostatic function or secretion due to decreases the availability of oxygen and nutrient required for sperm live. The

Krebs's cycle provides a major source of adenosine triphosphate energy and produce molecules that are starting point for a number of vital metabolic pathways for the cell, citrate that been chelated with zinc and their evidence that zinc required for oxygen consumption by sperm. Varicocele decreases the concentration of zinc and decreases the availability of oxygen for live sperm.

### **Conclusion:**

These findings suggest that the zinc has positive effect on sperm count and morphology. The fact that zinc plays a positive role in influencing the process of spermatogenesis cannot be ruled out. Varicocele decreases the concentration of zinc and decreases the availability of oxygen for live sperm. Further studies to elucidate the significance of zinc and other factors presents in seminal plasma in different types of causes of male infertility for the functional properties of human spermatozoa appear to be of importance since such studies may give hints to new ways of regulating male fertility.

#### **References:**

- Lin YC, Chang TC, Tseng YJ et al. Seminal plasma zinc levels and sperm motion characteristics in infertile samples. Chang Gung Med J 2000;23: 260-6.
- Bromuich, P., Cohen, J., Stewart, I., and Walker, A.: decline in sperm counts: An Artifact of Changed Reference Range of "Normal"? BMJ, 1994; 309: 19-22.
- Wyngaarden, J.B., Smith, L.H., and Bennett, J.C.: Endocrinolgic Diseases Unique to Men, Cecil Textbook of medicine, 20th edition. WB Saunders, Philadelphia, PA, 1996; Vol. 2: 1325-45.
- Shils, Maurice, E., And Young, Vernon, R.: Essential Trace Elements, Modern Nutrition in Health and Disease, 7th edition. Philadelphia, lea&febiger, 1988; p.p. 375-92.
  Chester's, J.K.: Metabolism and biochemistry of zinc, Clinical, Biochemical and
- Chester's, J.K.: Metabolism and biochemistry of zinc, Clinical, Biochemical and Nutrition Aspects of Trace Elements, New York: Alan liss, 1982; p.p. 487-509.
- Colzieher, J.W., and Dozier, T.S.: Improving the Diagnostic Reliability of Rapidly Fluctuating Plasma Hormones by Optimized Multiple Sampling Techniques. Clin Endocrinol Metab, 1976;43:824.
- Bonde, J.P.E., Ernst, E., and Jensen, T.: Relation between Semen Quality and Fertility. Lancet, 1998; 353: 1172-7.
- Sigman, M., Stuart, S.: Howard's, Male Infertility, Campbell's Urology. 7th edition, 1998; p.p. 1287-330.
- 9. Xu, B., Chia, S.E., and Ong, C.H.: Concentrations of Cadmium, Lead, Selenium, and Zinc in Human Blood and Seminal Plasma. Biol trace elem Res, 1994; 40 (1): 49-57.
- 10. Ward, R.J., and Hendry, W.F.: A Cooperative Study of Zinc, Copper, Cadmium, and Lead Levels In Fertile and Infertile Men. Fertile Sterile, 1983; 40 (5): 670-7.
- Schoenfeld, C., Ameral, R.D., Dubin, L., and Numberoff, M.: Prolactin, fructose and zinc levels found in human seminal plasma. Fertile Sterile, 1979; 32: 206-8.
  Lewis, D.L., Jones, L.A., Arid, M.M., and Billian. C. R.: Effects of Snerr Mactivity on Zinc and
- Lewis, D.I., Jones, I.A., Arid, M.M., and Biljan, C.R.: Effects of Sperm Activity on Zinc and Fructose concentrations in Seminal Plasma. Hum Reprod, 1996; 11:2465-7.
  Zvit, H., et al: The Cation Composition of The Seminal Plasma and Prostatic Fluid and
- Zvir, H., et al. The Cation Composition of the seminar Prasma and Prostatic Full and its Correlation to Semen Quality. Fertile Sterile, 1978; 29 (5):539-42.
  Euse, H. Kazaman, T. Ohta, S. and Euliuchi, Y. Relationship, between Zinc
- Fuse, H., Kazaman, T., Ohta, S., and Fujiuchi, Y.: Relationship between Zinc Concentrations in Seminal Plasma and Various Sperm Parameters, Int Urol Nephrol, 1999;31 (3):401-8.
- 15. Deadre, J., Johnson et al.: Improved Colorimetric Determination of Serum Zinc. Clin Chem, 1977; 23 (7): 1321-3.
- Fuentes, J., Miro, J., and Riera, J.: Simple Colorimetric Method for Seminal Plasma Zinc Assay. Andrologia, 1982; 14 (4): 322-7.