



## ANTIBACTERIAL ACTIVITY OF CASSIA OCCIDENTALIS L. LEAF AND SEED EXTRACTS

R. Lavanya

Department of Pharmacy, Government Polytechnic for women, Nizamabad, Telangana, India.

### ABSTRACT

The development of antibiotic resistance in harmful bacteria has led to the screening of plants for their antimicrobial activity. In the present work an attempt has been made to find out the antimicrobial activity of *Cassia occidentalis* L. (Caesalpinaceae). The various solvent extracts of leaves and seeds of the plant were screened for anti bacterial activity. The anti bacterial activity was done by agar well diffusion method.

**KEYWORDS** : *Cassia occidentalis* L., anti bacterial activity, solvent extracts

### INTRODUCTION

Infectious diseases are a major cause of death in many countries<sup>1</sup>. Resistance to antimicrobial agents is emerging in a wide variety of pathogens, this has necessitated a search for new anti microbial compounds. There are several reports on antibacterial activity of different plant extracts<sup>2-4</sup>. *Cassia occidentalis* leaf extracts exhibited antimalarial<sup>5</sup>, anticarcinogenic<sup>6</sup>, antidiabetic<sup>7</sup>, antimicrobial<sup>8-9</sup>, antiinflammatory<sup>10</sup>, analgesic & antipyretic<sup>11</sup> and hepatoprotective activity<sup>12</sup>.

### *Cassia occidentalis* L.

#### Taxonomic classification:

**Kingdom**- Plantae

**Class**- Magnoliopsida

**Order**- Fabales

**Family**- Fabaceae

**Genus**- Senna

**Species**- *S. occidentalis*.

**Synonyms:** *Cassia caroliniana*, *C. falcata* L., *C. macradenia*, *C. obliquifolia*, *C. planisiliqua*, *senna occidentalis*.

#### Vernacular name:

**English**- Coffee senna

**Hindi**- Kasaundi, Bari Kasaundi

**Telugu**- Kasinda

**Sanskrit**- Kasamarda

**Tamil**- Thagarai Tarun.

### MATERIALS AND METHOD

#### Description:

*C. occidentalis* is a small tree, somewhat branched, smooth, semi-woody, fetid herb or shrub, 0.8–1.5 m tall, taproot, hard, stout, with a few lateral roots on mid section. This plant species varies from a semi-woody annual herb in warm temperate areas to a woody annual shrub or sometimes a short-lived perennial shrub in frost free areas<sup>13-14</sup>. The fruit is a dry, dehiscent, transversely partitioned, faintly recurved, laterally compressed, sickle shaped legume (pod), 7–12 cm long, 8–10 mm wide, seeds are oval shaped, 3.5–4.5 mm wide, flattened, pale-dark brown, slightly shiny with rounded tip<sup>15</sup>.



Figure 1: *Cassia occidentalis* L. Plant.

#### Distribution:

*Cassia occidentalis* L. is found in tropical countries<sup>16</sup>. In India it is a perennial plant in south India, but as annual plant in North and North-west India<sup>17</sup>.

#### Plant material:

*Cassia occidentalis* L. plants were collected from the region of Nizamabad, Telangana, India, in the month of October. The plant was authenticated by Dr. Vidya vardini, Head of department, Department of Botany, Telangana University.

#### Preparation of extracts:

**Leaves extract**- *Cassia occidentalis* L. leaves were washed in water, shade dried, broken into coarse powder, grinded to fine powder using mechanical grinder and stored in air tight containers at room temperature till further use. Each solvent extract of sample was prepared by soaking 100 g of dried fine powdered samples in 200 ml of respective solvent (Ethyl acetate, Hexane, Dichloromethane) separately for 4 days at room temperature with occasional shaking. The extracts were filtered using Whatman filter paper and then concentrated. The residual extracts were stored in refrigerator.

#### Seeds extract-

The pods of the plant were shade dried until the seeds lose moisture. The seeds were collected and ground to fine powder. The powder was first defatted with petroleum ether, then each 100g of dried fine powder extracted with Ethyl acetate, Hexane, Dichloromethane separately for 4 days with occasional shaking.

#### Anti bacterial Activity Test by Agar well Diffusion Method:

In this study, one gram positive (*Bacillus subtilis*) and two gram negative bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*) were tested. The Anti antimicrobial assay was performed by agar well

diffusion method [18,19]. The sterilized nutrient agar (HiMedia) was inoculated with 200 µl of the bacterial inoculum and poured into the sterilized Petri plates. Three wells of 6 mm diameter were made on sterilized nutrient agar with a sterile borer. The prepared concentration of 100 mg/ml of each solvent extracts were transferred into the wells. The plates were incubated overnight at 37 °C. Anti bacterial agent Gentamicin (10 µg) used as a positive control and DMSO solvent as negative control. The diameter of clear zone of inhibition was measured.

## RESULTS AND DISCUSSION

The antibacterial activity of plant extracts is shown in Table 1. Among all tested plant extracts ethyl acetate fractions were found to be most effective.

## CONCLUSION

The present study reveals the antibacterial property of leaf and seed extracts of *Cassia occidentalis* L. This study paves the pathway for further research to identify the active chemical constituents responsible for the antibacterial activity.

**Table.1 Antibacterial activity of leaves and seeds extracts of *Cassia occidentalis* L. zone of inhibition in mm diameter.**

Extract	Bacillus subtilis	Escherichia coli	Pseudomonas aeruginosa	
Leaves	Ethyl acetate	12	10	11
	Hexane	7	6	-
	Dichloromethane	10	9	9
Seeds	Ethyl acetate	15	10	11
	Hexane	9	7	-
	Dichloromethane	10	8	8
Gentamicin (10µg as +ve ctrl)	40	38	38	
DMSO(-ve ctrl)	-	-	-	

Each solvent extract concentration at 100 mg/ml DMSO

(-) Value indicates no activity.

## REFERENCES:

- [1] WHO (World Health Organization). The world health report, Shaping the future (2003), Geneva, Switzerland, WHO;2003:pp.11-12.
- [2] Renisheya Joy Jeba Malar T, Johnson M, Mary Uthith M, Arthy A. (2011), "Antibacterial activity of ethanolic extracts of selected medicinal plants against human pathogens." *Asian Pacific Journal of Tropical Biomedicine*. ELSEVIER, 576-578.
- [3] Doriane E Djeussi, Jaurès AK Noumedem, Jackson A Seukep. (2013), "Antibacterial activities of selected edible plants extracts against multidrug-resistant Gram-negative bacteria." *The official journal of the International Society for Complementary Medicine Research (ISCMR)*, 13:164.
- [4] Mativandela SPN, Lall N, Meyer JJM: "Antibacterial, antifungal and antitubercular activity of (the roots of) *Pelargonium reniforme* (CURT) and *Pelargonium sidoides* (DC) (Geraniaceae) root extracts." *S Afr J Bot* 2006, 72:232-237.
- [5] Tona L, Ngimbi NP, Tsakala M, Mesia K, Cimanga K, Apers S, De Bruyne T, Pieters L, Totté J, Vlietinck AJ. (1999). "Antimalarial activity of 20 crude extracts from nine African medicinal plants used in Kinshasa Congo." *Journal of Ethnopharmacology*, 68: 193-203.
- [6] Sharma N, Trikha P, Athar M, Raisuddin S. (2000). "In vitro inhibition of carcinogen-induced mutagenicity by *Cassia occidentalis* and *Emblica officinalis*." *Drug and Chemical Toxicology*, 23:477-84.
- [7] Laxmi Verma, Anirudh Khatri, Basant Kaushik, Umesh K Patil, Rajesh S Pawar. (2010). "Antidiabetic activity of *Cassia occidentalis* (Linn) in normal and alloxan-induced diabetic rats." *224-228*.
- [8] Saganuwan AS, Gulumbe ML. (2006). "Evaluation of in vitro antimicrobial activities and phytochemical constituents of *Cassia occidentalis*." *Animal Research International*, 3:566-569.
- [9] Jain SC, Sharma RA, Jain R, Mittal C. (1998). "Antimicrobial screening of *Cassia occidentalis* L in vivo and in vitro." *Phytotherapy Research*, 12:200-204.
- [10] Sadique J, Chandra T, Thenmozhi V, Elango V. (1987). "Biochemical modes of action of *Cassia occidentalis* and *Cardiospermum halicacabum* in inflammation." *J Ethnopharmacol*, 19,201-12.
- [11] Sini KR, Sinha BN, Karpakavalli M, Sangeetha PT. (2011). "Analgesic and antipyretic activity of *Cassia occidentalis* Linn." *Annals Biol Res*, 2(1):195-200.
- [12] Bhattacharyya D, Mukherjee R, Pandit S, Das N, Sur TK. (2003). "Prevention of carbon tetrachloride induced hepatotoxicity in rats by Himoliv." A polyherbal formulation. *Ind J Pharmacol*, 35:183-5.
- [13] Henty EE, Pritchard GH. (1975). *Weeds of New Guinea and Their Control*, Botany Bulletin 7, Division of Botany, Lae, Papua New Guinea: Department of forests, 189 pp.
- [14] Holm L, Doll J, Holm E, Pancho J, Herberger J. (1997). *World Weeds*. New York, John Wiley and Sons, 129 pp.

- [15] Roy L, Holm G, Doll J, Holm E, Pancho J, Herberger J. (1997). *World Weeds: Natural Histories and Distribution* *Cassia occidentalis* L. and *Cassia tora* L. (Syn. *C. obustifolia* L.). New York: John Wiley & Sons, 1129 pp.
- [16] Saurabh Arya, Jogender Saini and Sumer Singh. (2013). "Antidiabetic activities of *Cassia occidentalis*." *Recent Research in Science and Technology*, 5(1), 51-53.
- [17] Vedapriya Arya, Sanjay yadav, Sandeep kumar, JP Yadav. (2010). "Antimicrobial activity of *Cassia occidentalis* L against various Human Pathogenic microbes." *Life sciences and Medicine Research, Volume 2010:LSMR-9*.
- [18] Murray, PR; Baron, EJ; Pfaller, M A; Tenover, FC and Tenover, HR (1995), "Manual of Clinical Microbiology", 6th Ed. ASM Press, Washington DC, 15-18.
- [19] Olurinola, PF (1996), "A laboratory manual of pharmaceutical microbiology", Idu, Abuja, Nigeria, 69-105.