



## EFFECTS OF DIFFERENT MODES OF YOGA PRACTICE ON PERCENTAGE OF BODY FAT AND BIOCHEMICAL VARIABLES

**Mr. S. Ananth**

Ph.D., Scholar, Department of Physical Education and Sports Sciences, Annamalai University.

**Dr. S. Chidambara Raja\***

Associate Professor, Department of Physical Education and Sports Sciences, Annamalai University. \*Corresponding Author

### ABSTRACT

The purpose of the present study was to find out the effect of different modes of yoga practice on percentage of body fat and bio-chemical variables among middle aged women. For this purpose, forty five middle aged women residing around Kumbakonam town, Thanjavur district, Tamilnadu, were selected as subjects. The age of the subjects were ranged from 40 to 45 years. They were divided into three equal groups, each group consisted of fifteen subjects, in which experimental group - I underwent Bihar School of yoga practice, experimental group - II underwent B.K.S. Iyengar yoga practice and group - III acted as control that did not participate in any special activities apart from their regular curricular activities. The training period for the study was six days (Monday to Saturday) in a week for twelve weeks. Prior and after the experimental period, the subjects were tested on percentage of body fat, high density lipoprotein and total cholesterol. Percentage of body fat was assessed by applying the Deurenberg et al formula. High density lipoprotein and total cholesterol were tested after taking 5 ml of blood samples by venous puncture method, by using Boehringer Mannheim Kit Method. The Analysis of Covariance (ANCOVA) was applied to find out any significant difference between the experimental groups and control group on selected criterion variables. The result of the study shows that the Bihar School of Yoga practice group and B.K.S. Iyengar Yoga practice group were decreased the total cholesterol, percentage of body fat and increased the high density lipoprotein significantly. It was concluded from the results of the study that Bihar School of Yoga practice and B.K.S. Iyengar Yoga practice has bring positive changes in percentage of body fat, high density lipoprotein and total cholesterol as compare to the control group. Moreover it was also concluded that there was no significant difference was found between the experimental groups in all criterion variables.

**KEYWORDS** : Bihar School of Yoga and B.K.S. Iyengar Yoga, Breath holding time, high density lipoprotein, total cholesterol, Boehringer Mannheim kit method, ANCOVA.

### INTRODUCTION

Yoga is one of the most ancient cultural heritages of India. The word yoga in Sanskrit means "to unite", and so yoga can be said to connote a unitive discipline. In this sense, it is an exercise in moral and mental cultivation that generates good health (*arogya*), contributes to longevity (*chirayu*), and the total intrinsic discipline culminates into positive and perennial happiness and peace.[1] Yoga is one of the orthodox systems of Indian philosophy. It was collated, coordinated and systematized by Patanjali in his classical work, the Yoga Sutra, which consists of 185 terse aphorisms. Yoga is a complete science of life that originated in India many thousands of years ago. It is the oldest system of personal development in the world, encompassing body, mind and spirit. [2]

Bihar School of Yoga (Swami Satyananda Saraswati) is a type of yoga which integrates intellect, emotion and action: the head, heart and hands. Known as Satyananda Yoga or Bihar Yoga (the School lies in Bihar in India), the system embraces many different philosophies and encourages students to examine the very essence of their being and make gradual changes to improve their awareness. Satyananda Yoga is considered truly holistic and suitable for everyone.[3] Iyengar Yoga, named after and developed by B.K.S. Iyengar, is a form of Hatha Yoga that has an emphasis on detail, precision and alignment in the performance of posture (*asana*) and breath control (*pranayama*).[4]

Recent scientific studies of the effects of yoga and meditation on health validates its ability to improve virtually every aspect of our functioning—brain function, hormonal function, sleep, mood, balance, etc. More active practices followed by relaxing ones lead to deeper relaxation than relaxing practices alone, documented by research from Swami Vivekananda yoga research foundation near Bangalore city and possibility of neuroplasticity bringing about changes in the hypo-pituitary–pancreatic axis.[5] The improvement in the lipid levels after yoga could be due to increased hepatic lipase and lipoprotein lipase at cellular level, which affects the metabolism of lipoprotein and thus increase uptake of triglycerides by adipose tissues.[6,7] Direct stimulation of the pancreas by the postures can rejuvenate its capacity to produce insulin.[8] Regeneration of

pancreatic beta cells could occur by yoga exercises that promote blood circulation in the region of the pancreas and yoga asanas that stimulate the meridian of pancreas also could assist in some diabetic patients.[9] Pranayama practices, stretches the lung tissue producing inhibitory signals from action of slowly adapting receptors and hyperpolarizing currents. These inhibitory signals coming from cardio-respiratory region involving vagi are believed to synchronize neural elements in the brain leading to changes in the autonomic nervous system; and a resultant condition characterized by reduced metabolism and parasympathetic dominance.[10]

### METHODOLOGY

Thirty middle aged women residing around Kumbakonam town, Thanjavur district, Tamilnadu were selected as subjects. The age of the subjects were ranged from 18 to 23 years. The selected subjects were divided into three equal groups, each group consisted of ten subjects, in which group - I (n = 15) underwent Bihar School of Yoga practice, experimental group - II (n = 15) underwent B.K.S. Iyengar Yoga practice and group - III (n = 15) acted as control, which did not participate in any special activities apart from their regular curricular activities. Different modes of yoga practices were conducted six days (Monday to Saturday) per week for twelve weeks. The researcher consulted with the yoga experts and selected the following variables as criterion variables: 1. percentage of body fat, 2. high density lipoprotein and 3. total cholesterol. Percentage of body fat was assessed by applying the Deurenberg et al [11] formula, high density lipoprotein and total cholesterol were assessed by using the Boehringer Mannheim Kit method. For the purpose of collection of data the subjects were asked to report at early morning, one day prior and one day after experimental period, in fasting condition. 5 ml of blood was collected from each subject by venous puncture method and the blood thus collected was stored in small bottles for pre and post-test for measuring the high density lipoprotein and cholesterol.

Analysis of covariance (ANCOVA) was applied to find out the significant difference if any, among the experimental groups and control group on selected criterion variables separately. In all the

cases, .05 level of confidence was fixed to test the significance, which was considered as appropriate. After applying the analysis of covariance, the result of this study shows that there was a significant increase in breath holding time, decrease in total cholesterol and high density lipoprotein levels.

**RESULTS**

The data collected on percentage of body fat, high density lipoprotein and cholesterol among experimental and control groups were analysed and the results were presented in Table – I.

**Table - I ANALYSIS OF COVARIANCE ON SELECTED CRITERION VARIABLES AMONG EXERCISE GROUPS AND CONTROL GROUP**

Variable Name	Group Name	Bihar School of Yoga Practice Group	B.K.S. Iyengar Yoga Practice Group	Control Group	'F' Ratio
Percentage of body fat (in percentage)	Pre-test Mean ± S.D	29.59 ± 1.58	29.98 ± 1.56	29.0 ± 1.61	0.96
	Post-test Mean ± S.D.	27.47 ± 1.16	27.76 ± 1.37	29.31 ± 1.11	15.79*
	Adj. Post-test Mean	27.121	27.361	29.692	45.586*
High Density Lipoprotein (mg/dl)	Pre-test Mean ± S.D	41.26 ± 1.19	41.36 ± 2.05	41.15 ± 1.95	0.59
	Post-test Mean ± S.D.	42.98 ± 1.86	42.56 ± 1.39	41.09 ± 1.15	21.166*
	Adj. Post-test Mean	43.154	43.081	41.101	56.87*
Total Cholesterol (mg/dl)	Pre-test Mean ± S.D	193.37 ± 3.78	193.55 ± 3.05	193.81 ± 3.55	0.861
	Post-test Mean ± S.D.	190.36 ± 2.87	190.77 ± 3.00	193.46 ± 3.58	21.532*
	Adj. Post-test Mean	190.056	190.266	193.573	80.375*

\*Significant .05 level of confidence. (The table values required for significance at .05 level of confidence with df 2 and 42 and 2 and 41 were 3.22 and 3.21 respectively).

Table – I shows that pre test mean 'f' ratio of Bihar School of Yoga practice group, B.KS Iyengar practice group and control group on percentage of body fat were 0.96 and which is insignificant at 0.05 level of confidence. The post and adjusted post test mean 'f' ratio value of experimental groups and control group were 15.79 and 45.586, which was significant at 0.05 level of confidence. The pre and post test means 'f' ratio of Bihar School of Yoga practice group, B.KS Iyengar practice group and control group on high density lipoproteins were 0.59, which is insignificant and 21.166, which is

significant at 0.05 level of confidence. The adjusted post test mean 'f' ratio value of experimental groups and control group was 56.87, which was significant at 0.05 level of confidence. The pre test means 'f' ratio of yogasana practice group, aerobic exercise group and control group on total cholesterol were 0.861 106 and 1.532 which is insignificant at 0.05 level of confidence. The post and adjusted post test means 'f' ratio value of experimental groups and control group were, 21.532 and 80.375, which was significant at 0.05 level of confidence.

**Table - II Scheffé S Test for the Difference Between the Adjusted Post-Test Mean of Selected Criterion Variables**

Adjusted Post-test Mean on Resting Pulse Rate				
Yogasana Practice Group	Aerobic Exercise Group	Control Group	Mean Difference	Confidence interval at .05 level
27.121	27.361		0.24	1.117
27.121		29.692	2.571*	1.117
	27.361	29.692	2.331*	1.117
Adjusted Post-test Mean on High Density Lipoproteins				
Yogasana Practice Group	Aerobic Exercise Group	Control Group	Mean Difference	Confidence interval at .05 level
43.154	43.081		0.073	0.861
43.154		41.101	2.053*	0.861
	43.081	41.101	1.98*	0.861
Adjusted Post-test Mean on Total Cholesterol				
Yogasana Practice Group	Aerobic Exercise Group	Control Group	Mean Difference	Confidence interval at .05 level
190.056	190.266		0.21	1.591
190.056		193.573	3.514*	1.591
	190.266	193.573	3.307*	1.591

\* Significant at .05 level of confidence.

Table – II shows that the Scheffé S Test for the difference between adjusted post-test mean on percentage of body fat of Bihar School of Yoga practice group and control group (2.571) and BKS Iyengar Yoga Practice group and control group (2.331), which were significant at .05 level of confidence. There was a significant difference on high density lipoproteins between Bihar School of Yoga practice group and control group (2.053) and BKS Iyengar Yoga Practice group and control group (1.98) and also there was a significant difference on total cholesterol between Bihar School of Yoga practice group and control group (3.514) and BKS Iyengar Yoga Practice group and control group (3.307) which was significant at 0.05 level of confidence after the respective training programme.

**CONCLUSIONS**

The experimental groups such as, Bihar School of Yoga practice group and B.K.S. Iyengar Yoga practice group have achieved a significant improvement in selected criterion variables such as percentage of body fat [12,13], high density lipoprotein [14,15.] and a significant reduction in total cholesterol [16], when compared with the control group. It was also found that there was no significant difference was found between the Bihar School of Yoga

practice group and B.K.S. Iyengar Yoga practice group on selected criterion variables.

**Reference:**

- Retrieved from <http://hinduism.about.com/bl-yoga-define.htm> on 24-04-2012.)
- Swami Vishnu Devananda, The Sivananda Companion to Yoga, (New York: Fireside Book, Simon and Schuster, 2000), p. 10.
- Retrieved from <http://samsaramindandbody.com/different-types-yoga-classes-ultimate-guide>
- Retrieved from [http://en.wikipedia.org/wiki/Iyengar\\_Yoga](http://en.wikipedia.org/wiki/Iyengar_Yoga) on 22-7-2014.
- McCall T. The Scientific Basis of Yoga Therapy. [Accessed Jun 16, 2012]. at [http://www.yogajournal.com/for\\_teachers/2016](http://www.yogajournal.com/for_teachers/2016).
- Delmonte MM. Biochemical indices associated with meditation practice: A literature review. Neurosci Biobehav Rev. 1985;9:557–61. [PubMed]
- Tulpule TH, Shah HM, Shah SJ, Haveliwala HK. Yogic exercises in the management of ischaemic heart disease. Indian Heart J. 1971;23:259–64. [PubMed]
- Ramaiah SA. Yoga Therapy for Diabetes: Washington, D.C. Study, International Conference on Traditional Medicine, 1986, Madras. Madras, India: Published by Siddha Medical Board, Govt. of Tamil Nadu;
- Yogalink. A community service donated by samyama yoga. [Accessed Jul 9, 2012]. at <http://www.yogalink.com.au>.
- Jerath RJ, Edry VA, Barnes VA, Jerath V. "Physiology of Long Pranayamic Breathing: Neural Respiratory Elements May Provide A Mechanism that Explains How Slow Breathing Shifts the Autonomic Nervous System", Med Hypotheses. 2006;67:566–71. [PubMed]
- Retrieved from <http://www.halls.md/bmi/fat.htm> on 15-08-2012.
- C. Kesavachandran, V. Bihari and N. Mathur, "Can Physical Activity Maintain Normal

- Grades of Body Mass Index and Body Fat Percentage?", *International Journal of Yoga*, 2:1, (January – June 2009), 26 - 29.
- [13] Ajay Singh Ruhall, Rakesh Bhandari and Ranjan Chakravarti, "Effect of Kapalabhati on Selected Body Composition Variables", *British Journal of Sports Medicine*, 44, (2010), i 70.
- [14] A. Pal, N. Srivastava, S. Tiwari, N.S. Verma, V.S. Narain, G.G. Agrawal, S.M. Natu and K. Kumar, "Effect of Yogic Practices on Lipid Profile and Body Fat Composition in Patients of Coronary Artery Disease", *Complementary Therapies in Medicine*, 19:3, (June 2011), 122-7.
- [15] Vijay Tundwala, R.P. Gupta, Surendra Kumar, V.B. Singh, B.R. Sandeep, Prabhu Dayal and Parul Prakash, "A Study on Effect of Yoga and Various Asanas on Obesity, Hypertension and Dyslipidema", *Internation Journal of Basic and Applied Medical Sciences*, 2:1, (January – April 2012), 93 – 98.
- [16] S. Telles, V.K. Naveen, A. Balkrishna and S. Kumar, "Short Term Health Impact of a Yoga and Diet Change Program on Obesity", *Medical Science Monitor*, 16:1, (January 2010), CR35-40.