



COMPARISON OF YOGA AND WALKING-EXERCISE ON CARDIAC TIME INTERVALS AS A MEASURE OF CARDIAC FUNCTION IN ELDERLY WITH INCREASED PULSE PRESSURE

Dr. Sameer Saraf

Department of Cardiology, U.P. University of Medical Sciences, Saifai, Etawah, Uttar Pradesh, India.

**Rajesh Kumar
Gautam***

Senior Resident, Department of Medicine, U.P. University of Medical Sciences, Saifai, Etawah, Uttar Pradesh, India. *Corresponding Author

ABSTRACT

Objective: Arterial aging along with increased blood pressure (BP) has become the major cardiovascular (CV) risk in elderly. The aim of the study was to compare the effects of yoga program and walking-exercise on cardiac function in elderly with increased pulse pressure (PP).

Methods: An open label, parallel-group randomized controlled study design was adopted. Elderly individuals aged 60 years with PP 60 mmHg were recruited for the study. Yoga (study) group (n = 30) was assigned for yoga training and walking (exercise) group (n = 30) for walking with loosening practices for one hour in the morning for 6 days in a week for 3 months. The outcome measures were cardiac time intervals derived from pulse wave analysis and ECG, resting heart rate (RHR), diastolic time (DT), ejection duration index (ED%), rate pressure product (RPP) and percentage of mean arterial pressure (% MAP).

Results: The mean within-yoga group change in RHR (bpm) was 4.41 (p = 0.031), PD (ms): 50.29 (p = 0.042), DT (ms): 49.04 (p = 0.017), ED%: 2.107 (p = 0.001) and %MAP: 2.08 (p = 0.04). The mean within-control group change in HR (bpm) was 0.35 (p = 0.887), PD (ms): 11.15 (p = 0.717), DT (ms): 11.3 (p = 0.706), ED%: 0.101 (p = 0.936) %MAP: 0.65 (p = 0.451). A significant difference between-group was found in RHR (p = 0.036), ED% (p = 0.049), DT (p = 0.02) and RPP (p = 0.001).

Conclusions: Yoga practice for 3 months showed a significant improvement in diastolic function with a minimal change in systolic function. Yoga is more effective than walking in improving cardiac function in elderly with high PP.

KEYWORDS : Yoga program, Walking-exercise, Cardiac function, Diastolic function.

Introduction:

Arterial aging along with increased blood pressure (BP) has become the major cardiovascular (CV) risk in elderly. Structural and functional changes take place in heart and blood vessels with advancing age. The reduction of vascular compliance with age due to stiffening of arteries is the major contributor for elevation of BP, especially systolic pressure resulting in isolated systolic hypertension (ISH) in elderly. Systolic BP increases and diastolic BP falls with age leading to widening of pulse pressure (PP). Pulse pressure is a best tool for measuring vascular aging and a good marker for CV risk in elderly. Pulse pressure is an independent indicator of arterial stiffness.¹ It is more closely associated to CV events than systolic BP or diastolic BP alone.²

Age-associated stiffening of aorta and increased systolic pressure increases left ventricle (LV) after-load. Increased ventricular load causes LV hypertrophy and increases LV oxygen demand. Due to these changes, LV contract and relax slowly, so that systolic time is increased while diastolic time is reduced.³ Left ventricle becomes stiff with decreased compliance and impaired relaxation, leading to increased end diastolic pressure and diastolic dysfunction. Age associated major changes in heart function occurs mainly in left ventricular diastolic function. So, elderly individuals (especially hypertensive patients) often manifest diastolic dysfunction.⁴ Although physical training programs are recommended for CV prevention and rehabilitation,⁵ their effects on cardiac function in elderly individuals remain unclear. Few authors demonstrated a beneficial effect of exercise training program on left ventricular function in elderly⁶⁻⁸ while others reported either least change⁹ or no change.¹⁰⁻¹⁴ Moreover, most of these studies are done on trained athletes and healthy elderly individuals. Yoga is emerging as an effective life-style modality and mind-body medicine. It is a skill to control mind and involuntary functions voluntarily. Many studies have showed beneficial effects of yoga program on CV health in young and older adults.¹⁵⁻¹⁸ Again, most of the studies have either included young and middle aged individuals or mixed population with wide range of age. We did not find any studies investigating the effect of yoga on cardiac function in elderly individuals with hypertension. In the present study, we compared the effects of yoga program and walking-exercise on cardiac function in elderly with

increased PP (as mentioned previously that PP is a better predictor of CV events than SBP or DBP in elderly).

2. Methods:

2.1. Participants and study design:

A total of 60 elderly individuals aged 60 years with increased PP >60 mmHg were recruited for the study. Volunteers were screened from geriatric health camp and Geriatric clinic of UPUMS, Saifai, Etawah. An open label, parallel-group randomized controlled study design was adopted. Exclusion criteria includes subjects with SBP > 159 mmHg and DBP > 99 mmHg; CV risk factors such as diabetes mellitus, hypercholesterolemia and high triglyceride level, history of secondary hypertension, neuromuscular disorders, alcoholism, practicing yoga for one hour/day for three days in a week and on any medications. Participants were instructed for not to consume any vitamin supplements or herbal drugs during the study period. This criterion for selection of elderly subjects with high PP for life-style changes intervention for three months was as per the 2007 guidelines of the task force for the management of arterial hypertension of the European Society of hypertension and of the European Society of Cardiology.¹⁹

During visit 1-3 (three consecutive days), volunteers were screened and their BP was measured. Baseline investigation, randomization and allocation of subjects were done at their 4th visit. They were randomly allocated to Yoga group (YG, n = 30) and exercise group (EG, n = 30) by using random number table. At visit 5, post-intervention investigations were made. All the investigations were done in the morning between 8.00 h to 11.00 h after supine rest for 10 min. No intervention was given on the day of investigation. Persons handling data analysis were kept blinded.

2.2. Ethics statement:

A prior study approval has been obtained from the Institutional ethical committee of UPUMS, Saifai, Etawah, India, as per the guidelines of Indian Council of Medical Research (ICMR).²⁰ The declaration of Helsinki has been followed during the entire study. Informed written consent was obtained from the participants.

2.3. Intervention:

The yoga program included Loosening practices, Asanas (maintaining postures), Pranayama (breathing exercises) and cyclic meditation (Table 1). Asanas were practiced for 15–20 min while Pranayama and relaxation technique/meditation for 40– 45 min. Emphasis was placed on practicing slow and paced breathing with asanas (maintaining postures) and other techniques of Yoga program. Cyclic meditation (CM)] is a cycle of alternating stimulation and relaxation that used to go into deep silence. It is a guided relaxation technique of about 23 min. During the practice of CM, the subjects followed the instructions with eyes closed. It includes stretching and relaxing the muscles consciously (in various postures) with internal awareness by observing changes in the system. The sequence of practice is as follows:

(1) It began by chanting a verse (40s) in supine position with Namaskar Mudra followed by isometric contraction of the muscles from toe to head and relaxation with awareness (1 min). (2) Linear awareness was observed in standing posture (Tadasana) and balancing the weight on both feet at ease (2 min). (3) slowly moving to the next posture (ardhakatichakr asana), bending to the right (1 min 20 s) followed with instructions about relaxation and awareness (1 min 20 s), then bending to the left (1 min 20 s) followed with relaxation (1 min 20 s). (4) Slowly lied down in supine posture, right arm stretched, turned to right side with head on the right biceps, (linear awareness) then rested on back, observed the abdominal movements and breathing (3 min). (5) Moved to the sitting posture (Vajra asana) and observed the changes (1 min 20 s). (6) Chanted (MMM) M-Kara in another sitting posture (Sasank asana) (1 min 20 s). (7) Chanted (AAA) A-Kara in backward bending on knees posture (Ustrasana) (1 min 20 s). (8) Relaxed in supine posture (Shavasana) and chanted A, U, M kara (7 min). The postures were practiced slowly, with awareness of all the sensations that are felt.²¹ The protocol for the walking-exercise group consisted of loosening practices like neck rotation, shoulder and hip rotation, wrist and ankle rotation, forward and side bending and walking (40–50 min) followed by rest (10 min). Intervention for both the groups was given for one hour/day for 6 days in a week in the morning from 06:00 h to 07:00 h for twelve weeks under the supervision of experienced authorized instructor.

2.4. Measurement of blood pressure:

As BP is more variable in older people, an average of nine BP readings (measured thrice with an interval of one minute on every visit for three consecutive days in a sitting posture) was taken using mercury sphygmomanometer (Diamond, Industrial Electronic and Allied products, India).²² Pulse pressure was estimated as the difference between systolic and diastolic BP. Mean arterial pressure (MAP), an average arterial pressure in an individual during single cardiac cycle was estimated by adding 1/3rd of PP to DBP.

Table 1

Integrated yoga module for elderly subjects with increased pulse pressure.

Sl. No	Practice		Duration
1.	Opening Prayer		1 min
2.	Sukshma Vyayama (Loosening Practices)	Loosening of Fingers	5 min
		Loosening of Wrist	
		Shoulder rotation	
		Ankle stretch/rotation	
		Drill walking	
3.	Breathing Practices	Hands in and out breathing	5 min
		Ankle stretch breathing	
		Straight leg raising breathing	
		Lumbar stretch breathing	
4.	Asana	Shashank asana	15 min

		Bhujang asana	
5.	Pranayama	Anuloma Viloma Pranayama	5 min
		Brahmari Pranayama	
6.	Cyclic Meditation [CM]		23 min
7.	Devotional Session – Chanting/songs	5 min	
8.	Closing prayer		1 min

2.5. Assessment of cardiac function:

Cardiac function was assessed by measuring the cardiac time intervals using 8-channel non-invasive automatic device (Periscope, Genesis Medical Systems, India). Periscope is a real time PC-based simultaneous acquisition (200 samples per second) and analysis system. This device uses four BP cuffs and two-channel ECG leads to record arterial pressure waveforms and ECG simultaneously. The recordings were made in supine position. BP cuffs were wrapped over both upper arms (brachial artery) and legs (tibial artery) above ankle. ECG electrodes were placed on the ventral surface of both wrists and medial side of the ankles. The BP cuffs were connected to oscillometric pressor sensor and plethysmographic sensor to determine pressure waveforms and volume pulse waveform. The data obtained in 10s was stored in the computer for further analysis. Periscope supports a sophisticated digital-signal algorithm to calculate all the results. As the device is fully automated and does not require any operator for handling any probe to record the waveforms, so it is devoid of any operator bias. It calculates cardiac time intervals using limb pressure waveforms and ECG.

2.5.1. Pre-ejection period (PEP):

It is a period of electromechanical delay and isovolumetric contraction. It is the time between the beginning of the depolarization of left ventricle and the opening of the aortic valve.

2.5.2. Ejection duration index (ED%):

The ratio of the duration of systolic ejection to the total duration of a cardiac cycle is the ejection duration index.

2.5.3. Diastolic time (DT):

It is a period of ventricular filling. It is the duration between opening and closing of mitral valve.

2.5.4. Heart rate (HR):

It was calculated from R-R interval of ECG.

2.5.5. %MAP:

It is the percentage of Mean Arterial Pressure height from diastolic pressure to the total pulse pressure. Higher the %MAP greater is the hardening of artery.

2.6. Rate pressure product (RPP):

It is a sensitive index of myocardial oxygen consumption (MVO₂) or demand and cardiac workload. It is also called as cardiovascular product or double product. RPP is a product of heart rate (bpm) and systolic BP (mmHg).²³

2.7. Estimation of biochemical parameters:

The blood sample was collected in the morning with overnight fasting. Fasting blood glucose (Trinder's method), serum triglyceride level (glycerol phosphatase-oxidase method), serum cholesterol (cholesterol oxidase-peroxidase enzymatic method) and HDL cholesterol (phosphotungstic acid method) were estimated using commercial diagnostic kits (ERBA-MANNHEIM).

2.8. Statistical analysis: The obtained data was expressed in mean and standard deviation. The differences between the post-intervention and the baseline measures were calculated to determine the significant improvement in the outcome measures using paired 't' test. Within group change was determined using mean within-person change and 95% confidence intervals (95% CI).

Between group differences were determined by calculating the mean between-group changes and confidence intervals. Further, Analysis of covariance (ANCOVA) was used to find the statistically significant differences in the post-intervention values between the two groups. ANCOVA determines the difference between the two treatment effects while controlling the baseline values. Statistical significance was established at $p < 0.05$. Data were analyzed by using SPSS software version 20.

3. Results:

3.1. Participant flow and baseline characteristics:

Altogether 44 subjects (study group-24; exercise group-20) with technically successful pressure waveform and ECG recordings were included for data analysis. Out of 60 male subjects recruited for the study, data of 44 subjects were analyzed. The baseline characteristics of the participants were shown in Table 2. There was no significant difference in age, BMI, HR, BP blood glucose and lipid profile between the yoga and walking groups. These findings indicate an equal distribution of subjects to both groups. Blood glucose and lipid profile of all participants values were within normal range. All the participants recruited for the study were residing surrounding the site of study and use to visit our campus ground regularly either morning or evening. Therefore, the adherence of the participants to intervention program was good: about 91% participants attended yoga program while 89% for walking-exercise. Most of the participants were retired employees and habitual of slow nature walking.

Table 2 Baseline characteristics of participants.

Variable	Yoga group (n = 24)		Exercise group (n = 20)		p-Value
Age (Years)	68.50	4.85	69.30	5.932	0.57
BMI (kg/m ²)	24.64	3.65	25.17	3.90	0.587
Heart Rate (bpm)	70.77	9.08	73.08	11.40	0.389
Systolic BP (mmHg)	146.87	5.72	145.83	6.33	0.51
Diastolic BP (mmHg)	74.2	4.60	75.57	5.685	0.31
Pulse Pressure (mmHg)	72.17	6.018	70.33	6.321	0.255
MAP (mmHg)	98.27	4.525	97.37	8.364	0.606
Fasting Blood Glucose (mg/dl)	93.83	11.63	91.73	11.94	0.493
Serum Triglyceride (mg/dl)	93.96	25.45	99.00	23.15	0.426
Total Cholesterol (mg/dl)	152.5	24.32	154.23	19.79	0.763
HDL Cholesterol (mg/dl)	46.8	4.21	46.16	4.29	0.567

3.2. Heart rate:

There was a significant reduction in HR ($p = 0.031$) and increase in PD ($p = 0.042$) within-yoga group while no change was noticed within-exercise group (Table 3). A significant mean between-group change was also observed in HR ($p = 0.036$) and PD ($p = 0.021$) (Table 4).

3.3. Ventricular function:

Table 3 shows cardiac time intervals in participants of yoga and walking-exercise group. Among the ventricular systolic function indices, there was no significant change in PEP, LVET and UT within-yoga and within-exercise group. Though statistically not significant but a mean increase in LVET was noticed within-yoga group while in walking group it was decreased. Ejection duration index was significantly reduced ($p = 0.001$) within-yoga group following yoga practice while no alteration in walking-exercise group was observed. ANCOVA showed significant between-group change in ED% and UT (Table 4).

Table 3 Heart rate and Cardiac time intervals: Baseline and post-interventional values in participants and within group changes.

Variables	Yoga group (n = 24)		p-Value	Exercise group (n = 20)		p-Value
	Before	After		Before	After	
	Mean SD	Mean SD		Mean SD	Mean SD	
RHR (bpm)	70.88 12.39	66.45 7.4	0.031*	72.15 11.08	71.8 10.06	0.887
ED%	33.24 4.59	31.13 2.69	0.001* *	32.83 4.32	32.93 4.22	0.936

Diastolic time (ms)	587.04 126.27	636.08 94.42	0.017*	581.9 119.85	570.6 105.1	0.706
RPP	10382.12 1708.72	8843.54 1001.8	0.001	9504.54 3378.74	9581.54 3307.82	0.978

RHR = Resting heart rate; ED% = ejection duration index; RPP = Rate pressure product.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Table 4 Change (gain) Scores in cardiac time intervals and rate pressure product.

Variables	Yoga group	Exercise group	Between group change
	Mean]FID\$T4[change (95% CI)	Mean]FID\$T4[Change (95%CI)	P[5T\$DIF]-value
RHR (bpm)	4.42 (0.43, 8.4)*	0.35 (- 4.71, 5.41)	0.036*
ED%	2.12 (0.92, 3.28)*	0.1 (- 2.7, 2.5)	0.049*
Diastolic time (ms)	49.04 (88.47, 9.61) *	11.3 (50.56, 73.16)	0.02*
Rate pressure product (RPP)	1538 (999.49, 2077.67)	10.45 (796.95, 776.05)	0.001

RHR = Resting heart rate; ED% = ejection duration index; RPP = Rate pressure product.

* $p[6T$DIF] < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

3.3. Ventricular function:

Table 3 shows cardiac time intervals in participants of yoga and walking-exercise group. Among the ventricular systolic function indices, there was no significant change in PEP, LVET and UT within-yoga and within-exercise group. Though statistically not significant but a mean increase in LVET was noticed within-yoga group while in walking group it was decreased. Ejection duration index was significantly reduced ($p = 0.001$) within-yoga group following yoga practice while no alteration in walking-exercise group was observed. ANCOVA showed significant between-group change in ED% and UT (Table 4).

Diastolic time was significantly increased ($p = 0.017$) within-yoga group while no significant difference was noticed within walking-exercise group ($p = 0.706$). Further, ANCOVA showed significant between-group change in DT indicating that yoga is effective in inducing favorable changes in diastolic function in elderly (Table 4).

3.4. Rate pressure product (at rest):

Within-group analysis has shown a significant decrease in RPP in yoga group while no significant change was noticed in exercise group (Table 3). Between-group analysis showed a significant difference in post-intervention RPP between yoga and exercise group indicating that yoga is better than walking-exercise in improving myocardial performance (Table 4).

3.5. Arterial stiffness:

Fig. 1 shows mean change in %MAP in Yoga and walking group. Within-yoga group, a significant reduction in mean %MAP ($p = 0.004$) was observed with no change within walking exercise group.




Fig.1. Arterial stiffness (%MAP): Baseline and after-intervention values in Yoga and walking-exercise group participants.

4. Discussion:

To our knowledge, influence of yoga on cardiac function in elderly with hypertension has not been studied previously. We believe that the present study may be the first randomized controlled study on influence of yoga on cardiac function in elderly with increased PP. First part of our study related to yoga effects on vascular function has been reported.²⁴ Present study has shown that yoga practice can induce favorable modulation in cardiac function in elderly with mild hypertension. Twelve weeks of yoga training have significantly increased diastolic time, reduced HR and increased PD, decreased RPP and arterial stiffness. Minimal change was noticed in systolic function with yoga practice. Yoga can be used as a physiological approach for improving the efficiency of cardiac function in elderly. Increased resting HR is an independent CV risk factor²⁵ and is associated with higher mortality and morbidity.^{26,27} Higher resting HR has been attributed to sympathetic over activity, impaired myocardial oxygen delivery, elevation in BP and increased arterial stiffness.²⁸⁻³⁰ Koskela et al. shown that lower resting HR is associated with a more beneficial hemodynamic profile.³⁰ Higher HR affect cardiac function by increasing left ventricular work load and myocardial oxygen demand. In addition, age-associated increase in aortic stiffness and BP in elderly increases after load on left ventricle leading to left ventricular hypertrophy and cardiac failure.⁵ Heart rate reduction is a choice of treatment in patients with coronary artery disease, which reduces myocardial oxygen consumption and improves subendocardial blood flow.³¹ In the present study, we found that yoga program has effectively reduced resting HR (6.25%; $p = 0.031$) and increased PD (unlike beta blockers), which may be attributed to reduction in sympathetic over activity and central arterial stiffness.²⁴ We and other researchers have shown that yoga can induce beneficial diastolic pressure and diastolic dysfunction.^{4,5} Aging has least impact on systolic function. Hypertension is also the most common cause of diastolic dysfunction and failure in elderly.¹⁹ Therefore, hypertension along with aging becomes the powerful risk for diastolic dysfunction in older individuals. Diastole is the time for cardiac perfusion. Decrease in DT reduces myocardial perfusion and its performance. Availability of time for ventricular relaxation, diastolic filling and cardiac perfusion is determined by heart rate.⁵ The present study has shown a significant increase in DT (8.35%, $p=0.017$) in study group participants following yoga practice indicating an improvement in the diastolic function (or relaxation). Increase in DT also enhances coronary blood flow or cardiac perfusion. Enhancement in left ventricular diastolic filling is an important factor for adequate cardiac output and working/exercise capacity. Yoga induced reduction in HR might have given more time for cardiac relaxation or diastole. Further, improvement in diastolic function in yoga practitioners can also be probably attributed to reduction in central arterial stiffness,²⁴ BP and oxidative stress.²⁴ We found minimal change in the systolic function with yoga practice. Among the systolic function markers, ED% was significantly reduced with no change in PEP, LVET and UT in yoga group.

Walking is a simple and effective physical exercise. However, our study has not found any beneficial modulation in cardiac function in participants of walking exercise group. The pattern, duration and dosage of physical exercise for cardiac health in modulation in age-associated cardiac autonomic changes and reduce sympathetic tone in elderly with hypertension²⁴ and healthy individuals as well.³² Further, reduction in RPP at rest in participants of yoga group (present study) demonstrates a decrease in myocardial oxygen demand and improvement in myocardial performance. Rate pressure product is the amount of oxygen required by the myocardium to perform a given activity.²³ Our finding suggests that yoga practice can improve the efficacy of cardiac function with less utilization of energy. Another study has showed that Yoga based cyclic meditation and relaxation technique can reduce oxygen consumption.³³ However, oxygen consumption may vary among the various yoga based practices or techniques.

Due to age associated changes in heart such as stiffness and hypertrophy, left ventricle contract and relax slowly, so that systolic time is increased and of diastolic time is reduced (impaired

relaxation) leading to increased end elderly are still not clear. Galetta et al., has demonstrated that the expected pattern of left ventricular diastolic function changes normally seen in response to age is modified in the elderly trained athletes and suggested that exercise training can effectively prevent aging effects on heart in older individuals.⁹ Arbab-Zadeh et al., evaluated left ventricular compliance in master athletes and sedentary seniors by echocardiography and has shown that prolonged, sustained endurance training preserves ventricular compliance with aging.⁸ While in another study, Doppler measures of diastolic function were compared in older athletes with age matched sedentary seniors and found that lifelong endurance training can only partially minimize age-associated changes in myocardial filling and relaxation.⁷ On the contrary, other studies showed that regular aerobic-endurance exercise does consistently modulate the age-associated changes that occur in left ventricular structure and diastolic function.^{11,12} An exercise training program for 6 months in elderly has showed a significant improvement in diastolic function in healthy during rest and exercise³⁵ while in hypertensive subjects minimal favorable change was noticed.¹⁰ Ehsani et al. has demonstrated that endurance exercise training for one year can enhance left ventricular systolic performance at peak exercise in healthy older men.³⁶

As most of our study participants were habitual of slow walking (mentioned earlier), so benefits of physical activity on cardiac health might have reached to threshold level and therefore, probably there was no change in cardiac function with walking-exercise intervention. However, it is also important to note that there was no further deterioration in cardiac function in walking group subjects of our study, indicating that walking-exercise has improved the coping ability of heart to sustain the effects of increased BP and aging. A combined program of physical and mental exercise associated with slow breathing; and relaxation techniques (meditation) for longer duration could induce more beneficial modulation in cardiac function in elderly.

5. Conclusion:

Yoga practice for 12 weeks has showed a significant enhancement in left ventricular relaxation period suggesting an improvement in diastolic function. A minimal change in systolic function has been observed. It is effective in reducing HR and RPP. Yoga is more effective than walking in improving cardiac function in elderly with high PP. Yoga practice can induce favorable changes in cardiac function possible by reducing HR and ventricular after-load through reduction in arterial stiffness and BP.

6. Limitations:

Pulse wave derived indices provide relevant information about cardiac function and moreover it is economical, easy to handle and does not require any operator for handling any probe, so it is devoid of any operator bias. However, echocardiography measures may provide more detailed information on cardiac function than pulse wave derived indices. So, further studies using echocardiography are essential to confirm and explore more information about the yoga effects on cardiac function with large sample size in elderly with hypertension. Another limitation of the study is, although, participants have practiced either yoga or walking under the supervision of authorized instructors for one hour daily, their involvement in other activity (other practices) has not been observed.

Financial disclosure:

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest: None

Financial support: UPUMS, Saifai, Etawah.

References:

1. Lim MA, Townsend RR. Arterial compliance in the elderly: its effect on blood pressure measurement and cardiovascular outcomes. *Clin Geriatr Med.* 2009;25(2):191–205.

2. Franklin SS, Larson MG, Khan SA, Nathan D Wong, Eric P. Leip, et al. Does the relation of blood pressure to coronary heart disease risk change with aging? The Framingham Heart Study. *Circulation*. 2001;103(9):1245–1249.
3. O'Rourke MF, Hashimoto J. Mechanical factors in arterial aging: a clinical perspective. *J Am Coll Cardiol*. 2007;50(1):1–13.
4. Gutierrez C, Blanchard DG. Diastolic heart failure: challenges of diagnosis and treatment. *Am Fam Phys*. 2004;69(11):2609–2617.
5. Mancia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, et al. Guidelines for the management of arterial hypertension: the task force for the management of arterial hypertension of the European society of hypertension (ESH) and the European Society of Cardiology (ESC). *Eur Heart J*. 2007;28:1462–1536.
6. Prasad A, Popovic ZB, Arbab-Zadeh A, Qi Fu MD PhD, Dean Palmer, MS, et al. The effects of aging and physical activity on doppler measures of diastolic function. *Am J Cardiol*. 2007;99:1629–1636.
7. Arbab-Zadeh A, Dijk E, Prasad A, Fu Q, Torres P, et al. Effect of aging and physical activity on left ventricular compliance. *Circulation*. 2004;110:1799–1805.
8. Galetta F, Franzoni F, Femia FR, et al. Left ventricular diastolic function and carotid artery wall in elderly athletes and sedentary controls. *Biomed Pharmacother*. 2004;58:437–442.
9. Zheng H, Lou M, Shen Y, Hong Fang. Improved left ventricular diastolic function with exercise training in hypertension: a Doppler imaging study. *Rehabil Res Pract*. 2011;2011:497690.
10. Gates PE, Tanaka H, Graves J, Seals DR. Left ventricular structure and diastolic function with human ageing: relation to habitual exercise and arterial stiffness. *Eur Heart J*. 2003;24:2213–2220.
11. Nottin S, Nguyen L-D, Terbah M, P. Obert. Long-term endurance training does not prevent the age-related decrease in left ventricular relaxation properties. *Acta Physiol Scand*. 2004;181:209–215.
12. Baldi JC, McFarlane K, Oxenham HC, Whalley GA, Walsh HJ, et al. Left ventricular diastolic filling and systolic function of young and older trained and untrained men. *J Appl Physiol*. 2003;95:2570–2575.
13. Fleg JL, Shapiro EP, O'Connor F, Taube J, Goldberg AP, et al. Left ventricular diastolic filling performance in older male athletes. *JAMA*. 1995;273:1371–1375.
14. Guirado GN, Damatto RL, Matsubara BB, Roscani MG, Fusco DR, et al. Combined exercise training in asymptomatic elderly with controlled hypertension: effects on functional capacity and cardiac diastolic function. *Med Sci Monit*. 2012;18(7) CR 461-5.
15. Patil SG, Dhanakshirur G, Aithala MR, Kusal Kanti Das. Comparison of the effects of yoga and lifestyle modification on grade-I hypertension in elderly males: a preliminary study. *Int J Clin Exp Physiol*. 2014;1:68–72.
16. Bharshankar JR, Bharshankar RN, Deshpande VN, Kaore SB, et al. Effect of yoga on cardiovascular system in subjects above 40 years. *Indian J Physiol Pharmacol*. 2003;42:202–206.
17. Manchanda SC, Narang R, Reddy KS, Sachdeva U, Prabhakaran D, et al. Retardation of coronary atherosclerosis with yoga lifestyle intervention. *J Assoc Phys India*. 2000;48:687–694.
18. Cramer H, Lauche R, Haller H, Steckhan N, Michalsen A, et al. Effects of yoga on cardiovascular disease risk factors: a systematic review and meta-analysis. *Int J Cardiol*. 2014;173:170–183.
19. The Task force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC) 2007 Guidelines for the management of arterial hypertension. *Eur Heart J* (2007) 28, 1462–1536.
20. Indian Council for Medical Research. ICMR ethical guidelines for biomedical research on human participants. Available at: http://icmr.nic.in/ethical_guidelines.pdf. (Accessed 10 July 2013).
21. Telles S, Reddy SK, Nagendra HR. Oxygen consumption and respiration following two yoga relaxation techniques. *Appl Psychophysiol Biofeedback*. 2000;25(4):221–227.
22. Supiano MA. Hypertension. In: Halter JB, Ouslander JG, Tinetti ME, Studenski S, High KP, Asthana S, eds. *Hazard's geriatric medicine and gerontology*. 6th ed. McGraw Hill Medical publishers; 2009:975–982.
23. Gobel FL, Nordstrom LA, Nelson RR, Jorgensen CR, Wang Y. The Rate pressure product as an index of Myocardial oxygen consumption during exercise in patients with angina pectoris. *Circulation*. 1978;57(3):549–556.
24. Patil SG, Aithala MR, Das KK. Effect of yoga on arterial stiffness in elderly with increased pulse pressure: a randomized controlled study. *Complement Ther Med*. 2015;23(4):562–569.
25. Cooney MT, Vartiainen E, Laatikainen T, Juolevi A, Dudina A, et al. Elevated resting heart rate is an independent risk factor for cardiovascular disease in healthy men and women. *Am Heart J*. 2010;159(4):612–619.
26. Gilman MW, Kannel WB, Belanger A, D'Agostino RB. Influence of heart rate on mortality among persons with hypertension: the Framingham Study. *Am Heart J*. 1993;125(4):1148–1154.
27. Reunanen A, Karjalainen J, Ristola P, Heliövaara M, Knekt P, et al. Heart rate and mortality. *J Intern Med*. 2000;247(2):231–239.
28. Verrier RL, Tan A. Heart rate, autonomic markers and cardiac mortality. *Heart Rhythm*. 2009;6(Suppl. (11)):S68–S75.
29. Park BJ, Lee HR, Shim JY, Lee JH, Jung DH, et al. Association between resting heart rate and arterial stiffness in Korean adults. *Arch Cardiovasc Dis*. 2010;103(4):246–252.
30. Koskela JK, Tahvanainen A, Harring A, Tikkaoski AJ, Ilveskoski E, et al. Association of resting heart rate with cardiovascular function: a cross-sectional study in 522 Finnish subjects. *BMC Cardiovasc Disord*. 2013;13:102.
31. Bangalore S, Steg G, Deedwania P, Crowley K, Eagle KA, et al. Beta-Blocker use and clinical outcomes in stable outpatients with and without coronary artery disease. *JAMA*. 2012;308(13):1340–1349.
32. Santaella DF, Devesa CRS, Rojo MR, Amato MB, Drager LF, et al. Yoga respiratory training improves respiratory function and cardiac sympathovagal balance in elderly subjects: a randomized controlled trial. *BMJ Open*. 2011;1:e000085.
33. Telles S, Reddy SK, Nagendra HR. Oxygen consumption and respiration following two yoga relaxation techniques. *Appl Psychophysiol Biofeedback*. 2000;25(4):221–227.
34. Patil SG, Dhanakshirur GB, Aithala MR, Naregal G, et al. Effect of yoga on oxidative stress in elderly with grade-I hypertension: a randomized controlled study. *J Clin Diagn Res*. 2014;8:BC04–BC07.
35. Nottin S, Nguyen LD, Terbah M, Obert P. Long-term endurance training does not prevent the age-related decrease in left ventricular relaxation properties. *Acta Physiol Scand*. 2004;181(2):209–215.
36. Ehsani AA, Martin 3rd. WH3rd, Heath GW, Coyle EF. Cardiac effects of prolonged and intense exercise training in patients with coronary artery disease. *Am J Cardiol*. 1982;50(2):246–254.