

ZOOPLANKTON DENSITY AND DIVERSITY INDICES IN THE BHAKRA-YAMUNA LINK CANAL SYSTEM IN NARWANA REGION, HARYANA.

Parmila Devi

Department of Zoology, Kurukshetra University, Kurukshetra- 136119, Haryana

Anil K. Tyor

Department of Zoology, Kurukshetra University, Kurukshetra- 136119, Haryana

ABSTRACT

Present study was undertaken to analyse the qualitative and quantitative variations in abundance and diversity indices of zooplankton of Bhakra–Yamuna link canals in the Narwana region, Haryana. For analysis, samples were collected from three selected sites (S_1 , S_2 , and S_3) on Bhakra–Yamuna Link canals for a period of one year from January to December, 2013. For diversity index, Shannon and weaver diversity index, Simpson dominance index and Pielou evenness index were used and to calculate similarity index between selected sites, Jaccard similarity coefficient was used. Result revealed that a total of 13 species of zooplanktons were recorded belonging to group protozoa, rotifera, cladocera and insecta. Population density of zooplankton was low at site S_1 and higher at site S_2 . The diversity indices revealed a significant decline in the population of zooplankton from site S_2 to S_3 after linking with site S_1 . Similarity coefficient was more between S_2 and S_3 as compare to site S_2 and S_1 and site S_1 and S_2 . Cladocerans were dominant at site S_1 and Protozoans were dominant at site S_2 .

KEYWORDS : Zooplankton, Shannon diversity indices, Simpson dominance index, Pielou evenness index

Introduction

Zooplanktons are indispensable members of the aquatic food chain, connecting primary producers to higher trophic levels including economically important population of fish (Umadevi, 2013). They are sensitive to changes in the aquatic environment and any variation in their composition is often a reaction of significant alteration in ambient conditions within aquatic ecosystem. The factors regulating their abundance may be hydrological, chemical, physical and biotic (Ramesha and Sophia, 2013). Some studies on Yamuna river and Western Yamuna Canal have been undertaken by few workers which deals with heavy metal pollution (Jindal and Rumana, 2000; Kaushik *et al.*, 2009), physicochemical parameters (Khairwal *et al.*, 2003, Bhatnagar *et al.*, 2009, Chopra *et al.*, 2012) and biological assessment (Chopra *et al.*, 2011). However there is no information about diversity of zooplankton of Bhakra–Yamuna Link canals in Narwana region. Hence the present study was undertaken to analyse the zooplankton abundance and diversity indices in Bhakra–Yamuna link canals in the Narwana region of Haryana.

Materials and methodology

Study sites: For the present study, three sampling sites (S_1 , S_2 , and S_3) were selected on Bhakra–Yamuna link canal system (Figure 1). Site S_1 (29°63'N & 76°17'E) is located 1 Km upstream from Dhakal head on Sirsa branch of Western Yamuna Canal. Site S_2 (29°61'N & 76°15'E) is located 1 Km upstream from Dhakal head on Barwala link canal and site S_3 (29°59'N & 76°14'E) is located 2.5 km downstream on Sirsa branch after junction with Barwala link canal in Narwana region, Haryana.




Fig. 1: Map of Haryana showing study sites on selected canals.

Plankton Analysis: For plankton study, 50 litres of water was filtered through planktonic net of mesh size 50 μ m with demarcated

tube fitted at the bottom, the concentrated sample preserved in 4% formalin. The literature consulted for the identification of plankton were: Ward and Whipple (1959), Needham and Needham (1962) and Gupta (1972). Counting of plankton was done with the help of “drop count method” and biodiversity indices were calculated by Shannon and weaver (1963), Simpson (1949) and Pielou (1966). Similarity coefficient was calculated by using Jaccard similarity coefficient (Jaccard, 1912).

Result and Discussion

Zooplankton, by their heterotrophic activity, plays a key role in cycling of the organic matter in an aquatic ecosystem. Zooplankton contributed a minor portion of total plankton and comprised of protozoans, rotifers, cladocerans and insects. A total of 13 species were recorded from all study sites. Cladocerans were dominant at site S_1 with percentage distribution of 37.5% of total species diversity. Malhotra, 2014 was also reported the similar results in Western Yamuna Canal in Yamunanagar. Protozoans were dominant at site S_2 with percentage contribution of 50% whereas at site S_3 , their percentage decreases to 33.33% due to the influence of water from site S_1 (Fig. 1). Only two taxa i.e. *Peridinium* sp. and *Polyarthra* sp. were common at all sites (Table 1).

Species richness was found highest at site S_2 (five species) and lowest at site S_3 (one species) (Fig. 2). Zooplankton density in different months ranged from $49 \pm 1.98 \text{ L}^{-1}$ to $18 \pm 2.09 \text{ L}^{-1}$ at site S_1 , $94 \pm 2.38 \text{ L}^{-1}$ to $38 \pm 2.09 \text{ L}^{-1}$ at site S_2 and $86 \pm 4.78 \text{ L}^{-1}$ to $22 \pm 6.31 \text{ L}^{-1}$ at site S_3 . Comparatively, population density of zooplankton was low at site S_1 than site S_2 and S_3 . Bhatnagar *et al.* (2013); Malhotra and Kumar (2014), also observed low values of zooplankton density coupled with low species diversity at WYC. At site S_3 population density was lower from the month of July to September when water current and water level at site S_1 were high which affect their prevalent biota (Fig. 2).

Diversity indices are good indicator of pollution in aquatic ecosystem. Mean value of Shannon diversity index were found maximum at site S_2 (2.77 ± 0.58) and minimum at site S_1 (1.45 ± 0.39). Site S_3 shows decline in the values of diversity as compared to site S_2 but values were greater than site S_1 (Table 2). Simpson diversity index was high at site S_1 (0.33 ± 0.12) than site S_2 and S_3 . Species evenness of zooplankton varied from 0.78 to 0.85 showed little variation at site S_2 as compare to site S_1 (0.60–0.78) and S_3 (0.61–0.84). Evenness was lowest at site S_1 during rainy months. At site S_3 , evenness was low when it received water from site S_1 and when S_1 became dry species evenness approached similar to the site S_2 . Zooplankton tends to show a negative relationship with water flow and a positive relationship with water residence time (Basu and Pick,

1996; Lair, 2006; Strayer *et al.*, 2008). Basu *et al.* (2010) have reported positive correlation of all the physico-chemical parameters except the dissolved oxygen and TDS with zooplankton community.

Similarity coefficient calculated on the basis of zooplankton revealed that site S₂ and S₃ was more similar as compare to site S₂ & S₁ and site S₁ & S₂ throughout the study period (Table 2). This may be due to the fact that site S₂ carried higher volume of water from site S₂ throughout the year and more species were common at these two sites site while site S₁ was dry during some part of the year.

Conclusion: In present study, high values of population density of zooplankton coupled with high species diversity recorded at site S₂ which decreased at site S₃ after joining with site S₁ which altered the overall ecology of the canal.

Table 1: Zooplankton recorded at selected study sites (site S₁, S₂ and S₃) during study period

Sr. No.	Scientific name	Site S ₁	Site S ₂	Site S ₃
I				
Family Protozoa				
1	<i>Alona</i> sp.	+	+	+
2	<i>Ceratium</i> sp.	-	+	-
3	<i>Diffusia</i> sp.	-	++	-
4	<i>Peridinium</i> sp.	+	++	++
II				
Family Rotifera				
5	<i>Brachionus</i> sp.	+	-	+
6	<i>Notholca</i> sp.	-	+	-
7	<i>Polyarthra</i> sp.	-	+	-
8	<i>Trichocerca</i> sp.	-	+	-
III				
Family Cladocera				
9	<i>Daphnia</i> sp.	++	-	+
10	<i>Sida</i> sp.	++	-	-
11	<i>Simocephalus</i> sp.	++	-	-
IV				
Family Insecta				
12	Chironomous larvae	+	-	+
13	Water mite	-	++	+

+ Present, - Absent, ++ Abundant




Fig.2: Per cent distribution of different groups of zooplankton at study sites




Fig. 3: Monthly variation in population density (Nos. L-1) of

zooplankton at study sites.

Table 2: Range and mean ± S.E. (with in parenthesis) of Species richness, species diversity and evenness of zooplanktons reported at selected sites during study period.

Diversity indices/Sites	Site S ₁ *	Site S ₂	Site S ₃
Species richness	2-5	3-7	1-5
Shannon diversity index	1.12-1.92 (1.45±0.39)	2.61-3.01 (2.77±0.58)	1.98-2.85 (2.40±0.31)
Simpson's diversity index	0.28-0.41 (0.33±0.12)	0.16-0.23 (0.18±0.05)	0.17-0.26 (0.21±0.07)
Equitability index	0.60-0.78 (0.68±0.05)	0.78-0.85 (0.81±0.03)	0.61-0.84 (0.72±0.08)
Similarity coefficient	(S ₁ and S ₂) 0.311	(S ₂ and S ₃) 0.798	(S ₃ and S ₁) 0.520

Sampling could not be done from March to May as the canal was completely dry

REFERENCES

- Basu, B. K., and Pick, F. R. (1996). Factors regulating phytoplankton and zooplankton biomass in temperate rivers. *Limnol. Oceanogr.*, 41: 1572-1577.
- Basu, M., Roy, N. and Barik, A. (2010). Seasonal abundance of net zooplankton correlated with physico-chemical parameters in freshwater ecosystem. *Int. J. Lak. Riv.*, 3:67-77.
- Bhatnagar, A., Chopra, G. and Malhotra, P. (2009). Water quality indices and abiotic characteristics of Western Yamuna Canal in Yamunanagar, Haryana, India. *J. Nat and App Sci.*, 1(2): 149-154.
- Bhatnagar, A., Chopra, G. and Malhotra, P. (2013). Assessment of water quality of river Yamuna in Yamunanagar, India with reference to planktons and macrozoobenthos. *Sch. J. Eng. Tech.*, 1(4): 204-213.
- Chopra G., Bhatnagar A. and Malhotra P. (2012). Limnochemical characteristics of river Yamuna in Yamunanagar, Haryana, India, *Inter j. water res and environ eng.*, 4: 97-104.
- Gupta, R.S. (1972). Blue green algal flora of Rajasthan. *Nova Hedwigia*, 23:481-492.
- Jaccard, P. (1912). The distribution of the Flora of alpine Zone. *New Phytol.* 11: 37-50.
- Jindal, R. and Rumana, H.S. (2000). Biomonitoring of water pollution in Western Yamuna Canal at Yamunanagar, Haryana. *JPAS.* 2(1): 177-182.
- Khaiwal, R., Ameena, Meenakshi, Rani, M. and Kaushik, A. (2003). Seasonal variations in physico-chemical characteristics of river Yamuna in Haryana and its ecological best designated use. *Journal of Environment Monitoring.* 5: 419-426.
- Kaushik, A., Kansal, A., Meena, S., Kumari, S., and Kaushik, C.P. (2009). Heavy metal contamination of river Yamuna, Haryana, India: Assessment by metal enrichment factor of the sediments. *Journal of Hazardous Material.* 164: 265-270.
- Lair, N., (2006). Abiotic vs biotic factors: lessons drawn from rotifers in the Middle Loire, a meandering river monitored from 1995 to 2002, during low flow periods. *Hydrobiol.*, 546:457-472.
- Malhotra, P. (2014). Species diversity and distribution of zooplankton of Western Yamuna Canal in Yamunanagar (Haryana) with special reference to industrial pollution. *Int. Res. J. Environ. Sci.*, 3(8):1-4.
- Malhotra, P. and Kumar, A. (2014). Comparative studies on zooplanktonic diversity of River Yamuna and Western Yamuna Canal in relation to industrial pollution in Yamunanagar (Haryana), India. *Int. J. Sci. Res.*, 3(9): 1438-1441.
- Needham, J.E. and Needham, P.R. (1972). *A guide to the study of fresh water biology.* Holden Day Inc. San Francisco, California.
- Pielou, E.C. (1966). The measurement of diversity in different types of biological collections. *J. Theor. Biol.*, 13: 131-144.
- Ramesha M.M. and Sophia, S. (2013) Species Composition and Diversity of Plankton in the River Seeta at Seetanadi, the Western Ghats, India, *Adv. Bio Tech.*, 12(08): 20-27.
- Shannon, C.E. and Weaver, W. (1963). The mathematical theory of communication, University of Illinois, Press, Urbana, pp117.
- Simpson, G.H. (1949). Measurement of diversity. *Nature*, 163: 688.
- Strayer, D.L., Pace, M.L., Caraco, N.F., Cole, J.J. and Findlay, S.E.G. (2008). Hydrology and grazing jointly control a large-river food web. *Ecology*, 89: 12-18.
- Umadevi, T. (2013). Limnological investigation and zooplankton diversity of Karanjriver, Karnataka. *International Journal of Science and Research.* 2(3):133-136.
- Ward, G.B. and Whipple, G.C. (1959). *Freshwater Biology*, John Willey and Sons, New York, pp 1248.