



## ENDOMETRIAL PATTERN IN MENOPAUSE TRANSITION AGE GROUP PATIENTS WITH ABNORMAL UTERINE BLEEDING.

**Vasant Kawade\***

Professor and Head, Department of OBGY, B.K.L.Walawakar Rural Medical College and Hospital, Sawarde \*Corresponding Author

**Pawar Samir**

Consultant Gynecologist, AK Pawar Institute of Gynec and Obstetrics, Nashik

### ABSTRACT

**Introduction:** In both, benign and malignant gynaecological diseases Abnormal Uterine Bleeding (AUB) is an important symptom. It is associated with endometrial carcinoma in the menopause transition age group, ie perimenopausal and postmenopausal, in approximately 10% of cases. The present study was designed to study the histopathology pattern of the endometrium in women with abnormal perimenopausal and postmenopausal uterine bleeding. The endometrium was obtained by endometrial biopsy, the menopause transition age group.

**Aims and Objectives:** To study, the various histopathological patterns of endometrium, in menopause transition age group patients (those are in perimenopausal and postmenopausal age group) with abnormal uterine bleeding.

**Materials and Methods:** 100 patients were selected for the study after satisfying the inclusion and exclusion criteria. They underwent detailed history taking including the pattern of bleeding, general, systemic, and pelvic examination. They were subjected to routine laboratory investigations and pelvic ultrasound examination. Endometrial biopsy was done in all the patients. Endometrial biopsy was done either in the outpatient without anaesthesia or inpatient under short general anaesthesia depending on the patients general condition. Biopsy was performed using a manual vacuum aspiration syringe, and the material collected was sent for histopathological examination. Endometrium was also obtained from patients undergoing therapeutic dilatation and curettage. Analysis of Histopathology report was done and results were obtained.

**Results:** Most predominant findings of histopathological examination were Proliferative Endometrium in 29% Simple hyperplasia without atypia 28% Secretory 20% Followed by other patterns.

**Conclusion:** Study of endometrial pattern in menopausal age group, ie perimenopausal and postmenopausal women with abnormal uterine bleeding is helpful to diagnose hyperplasia and carcinoma of endometrium.

**KEYWORDS :** Postmenopausal Uterine Bleeding, Simple Hyperplasia without atypia, Secretory.

### Introduction:

Perimenopause is the phase, preceding the onset of menopause, generally occurring around 40 years of age during which the regular menstrual cycle of a woman changes from normal regular cycles to a pattern of irregular cycles(1,2). Menopausal transition includes a period of about 4–5 years before menopause, sometimes even several months, characterized by varying degrees of somatic and psychological changes that reflect the change in the ovarian cycle (3). In some women, the most significant symptom is an irregular menstrual pattern, which must be carefully evaluated to determine whether it is the consequence of low oestrogen levels or an associated pathology (4). The terminology of abnormal uterine bleeding includes the following clinical entities (2,5,6) Oligomenorrhea, Polymenorrhea, Menorrhagia, Menometrorrhagia, Metrorrhagia, Midcycle spotting, and dysfunctional uterine bleeding.

Abnormal Uterine Bleeding (AUB) is an important symptom of both benign and malignant gynaecological diseases. AUB is the single most common reason for gynaecological referrals (7). Excessive menstrual blood loss affects 10–30% of menstruating women and in the order of 70% of all gynaecological consultations in the Perimenopause and Postmenopause (8). Abnormal perimenopausal or postmenopausal bleeding is associated with endometrial carcinoma in approximately 10% of cases (9,10). In particular, postmenopausal bleeding may be an early symptom of endometrial carcinoma. Currently the most commonly used technologies for outpatient evaluation of the endometrium are biopsy, hysteroscopy, and Trans Vaginal Ultrasound (TVS). Endometrial biopsy by Dilatation and Curettage or office endometrial biopsy is considered the gold standard in AUB. The present study is designed to study the histopathological results of the endometrial biopsy in women with abnormal perimenopausal and postmenopausal uterine bleeding.

### Aims and Objectives:

To study, the endometrial histopathological patterns, in patients with abnormal uterine bleeding, those are in perimenopausal and

postmenopausal age group.

### Materials and Methods:

The present study "Endometrial pattern in Menopausal age (in perimenopausal and postmenopausal) patients with abnormal uterine bleeding patients" was a prospective study. It was carried out at 'Dr. AK Pawar Institute of Obstetrics and Gynaecology, Nashik'. The period of study was from April 2012 to July 2015. 100 patients were selected for the study after satisfying the following inclusion and exclusion criteria.

**Inclusion Criteria-** Women with abnormal uterine bleeding more than 40 years of age.

**Exclusion Criteria** –i) Hormone therapy in last 6 months. ii) Positive pregnancy test. iii) Cases with cervical, uterine, or adnexal pathology on clinical examination or ultrasound. iv) Women with cervical cancer. v) History/ evidence suggestive of active pelvic infection.

Written Informed consent was obtained from all participants. A structured proforma was used to collect the medical history and examination findings of the patients. They underwent detailed history taking including the pattern of bleeding, and general, systemic, and pelvic examination. They were subjected to routine laboratory investigations and pelvic ultrasound examination. Endometrial biopsy was done in all the patients. Endometrial biopsy was done either in the outpatient without anaesthesia or inpatient under short general anaesthesia depending on the patients general condition. Biopsy was performed using a manual vacuum aspiration syringe, and the material collected was sent for histopathological examination. Endometrium was also obtained from patients undergoing therapeutic dilatation and curettage. Analysis of Histopathology report was done and results were obtained.

### Results:

There were total 100 cases of abnormal uterine bleeding in

perimenopausal and postmenopausal patients. Out of these, 83 (83%) were perimenopausal patients and 17 (17%) were postmenopausal patients. The clinical characteristics of the study population and age distribution are shown in Tables 1 and 2. The maximum incidence of abnormal uterine bleeding of 85% was in

multiparous patients (parity 1-3), followed by 11% in grandmultiparous (parity>3) patients and least 4% in nulliparous patients. Distribution of patients according to bleeding pattern is as shown in Table 3

**Table 1. Age-wise distribution of patients**

Age	Perimenopausal (% in group)	Postmenopausal (% in group)	Total (%)
40-45	60 (72%)	1 (6%)	61 (61%)
46-50	18 (22%)	4 (24%)	22 (22%)
51-55	4 (5%)	4 (24%)	8 (8%)
56-60	1 (1%)	5 (29%)	6 (6%)
61-65	0 (0%)	2 (12%)	2 (2%)
66-70	0 (0%)	1 (6%)	1 (1%)
Total	83 (100%)	17 (100%)	100 (100%)

**Table 2. The clinical characteristics of the study population**

	Perimenopausal		Postmenopausal		Total (Perimenopausal + Postmenopausal)	
	Range	Mean ±SD	Range	Mean ±SD	Range	Mean ±SD
Age (years)	40-56	44.012 ± 3.71	45-68	55.00 ± 6.10	40 - 68	45.880 ± 5.88
Menarche (years)	11-16	13.397 ± 1.27	11-16	13.470 ± 1.23	11 - 16	13.410 ± 1.26
Menopause (years)	-	-	43-55	48.058 ± 3.32	43-55	48.058 ± 3.32
BMI (kg/m <sup>2</sup> )	18-35	23.720 ± 3.70	18-39	26.147 ± 5.51	18 -39	24.133 ± 4.13
Haemoglobin (g/dl)	5.5-13.3	10.200 ± 1.65	8.6-13.2	11.764 ± 1.11	5.5-13.3	10.466 ± 1.67
ET (mm) on TVS	4-29	13.388 ± 4.36	3.0-18.0	11.147 ± 5.09	3-29	13.007 ± 4.55

**Table 3. Distribution of patients according to histopathology of endometrium**

Histopathology	Perimenopausal (% in group)	Postmenopausal (% in group)	Total (%)
Proliferative	27 (33%)	2 (12%)	29 (29%)
Secretory	19 (23%)	1 (6%)	20 (20%)
Simple hyperplasia without atypia	22 (27%)	6 (35%)	28 (28%)
Simple hyperplasia with atypia	6 (7%)	0 (0%)	6 (6%)
Complex hyperplasia without atypia	1(1%)	1 (6%)	2 (2%)
Complex hyperplasia with atypia	2(2%)	1(6%)	3 (3%)
Atrophic	1 (1%)	4 (24%)	5 (5%)
Disordered proliferation	5 (6%)	0 (0%)	5 (5%)
Endometrial carcinoma	0 (0%)	2 (12%)	2 (2%)
Total	83(100%)	17(100%)	100(100%)

**Discussion:**

Abnormal uterine bleeding at any stage in a woman's life is disruptive and worrisome. But postmenopausal bleeding is of special concern because it is the only common clinical indication of the presence of endometrial Carcinoma. Several different approaches have been proved to be clinically useful screening methods for early detection of endometrial abnormality in women with irregular uterine bleeding. These include dilatation and curettage (D&C), hysteroscopy, and micro-hysteroscopy. With regards to age of the study population, Archana Bhosle(1) studied 112 perimenopausal women with abnormal uterine bleeding, where 76% were in the age group of 41-45, 2.6% were in group 46-50 and 2.6% in group >51. Cornitescu(3) studied 256 perimenopausal patients and reported 35.5% incidence in age 41-45 and 64.5% incidence in group 46-52. Bharti et al. (12) found in perimenopausal patients the mean age is 47.52 ± 1.88 (43-50) whereas in postmenopausal patients it is 55.254 ± 3.84 (52-65). Indu Kaul et al (11) studied postmenopausal patients and in their study the mean age is 54.06 ± 6.64 (41-70). According to Kaul et al (11), the mean age at menopause is 47.8 ± 2.82 (range 47-49), whereas according to our study it is 48.05 ± 3.32 years. With regards to the pattern of bleeding, (1) the maximum incidence was of menorrhagia (53.3%), followed by 28.2%, 12.2% and 6.5% with polymenorrhagia, intermenstrual bleeding and metrorrhagia respectively. Majority of patients, ie 34% have menometrorrhagia , while least incidence , ie 9% is of intermenstrual bleeding (3) . With regards to the histopathology of endometrium in perimenopausal patients 4.3% have proliferative endometrium, 78.26% have simple hyperplasia without atypia and 13.04% have complex hyperplasia without atypia (1). According to Bhosle (1) , 66.1% had proliferative endometrium, 16.1% had Secretory endometrium and 17.8% simple hyperplasia without

atypia. Cornitescu(3) found , 8.6% had Secretory endometrium, 1.2% had simple hyperplasia without atypia, 1.6% had complex hyperplasia with atypia, 7% had atrophic endometrium and 1.9% had disordered proliferation. Among postmenopausal patients, 56% had simple hyperplasia without atypia, 8% had complex hyperplasia without atypia, 8% had complex hyperplasia with atypia, 16% had atrophic endometrium and 12% had carcinoma in a study by Bharati et al (12). Kaul et al (11), concluded ,18% had hyperplasias, 50% had atrophic endometrium, 10% had carcinomas, 8% had polyp, 8% had insufficient sample, 4% had endometritis and 10% had hormonal effect. Endometrial carcinoma most often occurs in women in the sixth and seventh decades of life at an average age of 60 years. Seventy five percent cases occur in women older than 50 years of age and 90% of women with endometrial carcinoma have vaginal bleeding or discharge as their only presenting complaint. Less than 5% of women diagnosed with endometrial cancer are asymptomatic. The incidence of adenocarcinoma is 0.1% per year in post menopausal women but rises to 10% in presence of abnormal bleeding. In our study there were 2 cases of adenocarcinoma amongst 17 patients with post menopausal bleeding and none in perimenopausal group.

**Conclusion:**

Study of endometrial histopathology in women in Menopau transition period, ie perimenopausal and postmenopausal ages, with abnormal uterine bleeding is helpful to diagnose hyperplasia and carcinoma of endometrium. Endometrial biopsy has for many years been the methods of choice for the diagnosis of endometrial cancer in patients with peri and postmenopausal bleeding. It also reveals various endometrial patterns ranging from proliferative, secretory, simple and complex hyperplasia with or without atypia,

disordered proliferation and atrophic endometrium.

**Conflict of Interest:**

The authors declare, they have no conflict of interest.

**Acknowledgements:**

We thank the management and administration of AK Pawar Institute of Obstetrics and Gynaecology, Nashik (AKPI) for providing the facilities for the management of these patients.

**References:**

1. Bhosle A, Fonseca M. Evaluation and histopathological correlation of abnormal uterine bleeding in perimenopausal women. *Bombay Hospital J.* 2010;52(1):69-72.
2. Fritz MA, Speroff L. *Clinical gynaecologic endocrinology, and infertility.* 8th ed. Lippincott Williams & Wilkins; 2011.
3. Cornitescu FI, Tanase F, Simionescu C, Iliescu D. Clinical, histopathological, and therapeutic considerations in non-neoplastic abnormal uterine bleeding in menopause transition. *Rom J Morphol Embryol.* 2011;52(3):759-65.
4. Berek JS. *Berek's & Novak's Gynaecology.* 14th ed. Philadelphia: Lippincott Williams & Wilkins; 2007.
5. Goodman A. Abnormal genital tract bleeding. *Clin Cornerstone.* 2000;3(1):25-35.
6. Goodman A. Abnormal genital tract bleeding. *Clin Cornerstone.* 2000;3(1):25-35
6. Oriol KA, Schrager S. Abnormal uterine bleeding. *Am Fam Physician.* 1999;60(5):1371-82.
7. Gimpelson RJ, Rappold HO. A comparative study between panoramic hysteroscopy with directed biopsies and dilatation and curettage. *Am J Obstet Gynecol.* 1988;158:489-92.
8. Clark TJ, Khan KS, Gupta JK. The diagnosis of intrauterine pathology in postmenopausal women: an evidence-based approach. *Rev Gynaecol Pract.* 2002;2:109-16.
9. Creasman WT. Endometrial cancer: incidence, prognostic factors, diagnosis, and treatment. *Semin Oncol.* 1997;24:S1-140.
10. Smith-Bindman R, Kerlikowske K, Feldstein VA, Subak L, Scheidler J, Segal M. Endovaginal ultrasound to exclude endometrial cancer and other endometrial abnormalities. *JAMA.* 1998;280:1510-17.
11. Kaul I, Kalsi M, Anand AK, Jad R, Menia V. Transvaginal sonography versus Histopathology in Postmenopausal Bleeding: a prospective study. *JK Science.* 2012 Jul-Sep;14(3):129-33.
12. Bharti B, Satish R P. Feasibility, and yield of endometrial biopsy using suction curette device for evaluation of abnormal pre and postmenopausal bleeding. *J Obstet Gynecol India.* 2008 Jul/Aug;58(4),322-326.