



A CROSS SECTIONAL STUDY FOR SCREENING OF THE RISK FACTORS OF DIABETES AMONG RURAL POPULATION

Dr. Sushma Debuka

Asst Prof, Department of Physiology, Raipur Institute of Medical Sciences, Raipur, CG, India

Dr. Rahul Kewal Kumar*

Asst Prof., Department of Community Medicine, Raipur Institute of Medical Sciences, Raipur, CG, India *Corresponding Author

ABSTRACT

Background: The rise in diabetes mellitus (DM) has made it a major public health problem & a disaster of major proportion of Populations in India . This scenario becomes worse in rural setup where limited health care facilities are available. So, to detect risk of diabetes the determinants of type-2 DM in people who are ≥ 45 years of age a study was done by selective screening methodology in rural area of Arang Block of Raipur district

Methods: A cross-sectional population-based study was conducted in Arang block of Raipur District during 22nd April to 30th May 2019 among those who are at risk of developing DM, as per the WHO guidelines on Laboratory Diagnosis and monitoring of Diabetes Mellitus . Blood glucose estimation was done using a blood glucose meter. Personal Interview, Anthropometric measurements & Clinical Examination was done and a Pre-designed Questionnaire form was filled from every study subject. Analyzed on SPSS.

Results: 8.4 % of the persons above the age of 45 years were diabetic. Study also revealed the proportion of people diagnosed with DM increases with increasing age groups. The difference between the post meal blood glucose level among the diabetics and nondiabetics was statistically significant. The proportion of diabetics was more among those who had family history of diabetes , BMI more than 25 and those with sedentary lifestyle.

Conclusions: Implementation of preventive measures to reduce the burden of diabetes is needed. Identification of the various factors adversely related to glucose intolerance will help in evolving preventive strategies.

KEYWORDS : Cement Workers , Respiratory Functions , Occupational Hazards , Cement Dust

INTRODUCTION

Prevalence of diabetes mellitus (DM) and impaired fasting glucose are reported to be highly variable among rural and urban population in India. A number of epidemiological studies with varying sample sizes have reported prevalence of diabetes at different geographical areas ranging from 1.6–12.4%,¹ type-2 diabetes is a global health problem.² According to the recent global estimates of the World Health Organization (WHO), there will be 300 million people with diabetes by the year 2025.^{2,3} It is estimated that the developing countries will bear the brunt of diabetes epidemic to the extent of 77% of the global burden, in the 21st century.⁴ Number of diabetics in India was 31.7 million in the year 2002⁵ and it is estimated that number of diabetics in 2030 will be 79.4 million in India.^{6,7}

The prevalence of diabetes in Indian adults was found to be 2.4% in rural and 4–11.6% in urban dwellers.⁵ Today, diabetes no longer remains a disease of the high socioeconomic status or confined to urban area. However, not enough prevalence studies have been conducted in India for rural population. American Diabetes Association has proposed the screening of all patients aged over 45 years by measuring fasting blood glucose every three years, in addition to screening patients from high-risk groups and younger patients with hypertension, obesity, a family history of diabetes in a first-degree relative, or a family history of gestational diabetes.⁷ Considering the above facts, this study was undertaken to study the distribution of determinants of type-2 diabetes among rural population at risk of developing DM in Arang Block Raipur District, C.G.

MATERIALS & METHODS

Ethical Consideration- This study plan has got Ethical clearance from the Institutional Ethical Committee .Informed Written Consent was taken from each participating subjects and in case of any illiterate subjects this consent was read out to him/her.

METHODOLOGY

This study was undertaken in Godhi Village of Arang Tehsil Raipur District .

A cross-sectional population-based study was conducted among those who are at risk of developing DM as per the WHO guidelines on Laboratory Diagnosis and monitoring of Diabetes Mellitus.⁸

Sample Size - The final sample included 306 individuals, with 171 men and 135 women, after selective screening methodology.

Inclusion criteria

Selective screening methodology was used to detect those who are at risk for diabetes. Criteria for selective screening for diabetes for the individuals as per WHO guidelines included⁹

- typical symptoms of diabetes,
- first-degree relatives with diabetes,
- overweight (BMI ≥ 25 kg/m²),
- women who had delivered baby weighing ≥ 4.5 kg or had gestational DM,
- hypertensive ($\geq 140/90$ mm Hg),
- raised serum triglyceride and cholesterol levels and
- prior history of IGT or IFG

Exclusion criteria

- known cases of type-2 diabetes,
- individuals who declined for informed consent, and
- not available at home even after repeat second visit.

Sampling

To select the household, systematic random sampling was done. A list of the households was made in all the selected wards. Thereafter, a random number was chosen by taking the last digit of the currency note and that was selected as the first household, and subsequently, every alternate household was selected. Similar procedure was followed for all the wards. The final sample included 306 individuals, with 171 men and 135 women, after selective screening methodology.

Data collection

On visiting the selected house, the investigator found out if there are any members aged ≥ 45 years of age, having any of

the above mentioned risk factors. On affirmation, the participant was included in the study after the written informed consent.

Data were collected using a Predesigned Pretested structured interview schedule. During the interview, information about sociodemographic status, family history of DM, physical activity, and other known risk factors for diabetes were asked. After the interview, participants were informed about procedural details of the blood sugar investigation. Participants were motivated to fast overnight and its importance was stressed for correct blood sugar recording. On the next day, the healthcare worker and laboratory technician visited their houses for sample collection and test for the fasting blood glucose by Glucometer. Thereafter, 75 gm powdered glucose in a glass of water was given to participants to estimate the postprandial blood sugar after one hour. The participants were informed about the result of the blood test and necessary advice was given and referral was done to a tertiary health care centre for further management, if needed.

Blood glucose estimation was done using Glucometer (One Touch Select Simple, Johnson & Johnson). Two microliters of the sample was collected by sip-in technique. Test time was 30 sec, test interval g/L- 0.4- 0.5). During the field survey, all the instruments were calibrated each morning using standard solutions and were checked with check-strips after every 20 measurements. All control values were within recommended ranges. The Diabetes Control and Complications Trial (DCCT) clearly demonstrated the benefits of normal or near-normal blood glucose levels by using Glucometer and the instrument fulfils the defined essential requirements as per standardizing organizations (ISO, CEN).^{6,10}

Data were analysed and presented as simple percentages and proportions. Test of significance was applied wherever necessary. SPSS version 20 was used.

Operational definitions

Hypertension: as per JSM VII criteria.⁸

Obesity: BMI: Normal, 18.5–24.9 kg/m²; overweight, 25–29.9 kg/m²; obese, ≥30.0 kg/m².

Diabetes: Criteria for the diagnosis of diabetes, proposed by WHO.⁹

RESULTS

Of the 306 persons of more than 45 years of age and who had at least one or more risk factors for DM, as per WHO guidelines,⁹ participated in the study. All belonged to Hindu religion. Twenty six (8.5%), of the 306 participants were diagnosed as diabetics at the time of the survey, mean fasting and post meal blood sugar level (in mg%) of all the study participants were 81.7 (± 3.2) and 118.6 (± 2.5), respectively

The mean fasting and post meal blood sugar level (in mg%) among nondiabetics were 81.7 ± 3.2 and 118.6 ± 2.5 and among diabetics were 138.8 ± 2.1 and 220.7 ± 1.9, respectively. The difference in the post meal blood glucose level among diabetics and non-diabetics was found to be significant. The current study also revealed that the proportion of people diagnosed with DM increases with increasing age groups. Out of 51 participants aged >60 years, 11.8% were diabetic as compared to 9.2% and 6.4% in the age group of 51–60 years and 45–50 years, respectively.

Of the 26 diabetics, 15 were men and 11 were women. The gender-wise distribution of diabetes among study subjects was also not statistically significant ($P = 0.89$). Ten (38.4%) of 26 diabetics gave family history of diabetes and almost

similar numbers of diabetics were not aware of their family history. Twenty three percent diabetic participants did not have family history of diabetes.

The study also revealed that as the BMI increases, the proportion of diabetes also increases. Twenty four percent participants with BMI >25 were diabetics as compared to 6.5 and 1.9 among those with BMI of 18.5–24.9 and <18.5, respectively.

Nineteen (8.5%) of the 255 subjects with hypertension of various stages (as per JSM VII criteria) were diabetic and seven (8.6%) of the 81 were normotensive. The association of hypertension and type-2 DM in our study was not statistically significant.

Eleven (9%) of the 122 participants with history of alcoholism and smoking were diabetics as compared to four out of those who did not give history of any addiction.

DISCUSSION

Of the 306 individuals enrolled in the study, 26 (8.4%) were diabetic, almost similar to the study conducted in Turkey (7.2%).¹⁰ However, the prevalence is lower than that of rural population of Pakistan (11.1%) and Hawaii (20.4%),^{11,12} but higher than in Mongolia (2.9%).¹³

In this study, even though more males, as compared to females, were diabetic, the sex distribution among diabetics was not statistically significant ($P > 0.05$). A study conducted in Turkey and Pakistan has found high prevalence of type-2 DM in women.^{12,13} Most of the diabetics were in the age group of 50–60 years, this finding was almost similar to the study conducted by Ramachandaran in rural South India, wherein 9.9% out of 588 subjects of above 60 years of age had diabetes.¹¹

Most of the studies observed that, regardless of ethnicity, this metabolic disease had increased with economic development related to affluent lifestyle, excess calorie intake and less physical activities resulting from the embrace of a more modernized lifestyle in favor of a traditional lifestyle.¹⁵⁻¹⁸ The preponderance of diabetes in the rural rich indicates the same.

In rural India, developmental changes have influenced the lifestyle of rural people. Therefore, higher prevalence of type-2 DM in study population may be primarily due to environmental factors, apart from the genetic predisposition.^{19,20}

The study revealed that family history of diabetes, increasing age and BMI (obesity) are few important risk factors found to be more among diabetics as compared to nondiabetics. For type-2 diabetes, this is not an unusual finding.^{11,20,21}

CONCLUSION

Our study supports high risk Screening method for diabetes at mass level as it is cost effective as well as time saving procedure. In rural areas, a great number of diabetic cases remain left behind due to lack of investigation facilities. Therefore, all of those could be detected very easily by applying this method

ACKNOWLEDGEMENT

We would like to thank Dr.S.P Singh, Professor & Head, Dept. of Community Medicine, RIMS Raipur for his always available guidance to us.

Declarations

Funding: None

Conflict of interest: None

Ethical approval: Yes

REFERENCES

1. Sadikot SM, Nigam A, Das S, Bajaj S, Zargar AH, Prasannakumar KM, et al. The burden of diabetes and impaired fasting glucose in India using the ADA 1997 criteria: PODIS. *Diabetes Res Clin Prac.* 2004;66:293-300.
2. King H, Rewers M WHO AdHoc Diabetes Reporting Group. Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults. *Diabetes Care.* 1993;16:157-77.
3. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995-2025. *Diabetes Care.* 1998;21:1414-31
4. World Health Organization. *World Diabetes: A Newsletter.* 1997. Sep, pp. 3-6.
5. Available from: http://www.neoncarrot.co.uk/h_aboutindia/india_health_stats.html. (Cited on 22 April 2019)
6. WHO. 1998. Prevention and control of Diabetes Mellitus, Report of an Intercountry workshop, Dhaka, Bangladesh, 27-30 April 1998, SEA/NCD/40.
7. The expert committee on the diagnosis and classification of diabetes mellitus. Report of the expert committee on the diagnosis and classification of diabetes mellitus *Diabetes Care.* 1997;20:1183-97.
8. World Health Organization - Laboratory diagnosis and monitoring of Diabetes Mellitus. 2002:17.
9. WHO. 1985. Technical Report Series; No. 727; 1985.
10. Satman I, Yilmaz T, Sengul A, Salman S, Salman F, Uygur S, et al. Population-based study of diabetes and risk characteristics in Turkey: Results of the Turkish Diabetes Epidemiology Study (TURDEP) *Diabetes Care.* 2002;25:1551-6.
11. Ramachandran A, Snehalatha C, Shyamala P, Vijay V, Viswanathan M. High prevalence of NIDDM and IGT in an elderly south Indian population with low rates of obesity. *Diabetes Care.* 1994;17:1190-2.
12. Shera AS, Rafique G, Khwaja IA, Baqai S, Khan IA, King H. Pakistan National Diabetes Survey prevalence of glucose intolerance and associated factors in North West at Frontier Province (NWFP) of Pakistan. *J Pak Med Assoc.* 1999;49:206-11.
13. Suvd J, Gerel B, Otgooloi H, Purevsuren D, Zolzaya H, Roglic G, et al. Glucose intolerance and associated factors in Mongolia: Results of a national survey. *Diabetes Med.* 2002;19:502-8.
14. McKeigue PM, Bela Shah, Marmot MG. Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. *Lancet.* 1991;337:382-6.
15. Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, et al. Effect of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. *Diabetes Care.* 1997;20:537-44.
16. Astrand PO. Physical activity and fitness: physical activities and carbohydrate metabolism. *Am J Clin Nutr.* 1992;55:1231s-65.
17. Lovejoy J, DiGirolamo M. Habitual dietary intake and insulin sensitivity in lean and obese adults. *Am J Clin Nutr.* 1992;55:1174-9.
18. O'Dea K, Spargo RM, Nestel PJ. Impact of Westernization on carbohydrate and lipid metabolism in Australian aborigines. *Diabetologia.* 1982;22:148-53.
19. McKeigue PM, Miller GJ, Marmot MG. Coronary heart disease in South Asians overseas: a review. *J Clin Epidemiol.* 1989;42:597-609.
20. McKeigue PM, Pierpoint T, Ferrie JE, Marmot MG. Relationship of glucose intolerance and hyperinsulinemia to body fat pattern in South Asians and Europeans. *Diabetologia.* 1992;35:785-91.
21. McKeigue PM, Marmot MG, Syndercombe Court YD, Cottier DE, Rahman S, Riemersma RA. Diabetes hyperinsulinemia and coronary risk factors in Bangladeshis in East London. *Br Heart J.* 1988;60:390-6.