



## A STUDY OF TRANSPOSITION OF REVERSE SURAL FLAPS AND SOLEUS MUSCLE FLAPS IN CASES OF TRAUMATIC LOWER LIMB FRACTURES WITH LARGE SOFT TISSUE DEFECTS

**Dr. Anand kumar D**

Plastic Reconstructive and Aesthetic Surgeon TNMC & BYL Nair Ch. Hospital, Mumbai

**Dr. Gujrathi Chanchal\***

MD Biochemistry, Mumbai. \*Corresponding Author

### ABSTRACT

This study aims to study the effects of transposition of reverse sural flaps and soleus muscle flaps in cases of traumatic lower limb fractures with large soft tissue defects.

**Methods:** It's a record based study in which the data of patients undergoing transposition of the soleus muscle for treating exposed bone in the leg and simultaneous sural flap were retrospectively analyzed from January 2016 to December 2018 in a tertiary hospital in western India.

**Results:** During the duration of study a total of 22 patients with varying ages between 19 and 59 years (18 males and 4 females) were enrolled. The main initial injury was trauma (85%), consisting of tibia and/or fibula fractures.

**Conclusion:** The use of soleus and reverse sural muscle flaps for the treatment of bone exposure gave satisfactory results in covering of exposed structures, favoring local vascularization and improving the initial injury.

**KEYWORDS :** reverse sural flaps, soleus flaps.

### INTRODUCTION:

Plastic surgeons since ages have been troubled treating large tissue defects in cases of lower limb fractures especially of the middle and distal parts of lower limbs. The main goals of soft tissue reconstruction is to cover exposed bone, tendon, or hardware. Recently, many studies have shown no difference in efficacy between musculocutaneous and fasciocutaneous flaps in the treatment of lower limb bone defects. (1) Delayed distally-based fasciocutaneous reverse sural flaps were used in a 2-step procedure (2). A reverse sural flap can be used to effectively treat chronic osteomyelitis without significant donor site morbidity or the complications that may occur during the elevation of a muscle flap (3) Small and moderate-size wounds of the leg may be better served by the use of local and regional flaps, (4) but in large defects, these flaps are not available and more complex and different flap designs are required.

The main goals of soft tissue reconstruction is to cover exposed bone, tendon, or hardware. Small and moderate-size wounds of the leg may be better served by the use of local and regional flaps, (4) but in large defects, these flaps are not available and more complex and different flap designs are required. (5) This study focused on authors experiences when applying simultaneous sural and soleus flaps from one incision for soft tissue reconstruction in large soft tissue defect of the leg.

### MATERIALS AND METHODS:

A record based retrospective study was carried out in the department of plastic surgery in a tertiary teaching hospital from 1<sup>st</sup> January 2016 to 31<sup>st</sup> december 2018. During the period of 2 years, a total of 22 patients (18 males and 4 females) with large soft tissue defects in anterolateral of the leg were studied on which this procedure has been used. The patients' age ranged from 19 to 59 years at the time of operation. Before operation, local soft tissue of lower limb around the defect was assessed carefully. Intact patients were selected for this method of surgery. Bone condition was evaluated and if needed, it was stabilized before soft tissue coverage. Surgical repair was performed under general anesthesia.

The method involved was marking the sural flap with maximum soft tissue that we can harvest. Total belly of soleus and reverse sural were approached from longitudinal lateral incision. The surgery was started by observing the antiseptic and aseptic techniques, with preparation of the entire lower limb in which the muscle transposition would take place, and also the contralateral thigh, when we wanted to use skin graft in the same procedure. After lifting limb for a few minutes, ischemia was made using a crepe

bandage or Smarch tape, by inflating the cuff with a mean pressure of 100 mmHg above the blood systolic pressure.

### RESULTS:

In this study, all 22 patients were managed initially by the orthopedic trauma services. Their wounds were debrided by the primary service, then referred to the plastic surgery department. An open tibial wound in the junction of the middle and distal thirds of the leg was reconstructed successfully with the synchronous regional muscle and fasciocutaneous flaps. All 22 patients had primary healing of their tibial wounds were without any complications.

Only one patient developed insignificant distal flap necrosis and was treated subsequently with surgical debridement and flap readvancement. His tibial wound healed completely after reoperation. All patients had reliable healing of their tibial wounds and evidenced healing of their fractures, with good cosmetic outcome following flap reconstructions. Limb salvage in those patients was also achieved during follow-up. Patients were able to ambulate postoperatively as instructed by the physiotherapist.

### DISCUSSION:

Large tissue defect repair in the lower limbs especially on the middle and distal parts have been a matter of concern for the plastic surgeons since ages and also for the relatives of the patient. There have been surgeries using reverse sural flap for cases of osteomyelitis and good results were obtained (3). During the last 2 decades since free tissue transfer became a standard surgical procedure for limb salvage, the outcome for surgical management of complex lower extremity wounds has improved dramatically. (4-8) Although microsurgical flaps have been the method of choice for this reconstruction, many hospitals do not have equipment or microsurgical staff trained for this type of procedure and also an experienced surgeon is not available. Moreover, the patient's clinical condition does not allow a more complex surgery in some cases.

The management of the mid-third tibial wound can be challenging. If the defect is not too extensive, a soleus muscle flap such as medial hemisoleus muscle flap can be used successfully to cover such a wound. (9) However, if the soleus muscle is traumatized, a microvascular free tissue transfer should be used to cover such a wound. In addition, a combined medial gastrocnemius and medial hemisoleus muscle flaps can also be used to cover a relatively large or extensive mid third tibial wound with good success. (10)

The distally based (or reverse) sural artery fasciocutaneous flap has gained much of the attention recently. (11) The flap is found to be reliable and versatile and can be elevated quickly to cover a wound in the distal third of the leg. At the present time, the proper management of soft tissue coverage for a less extensive tibial wound in the junction of the middle and distal thirds of the leg has rarely been discussed in literature and therefore, the optimal reconstruction for this unique clinical problem has not been determined. (12)

A soleus muscle flap can be considered as an option. Based on the previous studies, the authors used the soleus muscle flap for reconstruction of an open tibial wound in the junction of the middle thirds of the leg and reverse sural with good success. For reconstruction of a wound in the distal third of the leg, a microvascular free tissue transfer has been considered as a standard surgical procedure of the choice because in general, there are no reliable local options available for reconstruction. (4, 7)


In general, a microvascular free tissue transfer should still be considered for a larger soft-tissue wound in the distal third of the leg, or for a less extensive wound, when either the soleus muscle or those minor pedicles from the posterior tibial vessels are traumatized. (12, 13) The distally based (or reverse) sural artery fasciocutaneous flap has gained much of the attention recently. (10) The flap is found to be reliable and versatile and can be elevated quickly to cover a wound in the distal third of the leg. It can even reach the foot and ankle area to cover both medial and lateral malleoli wounds and also a heel wound. (10) The reverse soleus flap had an excellent outcome with short surgical duration, easy implementation, excellent resolution, and low morbidity of the donor area.

Occasionally, the venous congestion can be a problem for the flap and thus either surgical delay or supercharge of the flap has been recommended. (14) Some other reverse adipose-fascial flaps can also be used to cover a wound in the distal leg. (15) Other advantages of this procedure are in risk factor existence. These risk factors could impair successful defect coverage such as diabetes mellitus, and peripheral vascular disease. The aim of this method is to present an alternative option for reconstruction of large defect with combination of fasciocutaneous and muscular flap instead of free flap. Although, the entire soleus muscle used as a flap has a limited arc of rotation. (15)

Therefore, it may not reach a relatively distal soft tissue defect over the tibia. Limitation of plantar flexion of the foot may be another disadvantage, when the whole muscle is sacrificed. More importantly, the distal portion of the soleus muscle after elevation has traditionally been considered unreliable for coverage of a relatively distal tibial wound because of the muscle's circulation pattern, but for proximal defect it is versatile flap. (15)

The reconstructive outcomes are usually quite good and a low cost-effective approach. Also it can be performed by most reconstructive surgeons in selected patients.

Figures: figures 1-3 showing a large sized tissue defect which was initially managed by orthopaedic trauma team and which was later repaired by using reverse sural and soleus muscle flaps and a good result was obtained.



#### REFERENCES:

1. Heppert V, Becker S, Winkler H, et al. Myocutaneous versus fasciocutaneous free flap in the treatment of lower leg osteitis. *Eur J Orthop Surg Traumatol.* 1995;5:27-31
2. Reddy V, Stevenson TR. Lower extremity reconstruction. *Plast Reconstr Surg.* 2008;121:1-7.
3. Kneser U, Bach AD, Polykandriotis E, et al. Delayed reverse sural flap for staged reconstruction of the foot and lower leg. *Plast Reconstr Surg.* 2005;116:1910-7
4. Yang C, Geng S, Fu C, et al. A minimally invasive modified reverse sural adipofascial flap for treating posttraumatic distal tibial and calcaneal osteomyelitis. *Int J Low Extrem Wounds.* 2013;12:279-85
5. Maghari A, Forootan KS, Fathi M, Manafi A. Free transfer of expanded parascapular, latissimus dorsi, and expander "capsule" flap for coverage of large lower-extremity soft-tissue defect. *Plast Reconstr Surg.* 2000;106:402-5.
6. Heller L, Levin LS. Lower extremity microsurgical reconstruction. *Plast Reconstr Surg.* 2001;108:1029-41.
7. Armstrong MB, Villalobos RE, Leppink DM. Free-tissue transfer for lower-extremity reconstruction in the immunosuppressed diabetic transplant recipient. *J Reconstr Microsurg.* 1997;13:1-5.
8. Pu LLQ. The medial hemisoleus muscle flap: a reliable flap for soft-tissue reconstruction of the middle-third tibial wound. *Int Surg.* 2006;91:194-200.
9. Thornton BP, Pu LLQ. Reconstruction of an extensive tibial soft tissue defect with multiple local muscle flaps for limb salvage when free tissue transfer was not an option. *Euro J Plast Surg.* 2004;27:217-21.
10. Hollier L, Sharma S, Babigumira E, Klebuc M. Versatility of the sural fasciocutaneous flap in the coverage of lower extremity wounds. *Plast Reconstr Surg.* 2002;110:1673-9.
11. Barbour JR, Schweppe M, O SJ. Lower-extremity burn reconstruction in the child. *J Craniofac Surg.* 2008;19:976-88.
12. Pu LLQ. Successful soft-tissue coverage of a tibial wound in the distal third of the leg with a medial hemisoleus muscle flap. *Plast Reconstr Surg.* 2005;115:245-51.
13. Pu LLQ. Further experience with the medial hemisoleus muscle flap for successful soft-tissue coverage of a tibial wound in the distal third of the leg. *Plast Reconstr Surg.* 2008;60:594-97.
14. Follmar KE, Baccarani A, Baumeister SP, Levin LS, Erdmann D. The distally based sural flap. *Plast Reconstr Surg.* 2007;119:138e-148e.
15. Gumener R, Zbrodowski A, Montandon D. The reversed fasciocutaneous flap in the leg. *Plast Reconstr Surg.* 1991;88:1034-43.

