



COMPARATIVE STUDY OF SERUM ELECTROLYTES IN SENILE CATARACT PATIENTS AND IN NORMAL INDIVIDUALS

Dr. S. Aruna

M.D(Bio), Assistant professor ,Dept of Biochemistry, Govt Karur Medical college, Karur.

Dr. N. Santhi*

M.D(Bio),Assistant professor , Dept of Biochemistry, K.A.P.V Govt Medical college, Trichy. *Corresponding Author

ABSTRACT

Background: Cataract refers to opacification of crystalline lens in the human eye. Age is the strongest known non-modifiable risk factor for cataract formation. One of the proposed risk factors for cataract formation is rise in serum sodium level. Though ageing cannot be prevented, physiological changes that occur in electrolytes can be modified.

Aim: To compare serum electrolyte levels in senile cataract patients with those without cataract.

Materials and Methods: Serum electrolytes were estimated in Fifty senile cataract patients and age matched Fifty healthy people without cataract by using an electrolyte analyzer.

Results: We noted significant ($p < 0.0001$) rise in sodium levels in cases (143.41 ± 1.69 meq/l) compared to controls (138.7 ± 1.66 meq/l) and chloride levels ($p < 0.0001$) in cases (103.47 ± 3.48 meq/l) compared to controls (99.38 ± 2.65 meq/l). Serum potassium levels were significantly low in cases (4.07 ± 0.46) compared to controls (4.31 ± 0.45) ($p = 0.0126$).

Conclusion: Serum sodium and chloride levels are elevated in senile cataract patients. So, dietary salt restriction may delay the cataract formation

KEYWORDS : Senile Cataract ,Serum Sodium, Serum Chloride.

INTRODUCTION

There are about 50 million blind people in the world, a third of them being due to cataract. (Thylefors, Négrel, Pararajasegaram, & Dadzie, 1995) Its prevalence in developing countries is much more than the developed ones. In India alone, cataract accounts for 80% of treatable blindness. One of the most common types of cataract is the senile cataract which occurs as a consequence of the aging process. Senile cataract usually occurs after the age of 45 years. Approximately 75 percent of population above the age of 75 years suffers from cataract.

Many risk factors such as age, sex, radiation, genetics, metabolic disorders, protein aggregates, oxidative stress, post translational protein changes, phase separation are proposed for cataract formation, though the exact pathogenesis is not yet known. Many risk factors such as age, sex, radiation, genetics, metabolic disorders, protein aggregates, oxidative stress, post translational protein changes, phase separation are proposed for cataract formation, though the exact pathogenesis is not yet known. (Adiga, Harris, Ezhilvathani, & Basu, 2014)

Aqueous humour is the main source of nourishment for the lens. This thin fluid, is produced from the serum. Therefore, serum electrolytes concentration in turn regulates lens metabolism. (Fawad, Maqsood, & Abbas) It has been proved that, in aqueous humour, potassium is replaced with excess sodium and hence there is an alteration in their ratio in cataract patients. (Spector, 1995) This was attributed to the changes that occur in the serum cations. As this fluid is derived from the plasma, derangement in serum electrolytes appears to be one of the risk factor for cataractogenesis. (Jain, Vyas, & Sogani)

Some studies have shown significant difference in serum electrolytes concentration in senile cataract patients when compared to those without cataract (Mirsamadi, Nourmohamadi, & Imamian, 2004). The purpose of the study is to estimate serum electrolytes in senile cataract patients as compared to the healthy age and gender matched people without cataract. To the best of our knowledge, there are a few studies in this field and the results are inconclusive. As cataract is one among the treatable causes of blindness, it is justifiable to make an attempt to identify a probable risk factor for the cataractogenesis. (Jadav Prashantkumar et al.)

AIMS AND OBJECTIVES:

1. To estimate the serum electrolyte levels in patients with senile

cataract

2. To compare the electrolyte levels with the age and gender matched healthy controls without cataract
3. To study the association of serum electrolytes with risk of cataract formation.

MATERIAL AND METHODS:

It is a cross sectional observational study conducted in the department of Biochemistry in collaboration with the department of Ophthalmology, Thanjavur medical college, Thanjavur. The duration of the study was one month. Senile cataract patients who were scheduled to undergo cataract surgery in the Department of Ophthalmology were taken as cases. The patients with complicated cataract, those with renal disorders, liver cirrhosis, hypertension, diabetes mellitus, thyroid disorders, infections, those who have sustained trauma, those on medications which alter electrolytes like steroids, smokers, and alcoholics were excluded from our study. Exclusion was done based on the relevant history, physical examination and minimum investigations required in making the diagnosis.

Based on this, two groups were made:

Group I: Senile cataract patients, who were scheduled to undergo cataract surgery.

Group II: Age and gender matched normal healthy individuals without cataract, who were attending the government medical college hospital for routine check up and attenders who accompany the patients.

Each group included 50 subjects. Approval from the institutional ethics committee was obtained. Informed written consent of the participants was taken. A pre structured proforma was used to collect the baseline data and the cases were examined by an ophthalmologist in the hospital

METHODOLOGY:

BIOCHEMICAL ESTIMATION: Two ml of blood sample is collected with aseptic precautions for the estimation of serum electrolytes. Electrolytes are estimated by using the electrolyte analyzer which works on the principle of ion selective electrodes.

STATISTICAL ANALY

Data analysis was done by Graph Pad online software. Means of electrolytes were compared between two groups by unpaired t-test. Interpretation was done according to p-value.

RESULTS

In this study the case group consisted of 50 patients suffering from age-related cataract and the control group consisted of 50 healthy subjects who were attending the OPD for regular check up or patient's bystanders. Reference range for sodium was taken as 136-145 mEq/l, for potassium is 3.5 – 5.2 mEq/l and that for chloride 96-106 mEq/l. Senile cataract patients had an extremely significant elevation in sodium levels 143.41 ± 1.69 mEq/l ($p < 0.0001$) as compared to controls, 138.7 ± 1.66 mEq/l.) and chloride levels ($p < 0.0001$) in cases (103.47 ± 3.48 meq/l) as compared to controls (99.38 ± 2.65 meq/l). Serum potassium levels were significantly low in cases (4.07 ± 0.46) as compared to controls (4.31 ± 0.45).

Table-1: Comparison of serum electrolytes in senile cataract patients versus controls

SERUM LEVELS(meq/L)	CASES(n=50)	CONTROLS(n=50)	P VALUE
SODIUM	143.41 ± 1.69	138.7 ± 1.66	$< 0.0001^*$
POTASSIUM	4.07 ± 0.46	4.31 ± 0.45	$= 0.0126^{**}$
CHLORIDE	103.47 ± 3.48	99.38 ± 2.65	< 0.0001

* - EXTREMELY SIGNIFICANT STATISTICALLY (P VALUE < 0.0001)

**-. SIGNIFICANT STATISTICALLY (P VALUE < 0.05)

Odds's ratio is calculated to find the risk associated with elevated sodium (> 145 meq/l), chloride (> 105 meq/l), and cataract formation.

Chances of senile cataract formation is 5 times greater in people with serum sodium > 145 meq/l. People with high serum chloride levels are 1.7 times at a higher risk of developing senile cataract. Potassium levels did not show any significant correlation with cataract formation.

DISCUSSION:

We have found a extremely significant elevation in serum sodium and chloride levels in senile cataract patients as compared to healthy controls and a fall in potassium levels in cases which is also statistically significant. A study among Ghanians showed a very strong association between the level of exposure to serum sodium and the probability of developing the senile cataract. Comparing the serum electrolyte levels, there was a statistically significant difference between the mean serum Na⁺ levels and serum Cl⁻ levels in senile cataract patients and normal individuals. The mean serum K⁺ levels of senile cataract patients was statistically significant as compared to controls. The study concluded that exposure to sodium in the absence of other biochemical risk factors remained the most significant risk factor for the development of senile cataract. A study in Iranian population by Mansour Mirsamadi et al, showed that the mean of serum Na⁺ of cataract patients were in the normal range but were in the upper limit of this range and in comparison with control group, the serum Na⁺ of the patients was significantly elevated.(Khan et al., 2014)But potassium did not show any significant difference. Elevation of serum electrolytes in senile cataract patients can be explained as follows:

Lens has a high content of potassium and low content of sodium. (Harris & Gehrsitz, 1951)These two cations are in balance with each other due to Na⁺- K⁺- ATPase pump and lens capsular permeability. Normally lens has high level of K⁺ (114-130 mmol/L) and low Na⁺ (14-26mmol/L). These two cations are in balance with each other due to action of Na⁺ K⁺ATPase pump, which in turn maintains permeability of lens membrane.(Anbarasi, Vani, Balakrishna, Devi, & toxicology, 2005) With ageing there is an increase in membrane permeability of the lens cells due to reduced activity of Na⁺ K⁺ ATPase pump, which leads to an increase in internal Na⁺. (Rhodes & Sanderson, 2009)Higher levels of extracellular Na⁺ might make it more difficult for Na⁺ K⁺ ATPase pump to maintain the low levels of intracellular Na⁺ required for lens transparency. (Mathias, Rae, & Baldo, 1997)Variation of electrolytes in the serum in turn alters cation concentration of aqueous humors, which ultimately affect

lens metabolism leading to cataract formation. Alteration in cation concentration of aqueous humor which is attributed to alterations in serum cation concentration, can be an important risk factor for cataract formation. (Becker, 1959)Previous studies notify the significant difference between serum sodium of those suffering from age-related cataract versus those without cataract. But serum potassium did not show significant variation. Diets with high sodium contents could be a risk factor for senile cataract formation. As it seems, a high level of serum sodium in turn contributes to cataract formation.

The permeability of lens is high for chloride ions. The Na⁺-K⁺ 2Cl⁻ co transporter has been identified defect affecting chloride handling by the lens. The ability of the lens to maintain its hydrated state is lost; imbalance in osmotic equilibrium may result in cataract.

CONCLUSION:

We can conclude from our study that,

- Serum sodium and chloride levels are significantly increased in senile cataract patients and can be used to determine the risk involved in senile cataract formation.
- Simple measures like dietary restriction of salt may prolong cataractogenesis, which can be a preventive measure useful to the community.
- This study may be useful in patient care as follows:Cataract surgery is the commonest surgery in the field of ophthalmology. Much cost spent on cataract surgery may be minimized.
- Prevents disabilities and helps to improve quality of life, useful for the health care system.

REFERENCES

1. Adiga, U. S., Harris, A., Ezhilvathani, T., & Basu, S. J. A. J. M. S. (2014). Serum electrolytes in senile cataract patients. 7(2), 164-168.
2. Anbarasi, K., Vani, G., Balakrishna, K., Devi, C. S. J. J. o. b., & toxicology, m. (2005). Effect of bacside A on membrane-bound ATPases in the brain of rats exposed to cigarette smoke. 19(1), 59-65.
3. Becker, B. J. A. j. o. o. (1959). Carbonic anhydrase and the formation of aqueous humor: The Friedenwald Memorial Lecture. 47(1), 342-361.
4. Fawad, H., Maqsood, M., & Abbas, A. International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR).
5. Harris, J. E., & Gehrsitz, L. B. J. A. j. o. o. (1951). Significance of changes in potassium and sodium content of the lens. 34(5), 131-138.
6. Jadav Prashantkumar, M., Sharma Hariom, M., Maniar Megha, A., Chaudhary Nitinkumar, G., Maheshwari Amit, V., & Javia, H. Relationship between altered level of serum electrolytes and risk of senile cortical cataract-A Case Control Study.
7. Jain, P. P., Vyas, A., & Sogani, S. ESTIMATION OF SERUM ELECTROLYTES (NA AND K) AND SERUM CALCIUM LEVELS IN SENILE CATARACT PATIENTS AND ITS COMPARISON WITH NON CATARACT PATIENTS.
8. Khan, A. A., Rizvi, S. W. A., Amitava, A. K., Moin, S., Siddiqui, Z., & Yusuf, F. J. S. J. o. O. (2014). Serum Na⁺ and K⁺ as risk factors in age-related cataract: An Indian perspective. 6(1), 10.
9. Mathias, R., Rae, J., & Baldo, G. J. P. r. (1997). Physiological properties of the normal lens. 77(1), 21-50.
10. Mirsamadi, M., Nourmohammadi, I., & Imamian, M. J. I. j. o. m. s. (2004). Comparative study of serum Na⁺ and K⁺ levels in senile cataract patients and normal individuals. 1(3), 165.
11. Rhodes, J. D., & Sanderson, J. J. E. e. r. (2009). The mechanisms of calcium homeostasis and signalling in the lens. 88(2), 226-234.
12. Specter, A. J. T. F. J. (1995). Oxidative stress-induced cataract: mechanism of action. 9(12), 1173-1182.
13. Thylefors, B., Négrel, A. D., Pararajasegaram, R., & Dadzie, K. J. B. o. t. w. h. o. (1995). Global data on blindness. 73(1), 115.