



A COMPARITIVE STUDY ON INDUCTION OF LABOUR VERSUS EXPECTANT MANAGEMENT IN TERM PREMATURE RUPTURE OF MEMBRANES IN TERTIARY CARE HOSPITAL , VISAKHAPATNAM .

Dr Kavitha Gayak

Ms Obg , associate Professor , Department Of Obstetrics And Gynaecology, Gitam Institute Of Medical Sciences And Research , Rushikonda , Visakhapatnam ,ap.

Dr Nirupama Palakodeti*

Ms Obg , Assistant Professor, Department Of Obstetrics And Gynaecology, Gitam Institute Of Medical Sciences And Research, Rushikonda, Visakhapatnam ,ap. * Corresponding Author

ABSTRACT

Premature rupture of membranes (PROM), also called Prelabour Rupture Of Membranes, is classically defined as rupture of membranes before labour and accounts for 0.8-0.9% of all pregnancies at term. this study was conducted over a period of 14 months (from March 2018 to May 2019) in a tertiary care medical college . A total of 100 term PROM patients were recruited in our study – and divided into two groups randomly--50 (Group A) patients were managed conservatively and 50 (Group- B) patients underwent induction of labour. Both these patient groups were studied to compare the fetomaternal outcome. Group-A (conservative management group) patients were observed to await the spontaneous onset of labour pains for at least 24 hours. Patients in group B were induced with either - PGE1 tab (misoprostol) 25 µgm 4 hourly orally or iv oxytocin infusion. the PROM-delivery interval was < 12 hours in 72 % of induced groups (Group B) and 10% (5) in group- A (conservative or expectant group). LSCS rate was 10 % in group-A (expectant group) & 15 % in group –B (induced group). Sepsis rate, maternal and fetal, hospital stay, NICU admission & duration of NICU stay were notably higher in group – A (expectant management group). therefore, from our study we concluded that immediate induction of labour in term PROM cases shortens the PROM- delivery interval, hospital stay, NICU stay and reduction in both maternal & neonatal sepsis.

KEYWORDS : Term PROM, induction of labour, fetomaternal outcome, neonatal sepsis, NICU, expectant management

INTRODUCTION:

Premature or Prelabour Rupture Of Membranes is classically defined as the rupture of integrity of the fetal membranes before the onset of labour and resulting in leakage of amniotic fluid. (2,3,4,).Pre-labour rupture of membranes without spontaneous uterine contractions complicate approximately 10 % of all pregnancies, out of which 80 % occurs at term (5).the management of PROM at term remains a matter of great debate till date. While induction of labour has resulted in decreased incidence of maternal & fetal sepsis, but it is also associated with a higher incidence of caesarean section rate due to fetal distress and uterine hyperstimulation. Kappy et al in their study, reported a higher incidence of operative deliveries in term pregnancies with ruptured membranes managed with labour stimulations compared with those who were managed expectantly.(6). Approximately 65-75 % of term PROM patients are followed by spontaneous onset of labour within 24 hours (7). Prostaglandins- PGE2& PGE1 have been used for cervical ripening and myometrial stimulation in unfavorable cervixes with low Bishop's pre- induction score. It is seen that in patients who had expectant management, with prolonged hospitalization without active intervention with uncertain fetal and neonatal prognosis, many a times leads to maternal psychological sequelae.(8)

MATERIALS & METHODS:

Study type: Hospital based prospective observational study.

Study duration: Fourteen months

Study period: March 2018 to April 2019

Study subjects: Women with term PROM as per inclusion & exclusion criteria

Study place – department of obstetrics and gynaecology , githam institute of medical sciences and research , rushikonda, visakhapatnam, AP, India.

INCLUSION CRITERIA:-

1. Gestational age 37-40 weeks
2. Singleton pregnancy

3. Age 20-30 years
4. Adequate pelvis
5. Vertex presentation

EXCLUSION CRITERIA:-

1. Age < 20 years , > 30 years
2. Gestational age < 37 weeks , > 40 weeks
3. Multiple pregnancy
4. Chorioamnionitis
5. Medical disorders
6. Obstetric complications
7. Grand multipara
8. Previous LSCS
9. Meconium stained liquor
1. Cephalopelvic disproportion
2. Malpresentation
3. Intrauterine fetal death

All the patients who fulfilled the inclusion and exclusion criteria were subjected to a detailed history taking followed by thorough clinical examination including general and systemic examination to rule out any exclusion criteria. A thorough obstetrical examination including per speculum examination and Bishop's scoring was conducted. An obstetric ultrasonography and CTG were performed. All the patients were given IV antibiotics of 1 gram ceftriaxone, 100 ml metronidazole and gentamycin 80 mg after negative skin test till delivery. Informed written consent was taken and patients were randomly allocated to either Group A (expectant or conservative management group) or Group B (active management group with induction of labour). Patients in Group-B (active management group) were induced according to Bishop's pre- induction score . PGE 1 tablet (misoprostol) 25µm orally was given if Bishop's score was less than 5 followed by repeat dose after 4 hours or iv oxytocin at the rate of 5 units in 500ml RL or Normal Saline (for primigravida) or @ 1- 2 units for multigravida) and the drip was titrated in escalating doses till optimal response was observed as evidenced by onset of effective uterine contractions. All these patients had vitals charting, pulse, temperature, blood

Submitted : 27th July,2019

Accepted : 15th August,2019

Publication : 15th October, 2019

pressure along with progress of labour- fetal heart rate, uterine contractions and descent of head. Patients in Group-A (expectant management group) were kept for observation with sterile vulval pad for 24 hours to await spontaneous onset of labour pains. Vitals charting (pulse rate, temperature, blood pressure) , uterine contractions fetal heart rate, colour of liquor were monitored carefully . Unnecessary vaginal examinations were avoided and P/V exam done only if the uterine contractions were good, to gauge the progress of labour. In this group, most of the patients went into spontaneous labor in 24 hours. Induction of labour was done after 24 hours if there were no contraction. Emergency LSCS was done for indications like fetal distress, non progression of labour, cord prolapse, failed induction, and chorioamnionitis. All labour and delivery events were noted example vaginal delivery, spontaneous or instrumental-forceps/ventouse, LSCS etc. e fetal outcomes were noted namely APGAR score, birth weight, neonatal sepsis, NICU admissions and care required. All the patients were followed up in puerperium to assess maternal pyrexia, sepsis and neonatal sepsis.

RESULTS :

the results and observations of the present study are as follows : Most of the patients were primigravidae, had low education levels, hailed from rural background and from low socio-economic stratum, young aged and mostly unbooked.

Table 1 : Distribution of patients according to parity

	Group –A (Expectant Management)		Group- B(Induced)	
	Number	%	Number	%
Primigravida	27	54.0	22	44.0
Multigravida	23	46.0	28	56.0
Total	50	100.0	50	100.0

Table 2 : Mode of delivery

	Group – A (Expectant)		Group-B (Induced)	
	Number	%	Number	%
Vaginal delivery	45	90.0	41	82.0
Spontan eous delivery	38	56.0	28	56.0
Ventouse delivery	05	10.0	8	16.0
Forceps delivery	02	4.0	5	10.0
LSCS	5	10.0	9	18.0
Total	50	100.0	50	100.0

Table 3: PROM- delivery interval (PDI)

Time	Group-A (Induced)		Group-B (Induced)	
	Number	%	Number	%
< 12 hours	5	10.0	36	72.0
12-24 hours	35	70.0	11	22.0
>24 hours	10	20.0	3	6.0
Total	50	100.0	50	100.0

Table 4 : Maternal outcome

	Group- A (Expectant)		Group-B (Induced)	
	Number	%	Number	%
Fever	3	6.0	1	2.0
PPH	5	10.0	6	12.0
Sepsis	5	10.0	3	6.0
Nil	37	74.0	40	80.0
Total	50	100.0	50	100.0

Table 5 : Neonatal outcome

	Group- A(Expectant)		Group-B(Induced)	
	Number	%	Number	%
Birth asphyxia	14	28.0	8	16.0
Mild APGAR <7	10	20.0	6	12.0
Severe APGAR <5	4	8.0	2	4.0
Sepsis	2	4.0	0	0.0
Stillbirth /early neonatal	0	0.0	0	0.0

Nil	36	72.0	42	84.0
Total	50	100.0	50	100.0

DISCUSSION

In our study, the women in both the Group A (expectant management group) and Group-B (Induced group) were comparable with respect to mean maternal age, gestational age , parity, educational status, socio-economic background, urban-rural distribution. Since their socio-demographic profile was similar, therefore, any difference in outcome in these two groups was primarily due to different management protocols and not due to demographic differences. In our study we observed that vaginal delivery occurred in 90% of Group A and 82.0% in Group-B (Induced) patients. A study by Shanti K et al stated LSCS rate as 5.7% in the expected group as compared to 12% in active group (9). In another study by Suneela K at al , there were 88.3% vaginal delivery in expectant group and 85.0% in active management group i.e , 11.7% LSCS rate in expectant group and 15% in induced group.(1)In our study, we observed intrapartum pyrexia in 65 of expectant group (Group-A) versus 2 % in Group –B (Induced group). Suneela K et al reported 3.3% pyrexia in induced group and 5 % in expectant group (1). Sumaira Yasmeen et al (Peshawar) reported 2.4 % in induced group & 16 % fever in expectant group (10). In our study, PPH occurred in 10% Group-A & 12% Group-B, whereas Suneela K et al reported 6.7% PPH in expectant group & 10% in induced group(1).this result may be because of the fact that- induction of labour has a higher incidence of PPH". (1, 2).In our study, 28.0% babies in Group-A suffered from birth asphyxia (severe-8% and mild-20%) as compared to 16.0% in Group- B (mild- 12% & severe 4%).Neonatal sepsis was observed in 4 % of Group A and in none (0 %) in Group B .this may be attributed to the fact that there was a prolonged - delivery interval in Group A (expectant group).there were no stillbirths or early neonatal deaths in either groups. In her study, Suneela K et al reported severe birth asphyxia requiring ventilation in 6.6 % in the expectant group and neonatal sepsis was reported as 11.7% in the expectant group.(1) Active management in cases of PROM at term has shown many benefits in terms of reducing the latent period and better maternal satisfaction (8).Induction of labour in such cases not only reduces the PROM -delivery interval but also reduces neonatal and maternal sepsis without much significant rise in LSCS rate. It is clear that nearly 70% of term PROM patients go into spontaneous labour without induction within 24 hours. e method of induction whether PGE1 (misoprostol) or oxytocin depends on Bishop's pre-induction score or cervical findings.

CONCLUSION -

In our study we concluded that with induction of labour in patients with PROM at term, the PROM delivery interval was reduced along with significantly better maternal satisfaction and feto maternal outcome. the expectant group who underwent conservative management had higher maternal and fetal morbidity, sepsis, longer hospital and NICU stay causing anxiety and distress to both patients and clinicians. Therefore, in all patients presenting with premature rupture of membranes at term should be actively managed with induction of labour after assessing the cervical condition according to the Bishop's pre-Induction score so as to reduce the incidence of maternal and fetal sepsis and morbidity.

REFERENCES

1. Suneela K et al, Sch.J.App.Med.Sci, April 2016, 4 (40): 1424- 1427.
2. Gary Cunningham F et al ; Normal Labour Chapter 22, William Obst- 24thEd, NY, Mc Gram Hill -2014; 886 -933; 948.
3. Larranga - Azcarate C at al ; Dinoprostone Vaginal slow release system compared to expectant management in the active treatment of PROM at term; impact on maternal and fetal outcome ; Actaobstetria, 2008 ; 87 (2) : 195-200.
4. Shah K et al- PROM at term: early induction versus expectant management; e J. ob. Gynec. India 2012; 62 (2): 172-175.
5. Duff P: PROM in term patients: Induction of labour versus expectant management; clinObstetGynecol1998;41: 883 - 891.
6. Kappa KA et al; PROM: conservative approach Am. J. obs. Gyn1979;134: 655.

7. Hoffmann RA et al: Oral misoprostol versus placebo in the management of PROM at term; Int. J of GynecObstet. 2001;72: 215-221.
8. Term Pre labour rupture of membranes: e Royal Australian & NZ College of Obs&Gyn(C-obs 36) 1-9.
9. Shanti K et al: Comparative study of active vs expectant management and maternal and neonatal outcome in PROM: IOSRJDMS; 2015; 14(4):34-9.
10. Sumaira Yasmeen et al: Active versus conservative management of PROM at term. J Postgrad Med Inst 2013; 27(1): 63-68.