



## THE CORELATION OF DURATION OF HOSPITALISATION AND ONSET OF VENTILATOR ASSOCIATED PNEUMONIA

**Dr Kiran Tandia**

Assistant professor department of medicine Bundelkhand medical college sagar

**Dr J. L wadhvani\***

Associate Professor Department Of Medicine Gmc, Bhopal.  
\*Corresponding Author

### ABSTRACT

Introduction Ventilator-associated pneumonia (VAP) is defined as pneumonia that occurs 48-72 hours or thereafter following endotracheal intubation, characterized by the presence of a new or progressive infiltrate, signs of systemic infection (fever, altered white blood cell count), changes in sputum characteristics, and detection of a causative agent. VAP contributes to approximately half of all cases of hospital-acquired pneumonia. VAP is estimated to occur in 9-27 % of all mechanically ventilated patients, with the highest risk being early in the course of hospitalization. Aims & objectives-to study the association between duration of stay and onset of VAP. Results and conclusion-Incidence is directly proportional to duration of mechanical ventilation. It is a strong risk factor for development of VAP. Therefore, duration of ventilation has to be reduced.

**KEYWORDS :** Ventilator Associated Pneumonia, Mechanical Ventilator, Duration Of Stay, Intubation, Nosocomial Infection.

### INTRODUCTION -

Infections are the most important and leading cause of mortality and morbidity among the patients admitted in ICU. Nosocomial infections is a critical issue among intubated patients which is responsible for significant morbidity and mortality of these patients (1). One of the most important types of this infection is pneumonia which commonly occurs in relation to the endotracheal intubation and mechanical ventilation named ventilation associated pneumonia (VAP) (2-4). Patients with mechanical ventilation have an increased risk for respiratory tract infection because the tube which has been inserted into the trachea reduces the clearance of bacteria and increases the leakage of secretion around the cuff of the tube and disables the ciliary tract by damaging it (10). Clinical diagnosis: Ventilator associated pneumonia usually suspected when the individual develops a new and persistent (>48 hour) or progressive radiographic infiltrate plus two of the following; temperature >38°C or <36°C, blood leukocyte count of > 10,000 cells/ml or < 5,000 cells/ml, purulent tracheal secretions and gas exchange degradation.

Ventilator-associated pneumonia (VAP) is defined as pneumonia that occurs 48-72 hours or thereafter following endotracheal intubation, characterized by the presence of a new or progressive infiltrate, signs of systemic infection (fever, altered white blood cell count), changes in sputum characteristics, and detection of a causative agent. VAP is estimated to occur in 9-27 % of all mechanically ventilated patients, with the highest risk being early in the course of hospitalization. VAP rates range from 1.2 to 8.5 per 1,000 ventilator days and are reliant on the definition used for diagnosis [6]. Risk for VAP is greatest during the first 5 days of mechanical ventilation (3 %) with the mean duration between intubation and development of VAP being 3.3 days. This risk declines to 2 %/day between days 5 to 10 of ventilation, and 1 %/day thereafter. Intubated patients with nosocomial pneumonia have been associated with a longer ICU stay.

### MATERIAL AND METHODS

The study was conducted in Department of Medicine emergency medical ward of a Hamidia hospital Bhopal. Over a period of 1 year, extending from July 2013 to December 2014, it was an observational cross-sectional study. Patients intubated for >48 hours in emergency medical ward due to insufficient self-ventilation were included in the study. Patients who already had any respiratory tract infection or any other disease related to the respiratory system and patients who are intubated for <48 hours were excluded from the study.

A questionnaire was prepared and each patient selected to be included in the study was screened and monitored according to the questionnaire. Age, sex, date of admission to ICU, date of initiating mechanical ventilation and mode of access to the patients' airway, i.e. orotracheal or tracheostomy, were recorded. Indication of mechanical ventilation was noted. In each patient, ventilator mode and settings were recorded and any change in setting was recorded daily. Patients' vitals, general and physical examination, oxygen saturation and position of the patients were monitored regularly. All necessary measures were taken for prevention of hospital-acquired infections.

Sputum from the patients were collected from the tip of the suction catheter and transported to the laboratory in a sterile tube under all aseptic precautions. Patients were monitored from the date of inclusion in the study to the final outcome in the ICU. VAP was diagnosed on clinical grounds based on the modified CPIS system originally developed by Pugin and others, giving 0-2 points each for fever, leukocyte count, oxygenation status, quantity and purulence of tracheal secretions, type of radiographic abnormality and result of sputum culture and Gram stain. The VAP group was classified into two groups, early-onset type (within 48-96 h) and late-onset type (>96 h). Clinical Pulmonary Infection Score (CPIS)

Parameter Score

Temperature (°C)

>36.5 and <38.40  
>38.5 and <38.9 1  
>39.0 or <36.52

White Blood Cell Count

>4,000 and <11,000  
<4,000 or >11,000  
<4,000 or >11,000 & band forms >50% 2

Tracheal secretions

None or scant 0  
Non-purulent 1  
Purulent 2

Pao<sub>2</sub>/Fio<sub>2</sub>

>240, ARDS or pulmonary contusion 0  
<240 and no ARDS 2

Chest Radiograph

No infiltrate 0  
Diffuse or patchy infiltrate 1

Localised infiltrate2

Data were analyzed by Microsoft excel and Statistical Package for Social Sciences and the values are reported as number (%). The level of significance was set at P<0.05.

**RESULTS**

**Table-1.** Distribution Of Cases According To Duration Of Stay

Durationofstay	Culturepositive	Negative	Total
<4days	9	52	61
	14.8%	85.2	100%
>4days	16	23	39
	41%	59%	100%
Total	25	75	100

P=0.003 [significant, Longer stay more vap]

**Table-2** Duration Of Stay According To Number Of Cases

Durationofstay	Numberofcases	%ofcases
3	61	61
4	29	29
5	7	7
6	1	1
8	1	1
9	1	1

**Table-3**Distribution Of Cases According To Type Of Tracheal Secretions

Trachealsecretion type	Culture positive	Negative	Total
Purulent	9	0	9
	100%	0%	100%
Scanty	16	75	91
	17.6%	82.4%	100%
Total	25	75	100

P < 0.00([ significant, all purulent tracheal secretions had vap)

**Table-4** Age And Sex Distribution Of Cases

Ageinyears	Female	Male	Total
<20	3	6	9
	33.3%	66.7%	100%
20-30	21	20	41
	51.2%	48.8%	100%
31-40	4	10	14
	28.6%	71.4%	100%
41-70	0	12	12
	0%	100%	100%
70andabove	5	19	24
	20.8%	79.2%	100%
Total	33	67	100
	33%	67%	100%

**DISCUSSION-**

It is analysed in our study that those requiring prolonged ventilator support had a significantly higher incidence of VAP (P-value, 0.003) of the 25 patients who developed VAP, 9 patients developed early-onset (14.8%) VAP and 16 patients developed the late-onset type (41.0%).In our study it is also found that purulent secretions associated with positive sputum culture (p value< 0.001).[6]

In the study of our set up, males predominated (67%). The study shows the incidence of VAP in males was 16%, and in females is 9%.

Although the incidence of VAP is high in males but it is statistically not significant (P=0.713).The study also shows that there is no significant difference in age as risk factor(P=0.276). The study also shows that patients with

longer duration of hospital stay, more chances of development of VAP (p value,0.03).[49] It was proved in our study that duration of mechanical ventilation is an important risk factor for VAP, which is similar to other studies[7] where the mean duration of ventilation was around 10 days and the incidence of VAP was found to be 9.3%.(7) Early-onset VAP in our study was found to be 14.8% while in various study it was found to be around 40%.[10] The low incidence in our study may be due to antibiotic use before admission to the ICU. Studies[9]

**CONCLUSION-**

Incidence is directly proportional to duration of mechanical ventilation iis a strong risk factor for development of VAP. Therefore, duration of ventilation has to be reduced .

**REFERENCES**

1. Cunnion ,K.M.,D.J. weber, W.E. Broadhead, L.C.Hanson , C.F. Pieper and W.A. Rutala.1996 Risk factors for nosocomial pneumonia: comparing adult critical populations . Amj. Respir. Crit. Care Med. 153 : 158 -162.
2. Hiffken G, Niederman MS. Nosocomial pneumonia: The importance of a de-escalating strategy for antibiotic treatment of pneumonia in ICU.
3. 58. Applengren P, Hellstrom I WeitzbergE, Sodurlund V,Bindslev L, Ransjo U. Risk factors for nosocomial intensive care infection: A long term prospective analysis.
4. Tejada Artigas A, Bello Dronda S, Chacon Valles E, Munoz Marco J, Villuendas Uson MC, Figueras Petal. Risk factors for nosocomial pneumonia in critically ill trauma patients.
5. Hina gadani, Arun Vyas,Akhya kumar Kar, A study of ventilator associated pneumonia: incidence,outcome, risk factors and measures to be taken for prevention.
6. Pugin J, Auckenthaler R, Mili N, Janssens JP, Lew PD, Suter PM: Diagnosis of ventilator associated pneumonia by bacteriologic analysis of bronchoscopic and non bronchoscopic "blind" bronchoalveolar lavage fluid.
7. Rello J, Sa-Borges M, Correa H, Leal SR, Baraibar J. Variations in etiology of ventilator-associated pneumonia across four treatment sites: Implications for antimicrobial
8. Kollef MH. What is ventilator-associated pneumonia and why is it important? Respir Care. 2005;50:714-21. [PubMed] [Google Scholar]
9. Park DR. The microbiology of ventilator-associated pneumonia. Respir Care. 2005;50:742-63. [PubMed] [Google Scholar]
10. Craven DE, Kunches LM,Klinsky V, LichtenbergDA, Make BJ,Mccabe WR. Risk factors for pneumonia and fatality in patients receiving continuous mechanical ventilation.