



SEPSIS AS A MORTALITY INDICATOR IN ACUTE KIDNEY INJURY PATIENTS IN ICU

Dr. V. R. Aditya Chainulu

Junior Resident, Department Of General Medicine, Sree Balaji Medical College And Hospital, Chennai

Dr. V. Padma*

Professor, Department Of General Medicine, Sree Balaji Medical College And Hospital, Chennai *Corresponding Author

Dr. Indhuja

Senior Resident, Department Of General Medicine, Sree Balaji Medical College And Hospital, Chennai.

ABSTRACT

Sepsis is an extreme and dysregulated inflammatory response to infection that is distant from the primary site of infection resulting in end-organ damage. Appearance of acute kidney injury (AKI) during sepsis raises patient morbidity, predicts higher death rates, has a significant effect on functions of multiple organs, is associated with a prolonged duration of stay in the intensive care unit, and hence warrants appropriate healthcare resources. Pathophysiology, diagnostic procedures and appropriate therapeutic interventions in sepsis are still highly debatable despite impressive advances in several medical fields. The overwhelmingly high mortality rate of sepsis is not reduced despite multiple immunomodulatory agents showing promise in preclinical studies. Major obstacles to progress in understanding, early diagnosis, and implementing effective therapeutic modalities in sepsis-induced AKI comprise minimal histopathological data, few animal models closely emulating human sepsis, and a relative lack of specialized diagnostic tools. Here we analyse 100 ICU patients who developed AKI and sepsis as a major etiological factor as well as mortality indicator.

KEYWORDS : Sepsis, Acute Kidney Injury, Mortality, Intensive Care Unit, Etiology.

INTRODUCTION

Acute Kidney Injury (AKI) is a complex condition that manifests in a number of clinical scenarios with its manifestations ranging from modest plasma creatinine escalation to anuric renal failure. It is often under-recognized and has severe consequences¹⁻⁴.

Recent epidemiological studies show the wide variation in etiology and risk factors^{1,5-7}, identify the increased mortality associated with this disease (especially when dialysis is required)^{1,4,6,8,9}, and indicate a relationship to the subsequent development of chronic kidney disease (CKD) and development of dialysis dependence^{1,4,8,10-12}.

Emerging evidence suggests that even minor serum creatinine changes are associated with higher in-patient mortality¹³⁻¹⁵. Over the past decades, AKI has been the focus of extensive clinical and basic research. A major limitation has been the absence of a universally recognized definition of AKI. Despite significant progress in understanding the biology and mechanism of AKI in animal models, there has been limited translation of this knowledge into improved patient management and outcomes.

Acute Kidney Injury Network (AKIN) is an independent collaborative network of experts selected by the participating societies to represent their field of expertise as well as their sponsoring organization. AKIN seeks to promote global, interdisciplinary and intersocietal partnerships in order to ensure success in the area of AKI and achieve the best results for patients with or at risk of AKI.

AIMS AND OBJECTIVES

To study and analyze the etiology of AKI and sepsis as a mortality indicator in AKI patients in ICU.

SAMPLE SIZE : 100

INCLUSION CRITERIA :

All patients admitted to ICU with AKI and those who developed AKI after admission to ICU.

EXCLUSION CRITERIA :

- Patients with pre-existing renal disease and with renal transplant.
- Patients with abnormal kidney size and abnormal cortico-medullary differentiation.

RESULTS AND DISCUSSION

100 patients have been analyzed in our research. There were 52 males and 48 females. The mean age was 59.5 years. In the sixth to eighth decade, the highest number of cases were reported. Mehta et al., in her study analysed a sample population which had 59% males and 41% were females with mean of 59.5 years.⁴⁶

Sepsis was found to be the leading cause of Acute Kidney Injury in the current study (40%) as has been time and again proven by multiple studies in the past like Muthusethupathi et al⁴⁷ in Chennai. This study has shown that among 40 patients with sepsis, 18 patients expired ($p < 0.001$) and sepsis was hence found to be an indicator of higher mortality rates than those without sepsis. Mortality due to septicemia was observed to be 75%. This was identical to the findings of Qutub HO, Saeed IA⁴⁸ in their three-year long prospective study (1996 to 1999) in which sepsis emerged as the leading cause of Acute Kidney Injury and also a major contributor for mortality. They also stated that prevention of sepsis would improve the prognosis of patients who got admitted into ICU due to any other cause.

In another study conducted by Agarwal et al⁴⁹ at CMC Vellore in 2004, similar observations were made in pediatric population vis-a-vis septicemia. It was found that mortality due to septicemia was 83% among ICU patients who had Acute Kidney Injury and was a major contributor to morbidity and mortality.




Figure 1 : Proportion of cases with sepsis

Table 1 : Outcome based analysis of sepsis

			Outcome		Total
			EXPIRED	RECOVERED	
Sepsis	PRESENT	Count	18	22	40
		% within Sepsis	45.0%	55.0%	100.0%
		% within Outcome	75.0%	28.9%	40.0%
		% of Total	18.0%	22.0%	40.0%
	ABSENT	Count	6	54	60
		% within Sepsis	10.0%	90.0%	100.0%
		% within Outcome	25.0%	71.1%	60.0%
		% of Total	6.0%	54.0%	60.0%
Total	Count	24	76	100	
	% within Sepsis	24.0%	76.0%	100.0%	
	% within Outcome	100.0%	100.0%	100.0%	
	% of Total	24.0%	76.0%	100.0%	

REFERENCES:

- Mehta RL, Pascual MT, Soroko S, Savage BR, Himmelfarb J, Ikizler TA, Paganini EP, Chertow GM, Program to Improve Care in Acute Renal Disease (PICARD). Spectrum of acute renal failure in the intensive care unit: the PICARD experience. *Kidney international*. 2004 Oct 1;66(4):1613-21.
- Palevsky PM. Epidemiology of acute renal failure: the tip of the iceberg.
- Ympa YP, Sakr Y, Reinhart K, Vincent JL. Has mortality from acute renal failure decreased? A systematic review of the literature. *The American journal of medicine*. 2005 Aug 1;118(8):827-32.
- Metnitz PG, Krenn CG, Steltzer H, Lang T, Ploder J, Lenz K, Le Gall JR, Druml W. Effect of acute renal failure requiring renal replacement therapy on outcome in critically ill patients. *Critical care medicine*. 2002 Sep 1;30(9):2051-8.
- Waikar SS, Curhan GC, Wald R, McCarthy EP, Chertow GM. Declining mortality in patients with acute renal failure, 1988 to 2002. *Journal of the American Society of Nephrology*. 2006 Apr 1;17(4):1143-50.
- Uchino S, Kellum JA, Bellomo R, Doig GS, Morimatsu H, Morgera S, Schetz M, Tan I, Bouman C, Macedo E, Gibney N. Acute renal failure in critically ill patients: a multinational, multicenter study. *Jama*. 2005 Aug 17;294(7):813-8.
- Liangos O, Wald R, O'Bell JW, Price L, Pereira BJ, Jaber BL. Epidemiology and outcomes of acute renal failure in hospitalized patients: a national survey. *Clinical journal of the American Society of Nephrology*. 2006 Jan 1;1(1):43-51.
- Clermont G, Acker CG, Angus DC, Sirio CA, Pinsky MR, Johnson JP. Renal failure in the ICU: comparison of the impact of acute renal failure and end-stage renal disease on ICU outcomes. *Kidney international*. 2002 Sep 1;62(3):986-96.
- Thakur CV, Worley S, Arrigain S, Yared JP, Paganini EP. Influence of renal dysfunction on mortality after cardiac surgery: modifying effect of preoperative renal function. *Kidney international*. 2005 Mar 1;67(3):1112-9.
- Druml W. Long term prognosis of patients with acute renal failure: is intensive care worth it?
- Liano F, Junco E, Madero R, Pascual J, Verde E. The spectrum of acute renal failure in the intensive care unit compared with that seen in other settings. *Kidney International-Supplements*. 1998 May 1;66(1):16-24.
- Mehta RL, Pascual MT, Soroko S, Chertow GM, PICARD Study Group. Diuretics, mortality, and nonrecovery of renal function in acute renal failure. *Jama*. 2002 Nov 27;288(20):2547-53.
- Chertow GM, Burdick E, Honour M, Bonventre V, Bates DW. Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *Journal of the American Society of Nephrology*. 2005 Nov 1;16(11):3365-70.
- Gruberg L, Mintz GS, Mehran R, Dangas G, Lansky AJ, Kent KM, Pichard AD, Satler LF, Leon MB. The prognostic implications of further renal function deterioration within 48 h of interventional coronary procedures in patients with pre-existent chronic renal insufficiency. *Journal of the American College of Cardiology*. 2000 Nov 1;36(5):1542-8.
- Lassnigg A, Schmidlin D, Mouhieddine M, Bachmann LM, Druml W, Bauer P, Hiesmayr M. Minimal changes of serum creatinine predict prognosis in patients after cardiothoracic surgery: a prospective cohort study. *Journal of the American Society of Nephrology*. 2004 Jun 1;15(6):1597-605.
- Qutub HO, Saeed IA. Acute renal failure in intensive care unit. *Saudi Med J*. 2001;22:999-1003.
- Agarwal I, Kirubakaran C, Markandeyulu V. Clinical profile and outcome of acute renal failure in South Indian children. *Journal of the Indian Medical Association*. 2004 Jul;102(7):353-4.
- Macedo E, Malhotra R, Bouchard J, Wynn SK, Mehta RL. Oliguria is an early predictor of higher mortality in critically ill patients. *Kidney international*. 2011 Oct 1;80(7):760-7.