Original Research Paper

Neurosurgery

Set Per		neurosurgery
Annual State	NON-ANEURYSMAL NON-TRAUMATIC SUBARACHN PATIENT CHARACTERISTICS, CLINICAL OUTCOME I CENTER EXPERIENCE IN 60 PATIEN	OID HEMORRHAGE: BASED ON A SINGLE- NTS
Subhasis Ghosh	Professor & Head Department Of Neurosurgery Institute Of Neurosciences, Kolkata West Bengal, In	[,] Ipgme&r And Bangun Idia.
Arvind Kumar Agrahari*	Mch Resident, Department Of Neurosurgery Ipgme Of Neurosciences, Kolkata West Bengal, India. *Con	ə&r And Bangur Institute rresponding Author
Shubhomitro Chaudhuri	Assistant Professor, Department Of Neurosurger Institute Of Neurosciences, Kolkata West Bengal, Ir	y Ipgme&r And Bangu Idia.
Sudipto Chatterjee	Professor, Department Of Neurosurgery Ipgme&r Neurosciences, Kolkata West Bengal, India.	And Bangur Institute O
Gitanjali Dutta	Associate Professor Department Of Neurosurger Institute Of Neurosciences, Kolkata West Bengal, In	y Ipgme&r And Bangu Idia.
Sanjeev Kumar Chhabra	Resident, Department Of Neurosurgery Ipgme&r . Neurosciences, Kolkata West Bengal, India.	And Bangur Institute O

ABSTRACT Objectives: Subarachnoid hemorrhage (SAH) with negative angiographic findings has a heterogeneous nature with variations in clinical course and outcome as compared to the aneurysmal SAH. It makes up to 15% of the spontaneous SAH and is characterized by milder clinical presentation. The purpose of this study was the analyses of risk factors, clinical observations, radiologic characteristics, and outcome in patients with nonaneurysmal SAH (NA-SAH) & Aneurysmal SAH.

Methods: From 2018 to 2020, data of 62 patients with non-aneurysmal SAH were prospectively entered into a database. All patients underwent repetitive cerebral DSA. Repeat DSA done to all patients. Outcome was assessed according to the modified Rankin Scale (mRS) (mRS 0–2 favorable vs. 3–6 unfavorable). Also, patients were divided in two groups according to the distribution of blood in the CT scan (perimesencephalic and non-perimesencephalic SAH).

Results: Out of 62 patients 02 patients comes to positive in repeat DSA. Out of 60 subjects, 21 (35%) were having nonperimesencephalic hemorrhage and 39 (65%) were having perimesencephalic hemorrhage. GCS grade ≤ 12 was reported among 23.81% of the subjects having non-perimesencephalic hemorrhage. Mean modified Fisher scale grading at admission among the study subjects was 2.92 ± 0.93 and 1.98 ± 0.87 with statistically significant difference as p<0.05. At the end of 6 month, unfavorable outcome was revealed only in one subject (4.8%) having nonperimesencephalic hemorrhage.

Conclusions: Patients suffering from non-aneurysmal SAH have better prognosis compared to aneurysm related SAH and poor admission status was the only independent predictor of unfavorable outcome in the multivariate analysis. Patients with a non-perimesencephalic SAH have an increased risk of a worse neurological outcome. These patients should be monitored attentively.

KEYWORDS : Non-aneurysmal, Aneurysm, Perimesencephalic, Non-perimesencephalic, , Subarachnoid hemorrhage, SAH, DSA

Background -

Spontaneous nonaneurysmal subarachnoid hemorrhage (NA-SAH) is a heterogeneous disease with variations in clinical course and outcome as compared to the aneurysmal SAH.[1]. The clinical course of the NA-SAH is not well defined in the literature. Few series exist to define the risk factors, clinical presentations, and outcomes in NA-SAH[2,3]. In 20 to 70% of patients with angiography negative SAH the blood distribution is described as perimesencephalic or prepontine [4]. Patients with perimesencephalic SAH are considered to achieve a good outcome and to have a lower risk of rebleeding [4,5]. Therefore, we aimed in our study to identify the risk factors, clinical and neurological status at the initial presentation, radiographic appearance, complications, and outcome of the patients with spontaneous NA-SAH at the time of admission and 6 months' follow-up. We conducted a retrospective study to investigate the clinical and functional outcomes of patients with spontaneous SAH in Npmn & PMN SAH in short-term follow-up.

Methods

After exclusion of patients with traumatic SAH we reviewed 62 patients with SAH admitted in the department of neurosurgery, Bangur Institute of Neurosciences Kolkata (BIN) & SSKM Hospital, IPGME & R, Kolkata from December 2018 to December 2020. The study comprised of spontaneous subarachnoid hemorrhage patients diagnosed with initial non-contrast CT followed by CTA/MRA Brain and Digital substraction angiography brain. Initial NCCT brain suggestive of subarachnoid hemorrhage and initial negative DSA patients can be divided into perimesencephalic and nonperimesencephalic pattern.

The data was collected by a preformed structured intervieweradministered questionnaire that was pretested with modifications made prior to its use in the study. The patients were interviewed that requests for the demographic, medical history and previous history of taking any medications, clinico-radiological status and clinical outcome. We assess the patients on detail history and clinical examinations. After this detail routine blood investigation with viral marker along with radiological investigation in form of NCCT brain, CT/MR angiography & Cerebral DSA. Secondary DSA positive patients excluded from study. Patients of non-aneurysmal spontaneous subarachnoid hemorrhage assessed by initial Hunt & Hess scale, GCS, modified fisher grade, WFNS grade (admission status) and modified Rankin scale for clinical outcome.

All patients admitted as case of spontaneous subarachnoid

in SSKM & IPGMER HOSPITAL Kolkata included in study but patients of traumatic SAH, lost to follow up, not giving consent , and diagnosed as structural or radiological aneurysm and AVM Or vascular malformation.

Patients were examined at admission, 6 week, 3 months & 6 months. Clinical outcome is studied with help of Modified Rankin Scale at regular interval 6 week, 3 months & 6 months .AT time of admission patients assess with help of GCS Score, Hunt & Hess scale, WFNS grading and bleeding pattern on Modified fisher grading on NCCT brain. Patients were divided into good grade (WFNS grades I – III) versus (vs.) poor grade (WFNS IV – V) on admission. Outcome was assessed according to the modified Rankin Scale (mRS) and stratified into favorable (mRS 0–2) vs. unfavorable (mRS 3–6) after 6 months.

Ethical Clearance:

The study protocol for all procedures was approved by the Institutional Review Board for Ethical Clearance of IPGMER, Kolkata and was performed in accordance with the Code of Ethics of the World Medical Association according to the Declaration of Helsinki of 1975, as revised in 2000. All patients were asked to sign a written consent form prior to commencement of the study.



Statistical Analysis

Data so collected was tabulated in an excel sheet, under the guidance of statistician. The means and standard deviations of the measurements per group were used for statistical analysis (SPSS 22.00 for windows; SPSS inc, Chicago, USA). Difference between two groups was determined using student t-test as well as chi square test and Fisher Exact Test. The level of significance was set at p < 0.05.

Results;-

Table 1 Patients characteristics

Charactersic	Perimesen-	Non-	Р
	cephalic	Perimesencep	value
	SAH(%)	halic SAH(%)	
No. of Patients	39(65%)	21(35%)	NS
Mean age	43.03 ± 15.44	45.67±11.39	NS
Male/Female	27/12	13/08	NS
Mean WFNS at	1.44 ± 0.72	$1.44 {\pm} 0.72$	0.04
admission			
GCS more or equal to	39(39)	21(16)	0.02
12 at admission			
Mean HHS at	1.87 ± 0.52	2±0.63	NS
admisssion			
Secondary DSA Positive	02(8.69%)	0	
Mean MRS scale at	1.98 ± 0.87	2.92 ± 0.93	0.36
admission			

Good status at	39(100%)	21(76.19%)	0.002
admission			
Favourable status at	39(100%)	20(95.2%0	Ns
admission			

Gender

Out of 60 subjects, 21 (35%) were having non-perimesencephalic hemorrhage and 39 (65%) were having perimesencephalic hemorrhage diagnosed on initial bleeding pattern on NCCT Brain. In our study, 61.9% and 69.23% of the subjects were male having non-perimesencephalic and perimesencephalic hemorrhage respectively. In both groups male were more than females.

Misdiagnosis

In our study, initially there were 23 subjects having nonperimesencephalic hemorrhage, out of which 2 (8.69%) turned out to be positive on second DSA. None of subjects with perimesencephalic hemorrhage turned out to be positive on second DSA.

In our study, headache was reported among all the subjects whether suffering from perimesencephalic or nonperimesencephalic hemorrhage. Vomiting and loss of consciousness was comparatively more in nonperimesencephalic hemorrhage group as compared to perimesencephalic hemorrhage with statistically significant difference as p<0.05. Neurological deficit and neck rigidity was comparable among both the groups. Hypertension, diabetes mellitus, antiplatelet & anticoagulant and smoking was found among 52.38%, 42.86%, 19.05%, 23.8% and 43.59%, 30.77%, 0%, 35.9% of the subjects having nonperimesencephalic and perimesencephalic hemorrhage respectively.

Table 2 Clinical symptoms

Symptoms	Perimesenceph	Non-	p value
	alic SAH	Perimesencephalic	
Headache	39	21	1
Vomiting	6	18	0.007*
Loss of	12	16	< 0.01*
Consciousness			
Neurological	1	3	0.12
Deficit			
Neck Rigidity	28	16	0.48

ADMISSION;-

Hunt & Hess grading 3 and 4 was reported among 9.52%, 4.76% and 2.56%, 0% of the subjects having nonperimesencephalic and perimesencephalic hemorrhage respectively. In this study, WFNS grading 3 and 4 was reported among 14.3%, 23.81% and 2.56%, 0% of the subjects having non-perimesencephalic and perimesencephalic hemorrhage respectively. In our study, GCS grade ≤ 12 was reported among 23.81% of the subjects having non-perimesencephalic hemorrhage. GCS grade 15 was found among 38.09% and 66.67% of the subjects having non-perimesencephalic and perimesencephalic hemorrhage respectively.

Table 3: Complications Among The Study Subjects

S.NO.	Complication	Non- PMN Type	PMN Type	p value
01	EVD Placement	02	00	NS
02	V .P. Shunt	01	00	NS
03	Early HYDROCEPHALUS	05	02	NS
04	Late Hydrocephalus	01	00	NS
05	Seizure	01	01	NS
06	Vasospasm	02	00	NS
07	Rebleed	00	00	NS

[7].

ble 4: Modified	Fisher Scal	e Grading
-----------------	-------------	-----------

Ta

Modified	Non	Perimesencephalic	P value
Fisher	Perimesencephalic	Spontaneous SAH	
Scale	Spontaneous SAH		
01	03	23	0.04*
02	02	02	
03	14	13	
04	02	01	
Mean±SD	2.92 ± 0.93	1.98 ± 0.87	0.036*

Figure 3-DSA Machine with monitor

RADIOLOGICAL FEATURES:-

In this study, mean modified Fisher scale grading at admission among the study subjects was 2.92 ± 0.93 and 1.98 ± 0.87 with statistically significant difference as p<0.05. Modified Fisher scale grading 1, 2, 3 and 4 was revealed among 14.29%, 9.52%, 66.67%, 9.52% and 58.97%, 5.13%, 33.3%, 2.56% of the subjects having non-perimesencephalic hemorrhage and perimesencephalic hemorrhage respectively. When modified Fisher scale grading at admission among the study subjects was compared according to non-perimesencephalic and perimesencephalic hemorrhage, it was found to be statistically significant as p<0.05.

Clinical outcomes;-

At admission, Moderate and slight disability was found among 3 (14.3%) and 5 (23.8%) subjects having nonperimesencephalic hemorrhage while no moderate/slight disability was reported among subjects having perimesencephalic hemorrhage. Most of the subjects (64.1%) with perimesencephalic hemorrhage didn't have any significant disability. When subjects were compared w.r.t. MR scale at different intervals among according to nonperimesencephalic and perimesencephalic hemorrhage, it was found to be statistically significant as p<0.05.

Complications ;-

Only 5.1% and 2.56% of the subjects with perimesencephalic hemorrhage were having complications viz. early hydrocephalus and seizure respectively. Complications viz. EVD placement, VP shunt, early hydrocephalus, late hydrocephalus, seizure and vasospasm was revealed among 9.52%, 4.76%, 23.8%, 4.76%, 4.76% and 9.52% of the subjects with nonperimesencephalic hemorrhage respectively. In overall study complication in form of hydrocephalus, need of VP shunt or Need of EVD or Vasospasm was more common among non-perimesencephalic however it comes to be insignificant. No mortality was seen in this study also no case of rebleed was seen among any patients

DISCUSSION:-

About 15% of patients with spontaneous SAH approximately have negative findings regarding the bleeding source.[1]. SAH patients with negative initial DSA findings, usually defined as NASAH, can be a management challenge as the optimal management scheme for these patients remains controversial. The present study analyzes the data of SAH patients with negative initial DSA findings from our center. We divided those patients into the two subgroups of PMNSAH and nPMN-SAH. It is wildly accepted that NASAH patients have a more favorable clinical course and a lower incidence of complications when compared to those with aSAH [6]. The subgroup of PMN-SAH was first described by van Gijn et al. Spontaneous subarachnoid hemorrhage (SAH), characterized by bleeding into the subarachnoid space in the absence of trauma, is most often caused by the rupture of an intracranial aneurysm[8,9]. A diagnosis of nPMNSAH is associated with higher rate of complications, such as hydrocephalus, cerebral vasospasm, and cerebral infarction [10].

Patients suffering from non-aneurysmal SAH are usually in good clinical condition [2,4,11,12,13]. Males were comparatively more as compared to females in the present study. Liang Xu et al [14], Asma Bashir et al [15] Cody L. Nesvick et $al^{_{[16]}}$ males were comparatively more as compared to females in the present study. Mean age among subjects having nonperimesencephalic and perimesencephalic hemorrhage was 45.67 ± 11.39 and 43.03 ± 15.44 years respectively. Cody L. Nesvick et al^[16]. Alhoobi, et al^[17]. Akcakaya MO et al^[18] in their study found approximately similar mean age among the study subjects. 2 patients means 8.69% turned out to be positive on second DSA. None of subjects with perimesencephalic hemorrhage turned out to be positive on second DSA. In previous studies, the misdiagnosis rate of the initial DSA in nPMN-SAH patients varied greatly (misdiagnose rate from 4.7% to 45.9%), which is likely due to several factors, as previously mentioned. After pooling all the published studies, the overall misdiagnose rate was 12.5%, which is little higher than the findings at our institution.

In our study, headache was reported among all the subjects whether suffering from perimesencephalic or nonperimesencephalic hemorrhage. Vomiting and loss of consciousness was comparatively more in nonperimesencephalic hemorrhage group as compared to perimesencephalic hemorrhage with statistically significant difference as p<0.05. Neurological deficit and neck rigidity was comparable among both the groups.

In a study by Alhoobi, et al¹²⁷, headache in 73 patients (94.8%), along with vomiting in 24 patients (31.6%) and low level of consciousness in 8 patients 10.4%, all (100%) of them were of NPM. Asma Bashir et al¹⁵ in their study revealed that seven (9.5%) patients briefly lost consciousness at the time of ictus; among them one (1.4%) had a seizure. Two (2.7%) patients presented with abducens nerve palsy, and one (1.4%) with oculomotor nerve palsy. Hypertension, diabetes mellitus, antiplatelet & anticoagulant and smoking was found among 52.38%, 42.86%, 19.05%, 23.8% and 43.59%, 30.77%, 0%, 35.9% of the subjects having non-perimesencephalic and perimesencephalic hemorrhage respectively in the present study. Liang Xu et al¹⁹ in their study revealed similar findings. David P. Bray et al¹²⁰ also noted that 40 % patients received antiplatelet drug and associated blood volume is significantly more in npmn than PMN type SAH.

When Hunt & Hess grading at admission among the study subjects was compared according to non-perimesencephalic and perimesencephalic hemorrhage, it was found to be statistically insignificant as p>0.05. Similarly Asma Bashir et al¹¹⁵ in their study reported that HH grades 3 or above were mostly seen in patients with nPMSAH.

When WFNS grading at admission among the study subjects was compared according to non-perimesencephalic and perimesencephalic hemorrhage, it was found to be statistically significant as p<0.05. Similarly Asma Bashir et al¹⁵⁵ in their study reported that WFNS grades 3 or above were mostly seen in patients with nPMSAH.

When GCS grading at admission among the study subjects was compared according to non-perimesencephalic and perimesencephalic hemorrhage, it was found to be statistically significant as p<0.05. In a study by Alhoobi, et al¹¹⁷, patients with NPM-SAH had initial severe symptoms with low GCS (GCS \leq 12) as all patients with WFNS Grade IV had NPM SAH, and 3 of them had GCS <10 as compared to the PM-SAH. In this study, mean modified Fisher scale grading at admission among the study subjects was 2.92±0.93 and 1.98±0.87 with statistically significant difference as p<0.05. Akcakaya MO et al¹⁶ in their study too reported significant differences in Fisher grades were found between study groups at admission.

At admission, Moderate and slight disability was found among 3 (14.3%) and 5 (23.8%) subjects having nonperimesencephalic hemorrhage while no moderate/slight disability was reported among subjects having perimesencephalic hemorrhage. Most of the subjects (64.1%) with perimesencephalic hemorrhage didn't have any significant disability. Similarly Asma Bashir et al^[15] in their study found that at discharge 11% of the subjects with nonperimesencephalic hemorrhage were having 3-6 mRS at discharge. Juergen Konczella et al^[21] also reported similar type of finding. . Complications viz. EVD placement, VP shunt, early hydrocephalus, late hydrocephalus, seizure and vasospasm was revealed among patients in study. In a study by Jobyna Whiting et al^[19], hydrocephalus necessitating temporary CSF diversion developed in 22 of these patients (25%); 12 (13%) ultimately required permanent CSF diversion. Clinically significant vasospasm developed in 4 patients (4%), and 2 (2%) had cerebral infarctions. Three patients (3%) died.Good and bad outcome status among the stud subjects was found among 76.19%, 23.81% and 100%, 0% of the subjects having non-perimesencephalic and perimesencephalic hemorrhage respectively with statistically significant difference. Juergen Konczalla et al^[21] also reported more good grade at admission in perimesencephalic vs. nonperimesencephalic type SAH.

Conclusion;-

Perimesencephalic and non-perimesencephalic SAH have similar presentation status. Favorable outcome is achieved in most patients. Poor admission status was the only independent predictor of unfavorable outcome in patients with spontaneous non-aneurysmal SAH.Managing SAH patients with negative initial DSA findings can be challenging. Based on the results of our present study and a review of the pertinent literature, the PMN-SAH subgroup usually has a benign clinical course and a repeat DSA very seldom reveals a ruptured intracranial aneurysm. More importantly, nPMN-SAH patients are associated with higher complication rate and higher incidence of an aneurysm. Therefore, we strongly recommend a repeat DSA in patients with nPMN-SAH pattern on initial imaging.

REFERENCES

- Van Gijn J, Rinkel GJ. Subarachnoid haemorrhage: Diagnosis, causes and management. Brain 2001;124:249-78.
- Gupta SK, Gupta R, Khosla VK, Mohindra S, Chhabra R, Khandelwal N, et al. Nonaneurysmal nonperimesencephalic subarachnoid hemorrhage: Is it a benign entity? Surg Neurol 2009;71:566-71
- Renou P, Tourdias T, Fleury O, Debruxelles S, Rouanet F, Sibon I. Atraumatic nonaneurysmal sulcal subarachnoid hemorrhages: A diagnostic workup based on a case series. Cerebrovasc Dis 2012;34:147-52
- Ildan F, Tuna M, Erman T, Gocer AI, Cetinalp E: Prognosis and prognostic factors in nonaneurysmal perimesencephalic hemorrhage: a follow-up study in 29 patients. Surg Neurol 2002, 57:160–165.
- Van Gijn J, van Dongen KJ, Vermeulen M, Hijdra A: Perimesencephalic hemorrhage: a nonaneurysmal and benign form of subarachnoid hemorrhage. Neurology 1985, 35:493–497.
- A. Kapadia, T. A. Schweizer, J. Spears, M. Cusimano, and R. L. Macdonald, "Nonaneurysmal perimesencephalic subarachnoid hemorrhage: diagnosis, pathophysiology, clinical characteristics, and long-term outcome," World Neurosurgery, vol. 82, no. 6, pp. 1131–1143, 2014.
 J. van Gijn, K. J. van Dongen, M. Vermeulen, and A. Hijdra,
- J. van Gijn, K. J. van Dongen, M. Vermeulen, and A. Hijdra, "Perimesencephalic hemorrhage: a nonaneurysmal and benign form of subarachnoid hemorrhage," Neurology, vol. 35, no. 4, pp. 493–497, 1985.
- Lucke-Wold BP, Logsdon AF, Manoranjan B, Turner RC, McConnell E, Vates GE, Huber JD, Rosen CL, Simard JM. Aneurysmal subarachnoid hemorrhage and neuroinflammation: a comprehensive review. International journal of molecular sciences. 2016;17(4):497.

- Hasegawa Y, Suzuki H, Uekawa K, Kawano T, Kim-Mitsuyama S. Characteristics of cerebrovascular injury in the hyperacute phase after induced severe subarachnoid hemorrhage. Translational stroke research. 2015;6(6):458-66.
- Y. Kong, J. H. Zhang, and X. Qin, "Perimesencephalic subarachnoid hemorrhage: risk factors, clinical presentations, and outcome," Acta Neurochirurgica Supplement, vol. 110, part 1, pp. 197–201, 2011
- Beseoglu K, Pannes S, Steiger HJ, Hanggi D: Long-term outcome and quality of life after nonaneurysmal subarachnoid hemorrhage. Acta Neurochir (Wien) 2010, 152:409–416.
- Nayak S, Kunz AB, Kieslinger K, Ladurner G, Killer M: Classification of nonaneurysmal subarachnoid haemorrhage: CT correlation to the clinical outcome. Clin Radiol 2010, 65:623–628.
- Pyysalo LM, Niskakangas TT, Keski-Nisula LH, Kahara VJ, Ohman JE: Long term outcome after subarachnoid haemorrhage of unknown aetiology. J Neurol Neurosurg Psychiatry 2011, 82:1264–1266
- Xu L, Fang Y, Shi X, Chen X, Yu J, Sun Z, Zhang J, Xu J. Management of spontaneous subarachnoid hemorrhage patients with negative initial digital subtraction angiogram findings: conservative or aggressive?. BioMed research international. 2017 May 2;2017.
- Bashir A, Mikkelsen R, Sørensen L, Sunde N. Non-aneurysmal subarachnoid hemorrhage: When is a second angiography indicated? Neuroradiol J. 2018;31(3):244-252.
- Nesvick CL, Oushy S, Rinaldo L, Wijdicks EF, Lanzino G, Rabinstein AA. Clinical complications and outcomes of angiographically negative subarachnoid hemorrhage. Neurology. 2019;92(20):e2385-94.
- Alhoobi M, Abu-Qadous F, Khan M, et al. Ten Years' Experiences in the Treatment of Nonaneurysmal Subarachnoid Hemorrhage: A Retrospective Analysis of Outcome Parameters in a Single-Center Study. Asian Journal of Neurosurgery. 2020;15(2):315.
- Akcakaya MO, Aydoseli A, Aras Y, et al. Clinical course of nontraumatic nonaneurysmal subarachnoid hemorrhage: A single institution experience over 10 years and review of the contemporary literature. Turk Neurosurg. 2017;27(5):732-42.
- Whiting J, Reavey-Cantwell J, Velat G, Fautheree G, Firment C, Lewis S, Hoh B. Clinical course of nontraumatic, nonaneurysmal subarachnoid hemorrhage: a single-institution experience. Neurosurgical focus. 2009;26(5):E21.
- Bray DP, Ellis JA, Lavine SD, Meyers PM, Connolly Jr ES. Analysis of hemorrhage volumes after angiogram-negative subarachnoid hemorrhage. World neurosurgery. 2016;94:453-7.
- Konczalla J, Platz J, Schuss P, Vatter H, Seifert V, Güresir E. Non-aneurysmal non-traumatic subarachnoid hemorrhage: patient characteristics, clinical outcome and prognostic factors based on a single-center experience in 125 patients. BMC neurology. 2014;14(1):1-8.