



PROXIMAL FEMORAL NAIL- OUTCOME AND COMPLICATIONS: A RETROSPECTIVE AND PROSPECTIVE STUDY OF 115 CASES OF PROXIMAL FEMORAL FRACTURES.

Dr.Bharath reddy Danda

Senior Resident, Department of orthopaedics, GMC, Suryapet, Telangana.

Dr Ramesh Naik L*

Assistant Professor, Department of Orthopaedics, GMC, suryapet.

*Corresponding Author

ABSTRACT

Background: Fractures of the proximal femur are relatively common injuries in adults and common source of morbidity and mortality among the elderly. Fractures of the proximal femur include trochanteric and subtrochanteric fractures. The present study was designed to evaluate and analyze the role of proximal femoral nail (PFN) in the treatment of proximal femoral fractures.

Results: In this study at 6 months follow up, union was achieved in 115 cases, open reduction was performed in 11% of cases (14 cases). Technical and mechanical complications were noted in 21% cases (27 cases). Reoperation rate was 4% (Five cases). According to Salvati and Wilson scoring system excellent results were seen in 36% of cases (45 cases), good results in 46% cases (58 cases), fair result in 13% cases (16 cases) and poor results in 5% cases (6 cases).

KEYWORDS : Proximal femoral nail, Intertrochanteric fractures, Subtrochanteric fractures, Proximal femur fracture

INTRODUCTION

Hip fractures are the most devastating injuries in all the age groups. Extracapsular fractures (intertrochanteric and subtrochanteric fractures) primarily involve cortical and compact cancellous bone.^{1,2} For the treatment of the unstable proximal femoral fracture with lacking medial support and the intertrochanteric fractures, two principal options exist. Either any kind of a sliding neck screw connected to a plate at the lateral femoral cortex and inserted after semi-open reduction or a sliding neck screw penetrating the head-neck fragment through an intramedullary nail implanted via a semiclosed technique.³

MATERIALS AND METHODS:

This was a Retrospective and prospective study on cases of proximal femoral fractures treated between august 2020 to July 2021, who were admitted in Department of Orthopaedics, Medical Government medical College, Suryapet. Fractures were classified according to AO/ASIF- classification.⁵ 125 cases were followed at regular intervals and final assessment was done at 6 months. The Salvati and Wilson score of hip function was used at the last clinical assessment.⁶

RESULTS

A total of 115 patients were enrolled for the study. There were 55 males and 70 females, with an average age of 63 years (range: 31 to 93 years). Domestic fall was the main injury mechanism accounting for 64% cases (80 cases). Forty five patients (36%) had RTA as mode of injury. Right side fractures were recorded in sixty five patients (52%) and left side fractures in sixty patients (48%). Ninety five patients (76%) had intertrochanteric fractures of femur and thirty patients (24%) had fractures of subtrochanteric femur. According to AO/OTA classification most intertrochanteric fractures were in Type 31A2.2 (36 cases), in subtrochanteric fractures most common type was 32A1.1 (20 cases). Associated injuries included twenty one patients (16.8%) of other bones fracture, blunt trauma abdomen in five patients (4%), head injury in four patients (3.2%) and facial injuries also in four patients (3.2%). The average time from injury to surgery was 7 days (range: 1 to 13 days). Average duration of surgery was 88 min (range 45 to 145 min). Closed reduction was achieved in eighty nine percent cases (111 cases). Open reduction was performed in eleven percent cases (14 cases). Mean intraoperative blood loss was 126 ml. In our study it was observed that

Table 1: Intraoperative complications of PFN.

S. No.	Intraoperative complications	No. of subjects
1	Failure to achieve closed reduction	14
2	Fracture of lateral cortex (shattering)	1
3	Varus angulation	0
4	Failure to put antirotation screw	7
5	Failure to lock distally	0
6	Jamming of nail	0
7	Drill bit breakage	0
8	Guide wire breakage	2

Table 2: Delayed complications.

S. No.	Delayed complications	No. of subjects
1	Reverse Z effect	2
2	Shortening	3
3	Loosening of hip pin causing persistent pain in lateral surface of thigh	8
4	Z- effect	3
5	Stiffness of hip	11

DISCUSSIONS

Unstable fractures of the proximal femur represent a significant challenge to the trauma surgeon. Surgical fixation is often technically difficult and poor surgical technique may lead to failure of primary fixation.^{7,8} The best treatment for these fractures remains controversial. DHS fixation is widely preferred but failure of fixation still occurs in up to 20% of cases.³ Common causes of fixation failure include fracture instability, osteoporosis, lack of anatomic reduction, implant failure, and incorrect placement of the lag screw in the femoral head (leading to cutting out of the screw).⁹ Cephalomedullary femoral reconstruction nails with a trochanteric entry point are biomechanically stronger than extramedullary implants.¹⁰ In unstable proximal femoral fractures, control of axial telescoping and rotational stability are essential. Intramedullary implants inserted in a less-invasive manner are better tolerated by the elderly.¹¹ A new device was developed by AO/ASIF: the proximal femoral nail (PFN), with an additional antirotational hip pin preventing rotation and collapse of the head-neck fragment and an especially shaped tip together with a smaller distal shaft diameter resulting in less stress concentration at the tip.¹²

Velasco and Comfort found that 63% of subtrochanteric fracture occurred in patients from 51 to more than 70 years old and 24% of patients between 17 to 50 years old.¹³ In a study by

Babst et al in 1998 in intertrochanteric fractures, mean age was 79.7 years (range 39-98 years).¹⁴ According to Klinger et al in 2005 the mean age was 74 years ranging from (27 to 98 years) in patients who were treated either with DHS or proximal femoral nail.¹⁵ Alyassari et al studied seventy patients and the average age was 84 years showing trochanteric fracture are more common in higher age group.¹⁶ In our study fifty patients (40%) were between 20-60 years and 66% of subtrochanteric fractures were below the age of 60 years. The mean age of unstable intertrochanteric fractures was 67.73 years with range from 41 years to 95 years which is slightly towards the older age group, mainly due to Osteoporosis.

Simmermacher in their study the mean duration of surgery (skin to skin) was 68.7 min (range 25-240 min).³ Pajarinen et al in their comparative study of DHS and PFN in proximal femoral fracture, the average time of surgery in DHS was 45 min (range 20-105 min) and in PFN was 55 min (35-200 min).¹⁷ Wang in their study, the average operating time was 90 min (Range 60-155 min).¹⁸ In our study duration of surgery was longer in the initial operated cases. With frequent use of proximal femoral nail surgery the duration decreased. In our study average duration of surgery was 88.24 minutes.

Fogagnolo et al reported 46 patients with an average rate of intra operative technical or mechanical complications of 23.4%, mostly problems with the distal nail locking and fracture of the lateral wall of the greater trochanter.¹⁹

Kamboj et al studied 30 cases, in one case with trochanteric fracture extending to diaphysis encircling wiring was done. One patient got intra operative fracture shaft of femur, three patients had poor placement of screw. The closed reduction was tried in all cases and achieved in 17 patients, in the rest of 13 cases fracture had to be opened. In their study, due to smaller diameter of the neck of Indian femora they were not able to pass anti rotational hip pin in four patients.²⁰ Alyassari et al in their study, two cases required open reduction, distal locking was difficult in three cases, nail insertion was difficult in one patient.¹⁶ In our study, there was shattering of the proximal fracture fragment in one patient while insertion of the nail. In fourteen patients, it was not possible to achieve closed reduction, so open reduction was done by opening the fracture site. In seven patients it was difficult to put the derotation screws. In three cases, it could not be accommodated in the neck after putting the neck screw and in other four cases, it had to be removed after inserting as it was penetrating the superior cortex of the neck. This suggests that in Indian population the neck of femur is not broad. In Two patients, there were guide wire breakages while reaming over guide wires in femoral neck.

Pajarinen et al in their study of 83 patients, there was one case of heterotopic ossification corresponding to Brooker class 4, where PFN was used.¹⁷ Werner et al was the first who introduced the term Z-effect, detected in five (7.1%) of 70 cases. The incidence of cut-out of the neck screw in this study was 8.6%. The Z-effect phenomenon is referred as a characteristic sliding of the proximal screws to opposite directions during the postoperative weight-bearing period.²¹ Reverse Z-effect described by Boldin et al occurred with movement of the hip pin towards the lateral side, which required early removal. In their prospective study of 55 patients with unstable intertrochanteric or subtrochanteric fractures, they had three cases with Z-effect and two with reverse Z-effect.²² Fogagnolo et al, who reported 46 patients with an average rate of intraoperative technical or mechanical complications of 23.4%. They also reported two implant failures and one fracture below the tip of the nail. They also reported heterotopic ossification in two patients fixed with PFN.¹⁹ Simmermacher et al in a clinical multicenter

study, reported technical failures of the PFN after poor reduction, malrotation or wrong choice of screws in 5% of the cases. A cut-out of the neck screw occurred in 0.6%.³ In our series there was shortening in three patients. In one patient, fracture was comminuted which caused shortening >2 cm on healing while in two patients it was of <2 cm where there was inadequate restoration of alignment and there was no medial buttressing that led to shortening. There was five cases of implant failure, three cases with 'Z Effect' and two cases with 'reverse Z effect'. Revision Surgeries were done in these cases. In eight patients, there was loosening of hip pin which caused persistent pain in lateral surface of thigh. In eleven patients, stiffness of hip joint was present.

REFERENCES

1. LaVelle GD. Fractures and Dislocations of the Hip. In: Canale ST, Beaty JH, editors. Campbell's operative orthopedics. 11th ed. Philadelphia: Mosby Elsevier publishers; 2008: 3237-3296.
2. Weinlein JC. Fractures and Dislocations of the Hip: Campbell's operative orthopedics. 12th ed. Elsevier; 2013: 2725.
3. Simmermacher RK, Bosch AM, Van der Werken C. The AO/ASIF-proximal femoral nail (PFN): A new device for the treatment of unstable proximal femoral fractures. *Injury*. 1999;30:327-32.
4. Prins S, Letsch R, BuÈscher D. Gamma-nagel und Classic-nagel (intramedulläre Stabilisierung) versus DHS (extramedulläre Stabilisierung) bei proximalen Femurfracturen. *Hefte Unfallchir*. 1996;262:14.
5. Mueller ME, Nazarian S, Koch P, Schatzker J, editors. The comprehensive classification of fractures of long bones. Heidelberg: Springer-Verlag Berlin; 1990: 120-121.
6. Bridle SH, Patel AD, Bircher M, Calvert PT. Fixation of intertrochanteric fractures of the femur. *J Bone Joint Surg*. 1991;73:330-4.
7. Baumgaertner MR, Curtin SL, Lindskog DM, Keggi JM. The value of the tip-apex distance in predicting failure of fixation of peritrochanteric fractures of the hip. *J Bone Joint Surg*. 1995;77:1058-64.
8. Sierra RJ, Cabanela ME. Conversion of failed hip hemi-arthroplasties after femoral neck fractures. *Clin Orthop Relat Res*. 2002;399:129-39.
9. Simpson AH, Varty K, Dodd CA. Sliding hip screws: modes of failure. *Injury*. 1989;20:227-31.
10. Moen CM, Verhofstad MH, Bley RL, van der Werken C. Soft tissue injury related to choice of entry point in antegrade femoral nailing: piriform fossa or greater trochanter tip. *Injury*. 2005;36:1337-4.
11. Baumgaertner MR, Curtin SL, Lindskog DM. Intramedullary versus extramedullary fixation for the treatment of intertrochanteric hip fractures. *Clin Orthop Relat Res*. 1998;348:87-94.
12. Euler E, Huber St, Heining S, Schweiberer L. Spannungsoptische Untersuchung unterschiedlicher Stabilisierungsverfahren bei peritrochanterischen Femurfracturen. *Hefte Unfallchir*. 1996;262:2.
13. Velasco RU, Comfort TH. Analysis of treatment problems in subtrochanteric fractures of the femur. *J Trauma*. 1978;18(7):513-23.
14. Babst R, Renner N, Bieder MM, Rosso R, Heberer M, Harder F, Reggioni P. Clinical results using the trochanteric stabilizing plate: the modular extension of the dynamic hip screw for internal fixation of intertrochanteric fractures. *J Orthop Trauma*. 1998;12(6):392-99.
15. Klinger HM, Baums HM, Eckert M, Neugebauer R. A comparative study of unstable per and intertrochanteric femoral fractures with DHS and PFN and TSP. *Zentralbl Chir*. 2005;130(4):301-6.
16. Al-yassari G, Langstaff RJ, Jones JW, Al-Lami M. The AO/ASIF proximal femoral nail (PFN) for the treatment of unstable trochanteric femoral fracture. *Injury*. 2002;33(5):395-9.
17. Pajarinen J, Lindahl J, Michelsson O, Savolainen V, Hirvensalo E. Peritrochanteric femoral fractures treated with a dynamic hip screw or a proximal femoral nail. A randomised study comparing post-operative rehabilitation. *J Bone Joint Surg Br*. 2005;87(1):76-81.