



## KNOWLEDGE, ATTITUDE AND PRACTICES ABOUT CONTRACEPTIVE AMONG REPRODUCTIVE AGE GROUP FEMALES

**Dr Mamta Choudhary\***

Senior resident, Department of Obstetrics & Gynaecology, Govt. S.K. Medical College, Sikar (Rajasthan). \*Corresponding Author

**Dr. Bajrang Lal Rar**

Professor, Department of Obstetrics & Gynaecology, Govt. S.K. Medical College, Sikar (Rajasthan).

### ABSTRACT

**Background-** India is the second most populous country in the world having a rapidly growing population which is currently increasing at the rate of 16 million each year. Uncontrolled population growth is recognised as the single most important impediment to national development. Aim of current study was to assess the knowledge, attitude & practices of contraceptives among married women of reproductive age group.

**Methods-** A cross sectional study was conducted on females between 15-45 years served as inclusion criteria at Department of **Obstetrics & Gynaecology-** Govt. S.K. Medical College, Sikar, while unmarried females were excluded. The participation was on voluntary basis. Questions regarding factors responsible for non use of contraception were also asked. All data were analyzed by Epi-info software.

**Results-** Out of 100 women, 64.00% had knowledge about family planning. 61.00% knew about condom, followed by OCPs & other methods. Contraceptive usage in our study was 62.00%.

**Conclusions-** Effort should be made to educate the public about the safety and convenience of modern, long term, reversible methods of contraception among both in health care professional and public.

**KEYWORDS :** Knowledge, Attitude, Practice, Contraception

### INTRODUCTION

India is the second most populous country in the world having a rapidly growing population which is currently increasing at the rate of 16 million each year.<sup>1</sup> Uncontrolled population growth is recognised as the single most important impediment to national development. Instead of the fact that India was the first country in the world to implement a national population control programme in 1952, we are still struggling to contain baby boom. A lot of efforts and resources have gone into the national family welfare programme but the returns are not commensurate with the inputs to control the population.<sup>2</sup>

In developing countries Couple Protection Rate (CPR) is still very less. According to national family welfare statistics 2011, CPR in India is 40.4%. In Pakistan CPR was 30% in 2011 while most developed country like USA has 71% CPR for all methods.<sup>3,4</sup>

Now a days fertility rate has come down due to wide spread use of family planning methods, whereas some developing countries like India, Pakistan, the use of contraception is still low because of lack of awareness, education, religion, cultural, economic and political barrier. Proper counseling for family planning necessary because 27% of women are at risk of unplanned pregnancy.<sup>5</sup>

The essential aim of family planning is to prevent the unwanted pregnancies. An unwanted pregnancy may lead to an induced abortion. From the point of view of health, abortion outside the medical setting is one of the most dangerous consequences of unwanted pregnancy.<sup>6</sup> Keeping this in mind one cross sectional study was carried out to assess the knowledge, attitude and practices of contraceptives of reproductive females.

### MATERIAL AND METHODS

**Study type-** Cross-sectional study **Inclusion criteria-** Married women 15-45 yrs age group and willing to participate in the study.

### EXCLUSION CRITERIA-

Unmarried, age more than 45 yrs and not willing to participate study.

### METHOD OF DATA COLLECTION-

A cross sectional study was conducted on females between 15-45 years served as inclusion criteria at Department of Obstetrics & Gynaecology, Govt. S.K. Medical College, Sikar, while unmarried females were excluded. The participation was on voluntary basis. Questions regarding factors responsible for non use of contraception were also asked. All data were analyzed by Epi-info software.

### RESULTS

A total of 100 married females of reproductive age were enrolled in the study. The study shows, mean age of respondent was  $29.64 \pm 9.12$  years.

**Table 1: Knowledge regarding different contraceptive methods**

Contraceptive methods	Number	Percentage
Condom	61	61.00%
OCP	60	60.00%
IUCD	51	51.00%
Injectable	6	6.00%
Tubectomy	48	48.00%

Out of 100 women, 64 (64.00%) had knowledge about family planning. 61.00% knew about condom, followed by OCPs & other methods.

**Table 2: Knowledge regarding emergency contraceptive methods**

Knowledge regarding emergency contraceptive methods	Number	Percentage
Yes	7	7.00%
No	93	93.00%

Only 7.00% Women were aware of emergency contraceptive. 68.00% respondent thought that contraceptives were used to prevent pregnancy and about 12.0% thought that they could be used to prevent infections like AIDS. Only 9.00% thought that they could be used to control birth interval.

**Table 3: Contraceptive practice among respondents**

Contraceptive methods	Number	Percentage
Condom	51	51.00%
OCP	40	40.00%
IUCD	30	30.00%

Injectable	0	0.00%
Tubectomy	20	20.00%
Not use any methods	38	38.00%

Contraceptive usage in our study was 62.00%. The gap between awareness and practices are seen to be prevalent across different reasons, where people are aware but reluctant to practices. The most common reason for non practice of contraception was fear of side effects. Other reasons for non practice were non access to health facility, preference of male child, religious beliefs, cost, and family pressure. Some respondent also felt that the process of acquiring contraceptive is often embarrassing.

## DISCUSSION

64.00% were aware of at least one of family planning methods in our study. Ambareen Khan et al<sup>7</sup> mentioned that 81% had awareness regarding any method of contraception. Tuladhara H et al<sup>8</sup> also observed that the most common source of information on contraception was media (55.5%), and both printed and electronic.

Contraceptive usage in our study was 62.00%. Sunita Ghike<sup>9</sup> also mentioned various for non-use of contraceptive methods. The main reason 59% were pressure from family that is from husband, in-laws, son preference and physical pressure.

Family planning services need to provide a range of quality method that can allow women to either limit or space birth and to fulfill the need of women with differing socio-demographic characteristics. In our study though majority of interviewer women had knowledge about family planning methods but practicing is still low because of lack of education, cultural, religion, economical and political barriers.

## CONCLUSION

Despite the high rate of sexual activity in the study group, the contraceptive usage rate is low. There is a need for aggressive advocacy about female reproductive health and dissemination of information on family planning methods among the reproductive females. Fear of side effects of contraception is one significant reasons for low compliance. This can be decreased by proper selection of contraception before starting its use and adequate follow up of women using contraception by the family planning services.

## REFERENCES

1. Sunita TH, Desai RM. Knowledge, attitude and practice of contraception among women attending a tertiary care hospital in India. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2013;2(2):172-176.
2. Renjhen P, Kumar A, Pattanshetty S, Sagir A, Samarasinghe CM. A study on knowledge, attitude and practice of Contraception among college students in Sikkim, India. *J Turkish-German Gynecol Assoc*. 2010;11:78-81.
3. WHO. Reproductive health indicator-guidelines for their generation, interpretation and analysis for global monitoring. In: WHO, eds. *WHO Guideline*. Geneva. WHO, Reproductive Health and Research; 2006: 1-63.
4. Bibi S, Memon A, Memon Z. Contraceptive knowledge and practices in two districts of Sindh Pakistan-A hospital based survey. *JPMA*. 2008;58:254-7.
5. Musarrat Jabeen, Fouzia Gul, Farmanullah Wazir, Nargis Javed. Knowledge, attitude and practices of Contraception in women of reproductive age. *Gomal J Med Sci*. 2011;9(2):223-9.
6. Tsui AO, McDonald-Mosley R, Burke AE. Family planning and the burden of unintended pregnancies. *Epidemiol Rev*. 2010;32:152-74.
7. Khan A, Hashmi HA, Naqvi Z. Awareness and Practice of Contraception Among Child Bearing Age Women. *Journal of Surgery Pakistan (International)*. 2011;16(4):179-182.
8. Tuladhara H, Marahatta R. Awareness and practice of family planning methods in women attending gyne OPD at Nepal medical college teaching hospital. *Nepal Med Coll J*. 2008;10(3):184-191.
9. Ghike S, Joshi S, Bhalerao A, Kawthalkar A. Awareness and Contraception Practices among Women An Indian Rural Experience. *Journal of South Asian Federation of Obstetrics and Gynecology*. 2010;2(1):19-21.