



## EVALUATION OF BLEEDING PER VAGINA AMONG POST-MENOPAUSAL PATIENTS ATTENDING A TERTIARY CARE CENTRE IN BIHAR

**Priti Kumari**

Senior Resident, Department Of Obstetrics And Gynecology , JLNMCB, Bhagalpur, Bihar

**Usha Kumari\***

Professor, Department Of Obstetrics And Gynecology, JLNMCB, Bhagalpur, Bihar. \*Corresponding Author

### ABSTRACT

World Health Organization (WHO) defines menopause as cessation of menstruation permanently for a period of more than one year, which is resulted from loss of ovarian activity Postmenopausal bleeding (PMB) is defined as bleeding from the genital tract, more than 12 months after the last menstrual period in a woman not on hormone replacement. **METHODOLOGY:** This descriptive study was carried out on 100 patients at Obstetrics and Gynaecology department of Jawahar Lal Nehru Medical College& Hospital during July 2020 to June 2021. **RESULT:** In the present series, 7 patients (14%) had malignancy, including endometrial carcinoma 4% and cervical carcinoma 10%, 27 patients (54%) had benign pathology, 9 had hyperplasia (18%) and 7 patients (14%) had endometrial atrophy. **CONCLUSION:** Postmenopausal bleeding should always be taken seriously and investigated meticulously no matter how minimal or insignificant it may appear. Early diagnosis makes successful treatment of endometrial hyperplasia and endometrial cancer in post-menopausal bleeding cases.

**KEYWORDS :** Postmenopausal, Bleeding per vagina

### INTRODUCTION

World Health Organization (WHO) defines menopause as cessation of menstruation permanently for a period of more than one year, which is resulted from loss of ovarian activity [1] Postmenopausal bleeding (PMB) is defined as bleeding from the genital tract, more than 12 months after the last menstrual period in a woman not on hormone replacement (HRT) [2] In our country a larger population, that is 71 million people are over 60 years of age. And postmenopausal women constituting to 43 million, according to the third consensus meeting of Indian Menopause Society (2008) [3]. Bleeding per vagina postmenopausally is one of the most common reason for referral to gynecological department, with a strong suspicion of malignancy various studies have proven that 90% of the postmenopausal bleed is due to endometrial carcinoma [4], whereas only 10-15% of the women on HRT and having bleed per vagina is diagnosed as endometrial carcinoma.

Endometrial cancer is the most common gynecological malignancy in the West, but in India, the incidence rates are low. 80 – 90% of the women have benign conditions like endometrial or cervical polyps, endometrial atrophy, infection, simple endometrial hyperplasia, medical disorders (e.g., liver cirrhosis), decubitus ulcer in cases of uterovaginal prolapse, neglected pessary and forgotten intra uterine device [5]. Serious conditions like endometrial cancer must be ruled out first. Women with risk factors like nulligravida, women with multiple sexual partners, obese, diabetic, taking exogenous estrogens/ tamoxifen, women who attain menopause late must be extensively investigated [6]. As the first line of investigation Transvaginal ultrasonography (TVS) is the recommended to assess the endometrial pathology, when the endometrial thickness is found to be more than 4 mm, it yields 98% sensitivity to detect endometrial cancer and pap smear for cervical pathology [7]. In suspected cases Dilatation and curettage or hysteroscopic guided biopsy are found to be the best modality to diagnose the aetiology for the bleed. Postmenopausal bleed until proved otherwise should be considered abnormal, except in those taking HRT [7]. Postmenopausal bleed is the most common symptom of endometrial carcinoma; hence every postmenopausal bleed should be investigated to rule out this first [8]. Hence, this study was planned at a government medical college in Bihar to explore the clinical significance of postmenopausal bleeding in term of its risk factors, incidence of malignancy and histopathological evaluation for the cause of postmenopausal bleed.

### METHODOLOGY

This descriptive study was carried out on 100 patients at Obstetrics and Gynecology department of Jawahar Lal Nehru Medical College& Hospital (JLNMCB) during July 2020 to June 2021. Postmenopausal women who presented clinically with complaint of vaginal bleeding, with their last menstrual period at least one year back and who were 45 years old or above were considered eligible for participation after taking informed consent. Patients having pre-mature menopause, surgical induced menopause, radiation induced menopause and chemotherapy induced menopause were excluded from the study. Detailed history was obtained from the patients including name, age, marital status, parity and postal address. Details regarding vaginal bleeding were recorded.

These included the timing of onset, duration and amount of bleeding. History of associated symptoms including presence of vaginal discharge, abdominal mass or pain and history of recent weight loss was obtained. Drug history especially, that of anticoagulants, hormones replacement therapy and tamoxifen therapy was also noted. Past medical and surgical history was checked especially regarding hypertension, diabetes mellitus and liver diseases. A thorough general physical examination was performed. Height and weight of the cases were measured and body mass index (BMI) calculated. Blood pressure was recorded. Specific clinical examination including abdominal, speculum and bimanual pelvic examinations were performed to assess the cervix and to determine size, position and mobility of the uterus. Cervical smears were taken. Transabdominal scan was done to assess endometrial thickness. All base line investigations including Full blood count, Random blood sugar, Urine routine examination, Coagulation profile, Xray chest and ECG were requested. Opinion regarding fitness for anesthesia was obtained from the anesthetist. After taking written informed consent Examination under anesthesia (EUA), cervical smear and dilatation and curettage (D & C) were performed.

Endometrial polyps, if found were avulsed. The specimens were collected in separate containers and sent for histopathological examination to pathology department. The data was entered into SPSS version 20 and results were obtained. The result has been expressed in terms of numbers and percentages.

### RESULTS

The mean age of women with postmenopausal bleeding (PMB) was 51.4 years. Age distribution of women with

postmenopausal bleeding is shown in Table 1. The parity of the study patients is shown in Table 2. The majority of the patients (45%) had parity of P2.

The histopathological findings of endometrium in patients with postmenopausal bleeding are A is bleeding in younger women. In the present series, 7 patients (14%) had malignancy, including endometrial carcinoma 4% and cervical carcinoma 10%, 27 patients (54%) had benign pathology, 9 had hyperplasia (18%) and 7 patients (14%) had endometrial atrophy.

**Table 1: Distribution of study subjects based of their age (N = 100)**

Age group (in years)	Number
40 - 44	41
45 - 49	29
50 - 54	20
55 - 60	6

**Table 2: Distribution of study subjects based of their parity (N = 100)**

Parity	Number
Para 1	8
Para 2	45
Para 3	27
Para 4	15
Para 5 or more	5

**Table 3: Distribution of study subjects based of their histopathological finding (N = 100)**

Histopathological finding	Number
Proliferative endometrium	16
Atrophic endometrium	14
Secretory endometrium	10
Complex hyperplasia with atypia	8
Squamous cell carcinoma of endometrium	10
Simple hyperplasia of endometrium	10
Endometrial carcinoma	4
Endo-cervicitis	14
Endometrial polyp	4
Cervical polyp	4

## DISCUSSION

Postmenopausal bleeding is an alarming sign that has a high possible of association with the cervical or uterine malignancy. It is one of the commonest symptoms the patient presents with, and hence should be worked up on priority bases to detect abnormalities if any present. The total number of cases we studied were 50, were in the age group of 40 to 44 years had the highest prevalence and the least was noted between the age greater than 70, whereas the study conducted by Wong SF et al, Sousa R et al, Bharani B et al, and Sheikh M et al it varied to 38-94, 43-82, 52-65, 42-84 years respectively [9, 10, 11, 12]. The mean age of the patient in the study we conducted was 51.4 years whereas it was much lower in the study conducted in the western population. The correlation between the association of age and PMB was reported by Gredmark T et al, which was not found in our study [13]. Women when presented with post-menopausal study were enrolled in the study and the endometrial curettage and cervical biopsies were performed. Benign conditions were noted in 62% of the cases which predominantly included atrophic endometrium, proliferative endometrium, secretory endometrium, benign endometrial polyp, Endo-cervicitis and simple cystic hyperplasia cases. Premalignant conditions like hyperplasia with atypia seen in 8%. Malignancy was noted in 14 women, out of whom 4% had endometrial and 10% had cervical cancer. Gredmark T et al, Lee WH et al, Dangal G et al and by Kaur Met al reported atrophic endometrium in 49.9%, 52.1%, 64.4% and 53% respectively, which the commonest cause for the postmenopausal bleeding where as in our study

the reported prevalence was only 14% to be atrophy [13, 23, 14, 15]. Meyer et al postulated the reason for the postmenopausal bleed in senile endometrium to be sclerotic degeneration of endometrial vessels whereas Hourihan et al stated that the anatomical vascular variations or local abnormal haemostatic mechanisms in the uterus to be the cause. Simple endometrial hyperplasia which is one of the most important predisposing factors for the development of endometrial cancer were tabulated as 10% in our study which ranged from 13.46% to 26.6% in various studies [17, 18]. The highest risk is noted in atypical endometrial hyperplasia which was observed in 8% of the study population, ranged from 1.8 - 8% in other studies [8, 10, 12]. Proliferative endometrium was observed in 16% in our study which is comparison to the study conducted by Phillip H et al found 20% [16]. Choo Y C et al well wrote that as a result of the conversion of adrenal androstenedione by peripheral fatty tissue to estrogen which can lead to proliferative endometrium, which due to the fluctuating low level of estrogen can bleed [20]. Endometrial adenocarcinoma which is the most threatened cause of postmenopausal bleeding, was found in 4% of the study population in our study, while various others found it to range from 6% to 12% [11, 13, 19]. Squamous cell carcinoma of the cervix was responsible for 10% cases of PMB in our study making it almost double that of the endometrial cancer, while it is reported to range from 8.8% to 39.6% in other studies [21]. The results support the fact that the diagnostic focus in post-menopausal bleeding in our country, should also focus on excluding cervical pathology. Cervical cancer is a preventable cancer because it has a long preinvasive state and also the preinvasive stage can be detected by cervical cytology screening program, and because the treatment for preinvasive lesions is effective [22].

## CONCLUSION

Postmenopausal bleeding should always be taken seriously and investigated meticulously no matter how minimal or insignificant it may appear. Early diagnosis makes successful treatment of endometrial hyperplasia and endometrial cancer in post-menopausal bleeding cases. Endometrial sampling is mandatory prior to therapeutic gesture. Cervical malignancies must be ruled out in all cases of PMB. Priority should be given to introduce screening methods for cervical cancer such as cervicography, gynecoscopy and pap smear at different levels of health care to effectively reduce the prevalence of cervical cancer among elder women.

## REFERENCES

- WHO. Research on the menopause in 1990s. Report of a WHO Scientific Group. World Health Organization technical report series. 1996; 866:1-107.
- Brand AH. The woman with postmenopausal bleeding. Australian Family Physician. 2007; 36:116-20.
- Jillani K, Khoro RB, Maqsood S, Siddiqui MA. Prevalence of malignant disorders in 50 cases of postmenopausal bleeding. J Pak Med Assoc. 2010; 60:540-43.
- Seebacher V, Schmid M, Polterauer S, Frischmuth KH, Leipold H, et al. The presence of post-menopausal bleeding as prognostic parameter in patients with endometrial cancer: a retrospective multinational study. BMC Cancer. 2009; 9:460-69.
- Wilailak S, Jirapinyo M, Theppisai U. Transvaginal Doppler Sonography: Is there a role of this modality in the evaluation of women with postmenopausal bleeding? Maturitas. 2005; 50:111-16.
- Panay N. Menopause and Postmenopausal women. Dewhurst' Obstetrics and Gynecology. 7th ed. UK. 2007; 479-93.
- Thijs I, Danesh HA, Bhal PS. The forgotten IUCD-a potential problem for postmenopausal women. J Obstet Gynaecol. 2002; 22:224-25.
- Yousaf S, Shaheen M, Rana T. Frequency of endometrial carcinoma in patients with postmenopausal bleeding. ANNALS. 2010; 16:290-94.
- Wong SF, Luk KL, Wong AY, Tang LC. Findings in women with postmenopausal bleeding investigated with hysteroscopy. J Obstet Gynaecol. 2001; 21:392-95.
- Sousa R, Silvestre M, Almeida e Sousa L, Falcão F, Dias I, Silva T, De Oliveira C, Oliveira HM. Transvaginal ultrasonography and hysteroscopy in postmenopausal bleeding: a prospective study. Acta Obstet Gynecol Scand. 2001; 80:856-62.
- Bharani B, Phatak SR. Feasibility and yield of endometrial biopsy using suction curette device for evaluation of abnormal pre and postmenopausal bleeding. J Obstet Gynecol India. 2008; 58:322-26.
- Sheikh M, Sawhney, Khurana A, Al Yatama M. Alteration of sonographic texture of the endometrium in postmenopausal bleeding a guide to further management. Acta Obstet Gynecol Scand. 2000; 79:1006-10.

13. Gredmark T, Sonja Kvint, Guillaume Havel, Lars Ake Maltsson. Histopathological findings in women with postmenopausal bleeding. *British Journal of Obstetrics and Gynaecology*. 1995; 102:133-36.
14. Dangal G. A Study of Endometrium of Patients with Abnormal Uterine Bleeding at Chitwan Valley. *Kathmandu Univ Med J*. 2003; 1:110-2.
15. Kaur M, Singh R, Sharma M. Endovaginal Sonographic Evaluation of Postmenopausal Uterine Bleeding. *Journal of Clinical and Diagnostic Research*. 2010; 4:2175-82.
16. Phillip H, Dacosta VE, Fletcher HM, Kulkarni SK, Reid ME. Correlation between transvaginal ultrasound measured endometrial thickness and histopathological findings in Afro-Caribbean Jamaican women with postmenopausal bleeding. *Journal of Obstetrics and Gynaecology*. 2004; 24:568-72.
17. Veena S Naik, Jyoti D Rege, and Kusum D Jashnani. Pathology of Genital Tract in Postmenopausal Bleeding. *Bombay Hospital Journal – Original/ Research*. [www.bhj.org/journal/2005\\_4703\\_july/html/original\\_pathology250.htm](http://www.bhj.org/journal/2005_4703_july/html/original_pathology250.htm)-17k.
18. Ahmed JA. Clinico Pathological Evaluation of Postmenopausal Uterine bleeding in Mosul City. *Tikrit Medical Journal*. 2007; 13:73-78.
19. Reddy Rani P, Devi JK, Papa D, Jayanthi S. Transvaginal sonography and progesterone challenge test for identifying endometrial pathology in postmenopausal women. *J of Obs and Gyn of India*. 2002; 52:135-38.
20. Choo YC, Mak KC, Hsu C. et al. Postmenopausal uterine bleeding of non-organic cause. *Obstet Gynecol*. 1985; 66:225-28.
21. Jilani K, Khoro RB, Siddiqui SMA. Prevalance of malignant disorders in 50 cases of postmenopausal bleeding. *JPMIA*. 2010; 60:540-43.
22. Sherris TD, Waeles ES, Tsu VD, Bishop A. Cervical cancer in developing countries a situation analysis. Working paper Washington: The World Bank, 1993.
23. Lee WH, Tan KH, Lee YW. The aetiology of postmenopausal bleeding—a study of 163 consecutive cases in Singapore. *Singapore Medical Journal*. 1995; 36(2):164-168.