



## C-REACTIVE PROTEIN LEVEL IN DIAGNOSIS OF ACUTE APPENDICITIS

**Dr. Anil Kumar Shah**

M.B.B.S., M.S. (SURGERY), Principle Specialist, Government Hospital Nathdwara, Rajsamand.

**Dr. Virendra Mahatma\***

Professor Medicine, Govt. Medical College Sirohi. \*Corresponding Author

**ABSTRACT**

**Background-** To study the c-reactive protein in acute appendicitis.

**Methods-** The source of data was from pretested proforma which takes into account clinical history, general physical examination, relevant investigations, imaging modalities. All data were analyzed by Epi-info software.

**Results-** Mean CRP level in 0-6 mm appendix diameter patients was  $1.63 \pm 2.32$  mg/dl, in 6-8 mm diameter cases was  $2.89 \pm 3.67$  mg/dl, 8-10 mm cases CRP level was  $3.60 \pm 3.24$  mg/dl and in 10-12 mm diameter cases CRP level was  $3.61 \pm 2.28$  mg/dl.

**Conclusion-** Raised serum C-reactive protein reducing the rate of negative explorations.

**KEYWORDS :** CRP, Appendix, Pain abdomen

**INTRODUCTION**

C-reactive protein (CRP) together with other acute phase proteins increases in response to tissue injury. Many studies have shown the value of raised serum CRP quantitative measurement in improving the diagnosis of acute appendicitis.<sup>1-4</sup>

In this study we correlate the quantitative serum levels of CRP with the diameter of appendix in acute appendicitis. This study emphasizes the impact of normal rather than raised serum C-reactive protein in reducing the rate of negative explorations.

**MATERIALS AND METHODS****Inclusion Criterion**

The inclusion criteria were following:

All the patients who admitted with acute appendicitis and posted for surgery were included in the study.

**Exclusion Criterion**

The exclusion criteria were following:

1. Children below 12 years and elderly above 60 years were excluded as the CRP response is not optimal.
2. Patients who managed conservatively or individuals who had undergone appendectomy excluded from this study.
3. Patients with past history of jaundice, signs and symptoms of liver disease, chronic alcoholic and with other coexisting acute inflammatory conditions were excluded, as CRP is exclusively produced in liver and raised in acute inflammatory condition.
4. Females taking oral contraceptive pill or pregnant were excluded as CRP is elevated in these individuals.
5. Patients, not willing to participate in the study (who refused to give consent).

**Statistical Analysis:**

Data was analysed in terms of demographic, clinical features, blood tests -white blood cells, serum CRP levels and diameter of appendix in acute appendicitis as per ultrasonography reports preoperatively.

**RESULTS**

**Table 1. Association Between CRP Level And Appendix Diameter**

Appendix diameter	No. of patients	CRP level (mg/dl)
0-6 mm	18	$1.68 \pm 2.30$
6-8 mm	32	$2.92 \pm 3.61$
8-10 mm	34	$3.22 \pm 3.02$
10-12mm	16	$3.70 \pm 2.63$
p-value=0.001		

Mean CRP level in 0-6 mm appendix diameter patients was  $1.68 \pm 2.30$  mg/dl, in 6-8 mm diameter cases was  $2.92 \pm 3.61$  mg/dl, 8-10 mm cases CRP level was  $3.22 \pm 3.02$  mg/dl and in 10-12 mm diameter cases CRP level was  $3.70 \pm 2.63$  mg/dl.

**DISCUSSION**

In our study serum CRP level was elevated in 90.00% of the patients and this rate corresponds to study done at Wishaw General Hospital, Lanarkshire UK by Khan MN et al<sup>5</sup> 2004 where CRP levels were elevated in 85.7% of patients with acute appendicitis, also with study done by Vinoth Kumar et al<sup>6</sup> where 90% of patients had elevated CRP levels.

But in contrast to our study according to Mikaelson et al<sup>7</sup> the elevation of serum CRP levels were found in only 47% and 72% of patients respectively.

Serum CRP levels increases after the onset of inflammatory pathology, so the levels might be normal until about 12hrs after the onset of symptoms of acute appendicitis so the time interval between the appearance of symptoms and the actual testing of serum CRP levels had some bearing on the result shown by Mickaelson et al in their studies.

According to our study the sensitivity and specificity of serum levels in diagnosis of acute appendicitis is comparable to the results given by other researchers. So it is derived from different studies that serum CRP test is highly sensitive and specific in making diagnosis of patients who truly had acute appendicitis.

**CONCLUSION**

No doubt surgeon's clinical diagnosis using time tested clinical signs is effective in diagnosing acute appendicitis. However elevated serum CRP levels and appendix diameter support the surgeon's diagnosis and hence avoids chances of error in diagnosis, due to atypical presentations and may avoid negative appendectomy.

**REFERENCES**

1. Hoffmann J, Rasmussen O. Aids in the diagnosis of acute appendicitis. Br J surg 1989; 76:774-779
2. John H, Neff U, Kelemen M. Appendicitis diagnosis today: clinical and ultrasonic deductions. World J surg 1993; 17:243-249
3. Jones PF. Suspected acute appendicitis: trends in management over 30 years. Br J surg 2001; 88:1570-1577
4. Lee SL, Walsh AJ, Ho HS. Computed tomography and ultrasonography do not improve and may delay the diagnosis and treatment of acute appendicitis. Arch surg 2001; 136:556-561
5. Khan MN et al. Study of role of WBCs CRP in diagnosis of acute appendicitis. 2004.
6. Vinoth Kumar. Study of diagnostic value of CRP in suspected acute appendicitis-prospective study. 2010.

7. Mikaelsson C, Arnbjornsson E. The value of C-reactive protein (CRP) determinations in patients with suspected acute appendicitis. *Ann Chir Gynaecol.*1984;73(5):281-4.
8. Shozoyokoyama. Large prospective study of CRP estimation in acute appendicitis done at Wakayama Medical Hospital.2007.
9. Asfar S, Safar H, Khoursheed M, Dashti H, al-Bader A. Would measurement of C-reactive protein reduce the rate of negative exploration for acute appendicitis? *JR Coil Surg Edinb.* 2000 Feb;45(1):21-4.