



STUDY OF EARLY INITIATION OF BREASTFEEDING AND IT'S OUTCOME ON THIRD STAGE OF LABOUR.

Dr. Rita. D

PROFESSOR & HOD, Department Of OBG, Navodaya Medical College Hospital & research Centre, Raichur.

Dr. Monika Devi Thokchom*

Junior Resident, Navodaya Medical College Hospital & Research Centre, Raichur. *Corresponding Author

ABSTRACT

Background: Nipple stimulation or breastfeeding are stimuli that can lead to the secretion of oxytocin and consequent uterine contractions and thus reduces bleeding during the third stage of labour.

Oxytocin and prostaglandin are hormones responsible for uterine contraction during the third stage of labour. Receptors in the uterine muscles are stimulated by exogenous or endogenous oxytocin leading to uterine contractions.

Aims And Objectives: To initiate breastfeeding of the newborn as soon as possible and to evaluate the effectiveness of early initiation of breast feeding on outcome third stage of labour in the experimental group .

Method: In this study, normal deliveries were observed in Labour room, Navodaya Medical College. Study participants were selected from delivery cases attended during the study period from march 2020 to march 2021 excluding caesarean section deliveries. Out of 100 patients 50 is for experimental and 50 for control groups, time of initiation of breastfeeding and its association was measured with spontaneous breathing, skin-to-skin contact and postnatal contact of mother and newborn in the third stage of labour.

Result: The result showed in cases 32%, 48%, 18%, 2% between 16-20, 21-25, 26-30, 31-35 years and in control 16%, 56%, 26%, 2% in mothers were in the age group of 16-20, 21-25, 26-30, 31-35 years respectively. The mean duration of third stage and amount of blood loss was 7.30 +/-1.31, 9.50 +/-2.58 and 197.60 +/-60.93, 298.98 +/-71.77 with the mean difference of 2.20 minutes and 101.38ml in the experimental and control group respectively.

Conclusion: There is positive effect of early initiation of breast feeding on duration of third stage and the amount of blood loss in the experimental group.

KEYWORDS : Early initiation; breast feeding; intra partum and post-natal .

INTRODUCTION

Breast feeding is the first fundamental right of the child. The initiation of breast feeding and the timely introduction of adequate safe and appropriate complementary foods in conjunction with continued breast feeding are of prime importance for the growth, development, health and nutrition of infants and children everywhere^[1,2]. It is found that early suckling and hand touching by babies stimulates oxytocin release which is significant for uterine contraction and early separation of placenta^[3,4].

In India child death accounts for two-third of mortality and half of the pediatric deaths occurring in infancy. Each year 26 million infants are born in India, around 10% of them do not survive to 5 years of age. India contributes to 25% of the 10 million under five deaths occurring worldwide every year. One of the main causes for this is delay in initiation of breast feeding^[5]. Only 24.5% women initiate breast feeding within one hour of delivery, exclusive breast feeding rate at 6 months is only 40%. Exclusive breast feeding has been identified as the single most effective, preventive intervention which could prevent 13 percent of all childhood deaths^[6]. WHO estimates that 1.5 million infant lives could be saved each year through increased breast feedings.

One -fifth of neonatal deaths could be prevented by early initiation of exclusive breast feeding. Breast feeding within the first hour^[7]. A descriptive study was conducted to assess the prevailing breast feeding and infant feeding practices among 1050 infants from 0 - 24 months of age in rural areas of central Karnataka. The results revealed that only 3(0.3%) infants were offered breast feeding within one hour after delivery. Delayed initiation of breast feeding was common; 35% of babies were not breast fed even at 48 hours of birth^[8,9]. Breastfeeding babies within the first hour of birth stimulate the production of breast milk. The first milk, also called as colostrums is extremely nutritious and helps to prevent diseases. Early initiation ensures that a newborn receives colostrum. Colostrum is often considered the baby's first immunization

because of its high level of vitamin A, antibodies, and other protective factors^[10].

Interestingly, the trend of early breastfeeding initiation, identified within the last decade from similar surveys, shows that mothers are less likely to practice so if they give birth in health facility than at home^[11,12]. This persistent lower coverage of early breastfeeding initiation within health facility is of concern since promoting institutional delivery (a proxy for skilled attendance) is a priority intervention to reach the targets of Sustainable Development Goals of reducing maternal and child mortality by 2030^[13].

OBJECTIVES

1. To initiate breastfeeding of the newborn as soon as possible in the experimental group.
2. To evaluate the effectiveness of early initiation of breastfeeding on outcome of third stage of labour in the experimental group in comparison with control group.

MATERIALS AND METHODS

Source Of Data: The study was carried out among selected 100 delivering patients in labour room of Obstetrics and Gynaecology of Navodaya Medical College, Raichur.

Study Design: prospective observational study with study period from March 2020-March 2021.

Sample Size: 100 patients in labour were selected excluding cesarean section deliveries after obtaining ethical committee clearance. 50 Patients were for cases (group A) and remaining 50 patients were for control (group B).

Inclusion Criteria

1. Mothers who are willing to participate in this study.
2. Mothers undergoing normal delivery between 37-40 weeks of gestation.
3. Singleton pregnancy
4. Live fetus

5. Neonates with APGAR score more than 7.

Exclusion Criteria

1. Mothers with severe pregnancy induced hypertension
2. Newborns of mother with gestational diabetes mellitus.
3. Mother with placenta previa.
4. Fetus with meconium stained amniotic fluid.

Methodology:

The prospective observational study was carried out among selected 100 delivering patients in labour room of Obstetrics and Gynaecology of Navodaya Medical College, Raichur excluding cesarean section deliveries. 50 Patients were selected for cases(group A) and remaining 50 patients were for control(group B). Time of initiation of breastfeeding was observed and its association was measured with spontaneous breathing, skin to skin contact and postnatal contact of mother and newborn in the third stage of labour .

RESULTS

In relation to the demographic and obstetrical variables it showed that mother in cases 32%, 48%,18%, 2% in the age group of 16-20 21-25, 26-30, 31-35 years and in control 16%, 56%, 26%, 2% were in the age group of 16-20 21-25, 26-30, 31-35 years respectively.The mean duration of third stage and amount of blood loss was 7.30+/-1.31, 9.50+/-2.58 and 197.60+/-60.93, 298.98+/-71.77 with the mean difference of 2.20 minutes and 101.38ml in the experimental and control group respectively. The calculated 't' value is -5.569 and -7.582 (p<0.000) was statistically significant in the duration of third stage of labour and amount of blood loss in the experimental group.

Table 1:Age Of The Patient

Age (years)	Group A (CASES)		Group B (CONTROL)	
	No. of patient	percentage	No. Of patient	percentage
16-20	16	32%	8	16%
21-25	24	48%	28	56%
26-30	9	18%	13	26%
30-35	1	2%	1	2%

The mean age of patient in labour in Group A was highest in 20-25 years with 48% and lowest with 30-35 years age group with 2%. While in Group B highest in 20-25 years with 56% and lowest with 30-35 years age group with 2%.

Table 2: Parity

Parity		Group A		Group B		Total	Chi-square value	P value
		No. of cases	Perce ntage	No. of cases	Perce ntage			
Primigravida	19	38%	21	42%	40	0.167	0.683	
	31	62%	29	58%				
Multigravida	31	62%	29	58%	60			
Total	50	100%	50	100%	100			

P=0.167>0.05,NS

The no. of patients in multigravida is 62% and 38% in primigravida in experimental group(A).The no. of patients in multigravida is 58% and primigravida is 42% in control group(B) respectively.

The no. of patients belonging to 37-40 weeks with 70% and >40 weeks with 30% in experimental group (A). The no. of patients belonging to 37-40 weeks with 82% and >40 weeks with 18% in Control group (B)

The no. of patients with previous breastfeeding is 30 with 60% and no previous feeling is 20 with 40% in experimental

group(A). The no. of patients with previous breastfeeding is 29 with 59% and no previous feeling is 21 with 42% in control group(B).

Table 3: Gestational Age

		Group A (CASES)		Group B		Total	Chi-square value	P-value
		No. of cases	Perce ntage	No. of cases	Perce ntage			
Gestational Age(weeks)	37-40 weeks	35	70%	41	82%	76	1.974	0.16
	>40 weeks	15	30%	9	18%	24		
Total		50	100%	50	100%	100		

P=0.16>0.05,NS

Table 4: Previous Breast Feeding

		Group A (CASES)		Group B		Total	Chi Square value	P value
		No. of cases	Perce ntage	No. of cases	Perce ntage			
Previous Breastfeeding	Yes	30	60%	29	58%	59	0.041	0.839
	No	20	40%	21	42%	41		
Total		50	100%	50	100%	100		


P=0.839>0.05 NS

Table :5 Gender Of The Baby

		Group A (CASES)		Group B		Total	Chi square test	P value
		No. of patient	Perce ntage	No. of patient	Perce ntage			
Sex	Female	22	44%	27	54%	49	1.000	0.317
	Male	28	56%	23	46%			
Total		50	100%	50	100%	100		


P=0.317>0.05 NS

The no. of female baby is 22 with 44% and male baby is 28 with 56% in experimental group (A). The no. of female baby is 27 with 54% and male baby is 23 with 46% in control group (B).



P=0.001<0.05,Significant

Graph 1: Duration Of Third Stage Of Labour(HRS)



P=0.001<0.05,Significant

Graph 2: Amount Of Blood Loss

The mean duration of third stage of labour in mins in experimental group(A) is 7.30 mins and in control group(B) is 9.58 mins with difference of 2.20 mins with is statistically significant.

The mean amount of blood loss in experimental group(A) is 197.60ml while in control group(B) is 298.98 ml with difference of 101.38 ml which is statistically significant.

DISCUSSION

The present study result revealed that there is a significant difference in the maternal outcomes among the parturients between case and control groups. The component duration of third stage of labour in case is case (group A) 7.30 ± 1.31 and control (group B) 9.50 ± 2.50 respectively and amount of blood loss 197.60 ± 60.93 in cases and 298.98 ± 71.72 in control groups were consistent with the findings Sumithra et al and A Prishanthini et al.

The present study observes that more number of participation is there from multigravida which is 62% and primigravida with 38% which is comparable to the study by Farhana et al with 50.6% in primigravida group and multigravida with 48.2%. It is also comparable to the study by Mahabat et al there from multigravida which is 52.7% and primigravida with 47.3% respectively.

Table 8: Present Study Vs Other Studies

	Present Study	Sumithra et al	A Prishanthini et al
Duration of third stage of labour(min)			
Case group	7.30 ± 1.31	9 ± 1	3.57 ± 1.52
Control group	9.50 ± 2.50	12 ± 1	10.27 ± 2.92
Amount of bleeding(ml)			
Case group	197.60 ± 60.93	125 ± 25	198.33 ± 25.74
Control group	298.98 ± 71.72	272.60 ± 60.2	302.67 ± 33.41

CONCLUSION

The study shows that early initiation of breastfeeding was effective on the duration of third stage and amount of blood loss among mothers in the third stage of labour. And there was positive correlation between the amount of blood loss and the duration of third stage of labour. All the patients in Group A has decreased amount of blood loss thereby preventing Post partum haemorrhage, the duration of third stage of labour, manual removal of placenta, uterine inversion, infections and puerperal sepsis.

In conclusion the early initiation of breastfeeding could be promoted by health care professionals in day to day caring the mothers in hospital.

REFERENCES

1. Renu Misra. "Practical Obstetrical Problems". New Delhi: B.I. Publications (2007).
2. Sudip Chakravarty. "Manual of Obstetrics". London: Elsevier Publications (2003).
3. Pandit Bhide V. "Textbook of Obstetrics". New Delhi: B.I. Publication Private limited (2006).
4. Pillitteri Adele. "Maternal and Child Health Nursing". Philadelphia: Lippincott (2007).
5. WHO. "The Role of Breastfeeding on the Prevention of infant mortality". Geneva. WHO (2010).
6. World Health organization. "Maternal and Newborn Health safe Motherhood; Essential Newborn care". Report of a Technical Working Group 72.4 (2006): 32-34.
7. Erickson's. Effect of early initiation of breastfeeding (2000).
8. Jane F Thompson. "Initial management of breastfeeding" (2001).
9. Marcelo Febo., et al. "Early initiation of Breast feeding and its benefit in Rural Egypt". International Breast Feeding Journal 104.1 (2007): 3-4.
10. "Effect of breastfeeding on infant and child mortality due to Infectious diseases in less developed countries". Lancet 355.9202 (2000): 451-455.
11. National Institute of Population Research and Training (NIPORT), Mitra and associates, ICF International Bangladesh Demographic and Health Survey 2014. Dhaka, Bangladesh and Calverton, Maryland, USA: NIPORT, Mitra and Associates, and ICF International; 2015.
12. National Institute of Population Research and Training (NIPORT), Mitra and associates, Macro International. Bangladesh Demographic and Health Survey 2007. Dhaka, Bangladesh and Calverton, Maryland, USA:

13. NIPORT, Mitra and Associates Macro International; 2009
Campbell OM, Graham WJ, group LMSSs. Strategies for reducing maternal mortality: getting on with what works. Lancet 2006;368(9543):1284-99.
[http://doi.org/10.1016/S0140-7636\(06\)69381-1](http://doi.org/10.1016/S0140-7636(06)69381-1) PMID:17027735.