Original Research Paper



Dr. Shaik Ahmad Sheriff	Associate Professor of General Surgery, GGH, Kadapa.
Dr. P. Sabitha	Associate Professor of General Surgery, ACSR, Nellore.
Dr. Syed Imam Sadiq	Assistant Professor of General Surgery, GGH, Kadapa.
Dr. K. Lokesh	Assistant Professor of General Surgery, GGH, Kadapa.
Dr. K. Devika	Post Graduate in General Surgery, SVMC, Tirupathi.
	ion. The modern approach to broast careinong management is multidissiplinger, including

ABSTRACT Introduction: The modern approach to breast carcinoma management is multidisciplinary, including surgery, radiotherapy, hormonal therapy, and chemotherapy. However, surgical management is the hallmark of the treatment of breast carcinoma. "Modified radical mastectomy" is the most commonly employed surgery for breast carcinoma.2 In Some cases, complications will occur after surgery. Complications contribute to morbidity, prolong the hospital stay, cost, and interfere with the management of breast carcer. e-operative evaluation, meticulous techniques, hemostasis, and wound closure can decrease complications. Aims & Objectives: To study the postoperative complications within 30 days of modified radical mastectomy, To identify the triggering factors of early complications, and to know the preventive and treatment modalities for complications. Patients And Methods: Prospective observational study on 100 Patients diagnosed with breast carcinoma who undergo Modified radical mastectomy in the Department of General Surgery at a tertiary care hospital over a period of one year. Conclusion: Seroma formation (18%), surgical site infection were common complications. The patient's age, higher BMI, comorbidities like diabetes and hypertension and electrocautery dissection, and early removal of drain were found as risk factors for seroma formation.

KEYWORDS:

INTRODUCTION

Over 2.3 million females were diagnosed with breast carcinoma and 685 000 deaths globally in 2020. Woman dies from breast cancer, mainly because of widespread metastasis.³

The modern approach to breast carcinoma management is multidisciplinary, including surgery, radiotherapy, hormonal therapy, and chemotherapy.¹ However, surgical management is the hallmark of the treatment of breast carcinoma. "Modified radical mastectomy" is the most commonly employed surgery for breast carcinoma.²

Early surgery complications are defined here as complications occurring within 1 month after surgery, include wound infection, seromas, altered sensation along the anterior axillary fold, wound dehiscence, flap necrosis, and hematomas. Based on different surveys, surgical site complications after breast surgeries are 0.8 to 26% wide-ranging incidence.⁴⁵

Seroma formation is the most frequent postoperative complication with an incidence of 3% to 85%.⁶ Infection developing within seroma increases morbidity and often results in the need for readmission, reimaging, drainage, and antibiotic usage.⁷

Incidence rates of Postoperative wound infections vary and range from 3% to 19%. The incidence of Flap necrosis is between 3% and 32%.^e Incidence of functionally significant lymphedema is <10%.

Thorough pre-operative evaluation, meticulous techniques, hemostasis, and wound closure can decrease complications.

RESULTS Table 1: Distribution Of Cases By Age

Table 1: Distribution Of Cases By Age							
Āge	Frequency	Percentage					
31 – 40	8	8%					

41 – 50	35	35%
51 – 60	41	41%
61 – 70	16	16%
Total	100	100%
Mean ± SD	50.31 ± 7.91	

In the present study 63% of the patients had BMI between the 18.6 - 24.9kg/m² and 28% of the patients had 25-29.9kg/m² and 7% of the patients ≥ 30 kg/m², 2% of the patients less than 18.5 kg/m². The mean and standard deviation of the patient's BMI were 24.27kg/m² and 3.56, respectively.

In the present study, 20% patients were Diabetic, 8% hypertension, and 6% had both hypertension and diabetes. Mean and standard deviation of Hb value was 10.67gm/dl and 1.39, respectively. 49% patients were found to be right-sided tumor. The mean tumor size and standard deviations were 5.15cm and 2.17cm, respectively. In this study, 41% of the patients were treated by neoadjuvant chemotherapy. The mean duration of surgery and standard deviation were 124.85minutes and 19.29minutes, respectively. In this study,50% of the usage of electrocautery during surgery and 50% of scissor usage to dissect the tissue during surgery.

Table 2: Comparison Of Complications With Risk Factors On POD 4

		No	Sero	FN	WD	SSI	AS	Hema
			mα					tomα
Age(yrs)	<50	7	3	0	0	2	1	1
	>50	13	8	1	1	0	2	1
BMI(kg/m ²)	<25	8	1	1	1	1	3	1
	25 – 30	8	6	0	0	1	0	1
	>30	4	4	0	0	0	0	0
DM		5	3	0	1	0	0	1
HTN		3	3	0	0	0	0	0
DM + HTN		4	3	1	0	0	0	0
Duration of	<150	19	10	1	1	2	3	2
surgery(min)	>150	1	1	0	0	0	0	0

36 ★ GJRA - GLOBAL JOURNAL FOR RESEARCH ANALYSIS

VOLUME - 11, ISSUE - 08, AUGUST - 2022 • PRINT ISSN No. 2277 - 8160 • DOI : 10.36106/gjra

Hematoma:

Dissection	EC	15	13	0	1	1	0	0
device	SC	9	2	1	0	1	3	2
Total drain	<500	1	1	0	0	0	0	0
output(ml)	500 - 1000	18	10	1	1	2	2	2
	>1000	1	0	0	0	0	1	0
Drain removal	<10	3	1	0	0	0	2	0
day(days)	10-15	17	11	1	1	1	1	2
	>15	4	3	0	0	1	0	0
lymph nodes	Nil	0	0	0	0	0	0	0
removed	<15	9	5	1	1	0	1	1
	>15	15	10	0	0	2	2	1

Table 3: Comparison Of Complications With Riask Factors On POD - 10

		No	Sero	FN	WD	SSI	AS	Hema
			mα					toma
Age(yrs)	<50	8	4	2	0	2	0	0
	>50	14	3	0	2	9	0	0
BMI(kg/m ²)	<25	6	3	0	0	3	0	0
	25 – 30	11	3	2	2	4	0	0
	>30	5	1	0	0	4	0	0
DM	•	11	1	1	1	8	0	0
HTN		0	0	0	0	0	0	0
DM + HTN		3	0	1	1	1	0	0
Duration of	<150	16	7	2	2	5	0	0
surgery(min)	>150	6	0	0	0	6	0	0
Dissection	EC	9	2	2	1	4	0	0
device	SC	9	1	0	1	7	0	0
Total drain	<500	5	0	0	1	4	0	0
output(ml)	500 - 1000	16	7	2	1	6	0	0
	>1000	1	0	0	0	1	0	0
Drain removal	<10	3	2	1	0	0	0	0
day(days)	10-15	9	1	0	2	6	0	0
	>15	6	0	1	0	5	0	0
Lymph nodes	Nil	0	0	0	0	0	0	0
removed	<15	6	1	1	1	3	0	0
	>15	12	2	1	1	8	0	0

Table 4: Comparison Of Complications With Risk Factors On POD -28

		No	Sero	FN	WD	SSI	AS	Hema
			mα					toma
Age(yrs)	<50	2	0	0	1	0	1	0
	>50	1	0	0	1	0	0	0
BMI(kg/m ²)	<25	1	0	0	0	0	1	0
	25 – 30	1	0	0	1	0	0	0
	>30	1	0	0	1	0	0	0
DM		1	0	0	1	0	0	0
HTN		0	0	0	0	0	0	0
DM + HTN		0	0	0	0	0	0	0
Duration of	<150	2	0	0	1	0	1	0
surgery(min)	>150	1	0	0	1	0	0	0
Dissection	EC	1	-	-	1	-	-	
device	SC	2	-	-	1	-	1	
Total drain	<500	1	0	0	1	0	0	0
output (ml)	500 - 1000	1	0	0	0	0	1	0
	>1000	1	0	0	1	0	0	0
Drain removal	<10	1	0	0	1	0	0	0
day(days)	10-15	2	0	0	1	0	1	0
	>15	0	0	0	0	0	0	
Lymph nodes	Nil	0	0	0	0	0	0	0
removed	<15	3	0	0	1	0	0	0
	>15	0	0	0	1	0	1	0

DISCUSSION

Seroma Formation: In the present study, seroma was the most common complication found with an incidence of 18%; this was in the range of other studies like Altinyollar H et al⁹ with 15.5% and Bhatty I et al¹⁰ with 20%, and Dahri FJ et al¹¹ with the 33.33%. On the contrary, Kuroi K et al¹² quoted that existing evidence was inconclusive for seroma formation.

In the present study, hematoma developed in 2 patients(2%). In Obadiel Y A et al¹³ studies, hematoma was found in one case(2%). This is comparable with the Rajiv Sharma et al¹⁴ studies in which bleeding was found more with scissor dissection and post-operative development of a hematoma.

Altered Sensation:

In this study, altered sensation along the anterior axillary fold was found in 4 patients. Couceiro et al¹⁵ study explained why occurring in the younger group, as increased nerve sensitivity and lower threshold due to anxiety and more aggressive nature of breast carcinoma needs aggressive surgical treatment.

Surgical Site Infection (SSI):

This study found SSI in 2 patients on the 4th postoperative day and 11 patients on the 10th postoperative day. Surgical site infection was found in 13 cases (13%). In the study conducted by Chandrakar N et al¹⁶SSI were found in 10 cases (24.39%). Obadiel Y A et al¹³ studies were found in 14 patients (28%).

Flap Necrosis:

In this study, flap necrosis was found in 3 cases (3%), one case on the $4^{\mbox{\tiny th}}$ postoperative day and 2 cases on the $10^{\mbox{\tiny th}}$ postoperative day. Flap necrosis was observed with a higher BMI in this study. Robertson SA et al¹⁷ study found higher BMI was the risk factor.

CONCLUSION

1. Seroma (18%) was the most common early postoperative complication. It is due to multiple risk factors like increased age, higher BMI and comorbidities, early drain removal and excessive lymphatic dissection, and electrocautery dissection without ligation of lymphatic vessels.

2. Surgical site infection (13%) is the next common complication, and risk factors are older age group, diabetes, and prolonged drainage.

3. Wound dehiscence (5%) occurs in older people with diabetes and seroma formation with infection cases.

4. Flap necrosis (3%) can be seen in higher BMI, diabetes and hypertension, electrocautery dissection.

5. Hematoma and altered sensation along the anterior axillary fold complications are less than other complications.

REFERENCES

- Fujii T, Le Du F, Xiao L, Kogawa T, Barcenas CH, Alvarez RH, Valero V, Shen Y, Ueno NT. Effectiveness of an Adjuvant Chemotherapy Regimen for Early-Stage Breast Cancer: A Systematic Review and Network Meta-analysis. JAMA Oncol. 2015 Dec; 1(9):1311-8.
- 2 Henke, G., Knauer, M., Ribi, K. et al. Tailored axillary surgery with or without axillary lymph node dissection followed by radiotherapy in patients with clinically node-positive breast cancer (TAXIS): study protocol for a multicenter, randomized phase-III trial. Trials 19, 667 (2018).
- https://www.who.int/news-room/fact-sheets/detail/breast-cancer
- Prospero E, Cavicchi A, Bacelli S, Barbadoro P, Tantucci L, D'errico MM. 4. Surveillance for surgical site infection after hospital discharge: a surgical procedure-specific perspective. Infect Control Hospital Epidemiol. 2006 . Dec:27(12):1313-7.
- Ruvalcaba-Limón E, Robles-Vidal C, PoitevinChacón A, Chávez-MacGregor 5. M, GamboaVignolle C, et al. Complications after breast cancer surgery in patients treated with concomitant preoperative chemoradiation: a case-controanalysis. Breast Cancer Res Treatment. 2006 Jan 1;95(2):147-52.
- Agrawal A, Ayantunde AA, Cheung KL. Concepts of seroma formation and prevention in breast cancer surgery. ANZ J Surg. 2006 Dec; 76(12):1088-95.
- 7 Stanczyk, M., Grala, B., Zwierowicz, T. et al. Surgical resection for persistent seroma, following modified radical mastectomy. World J Surg Onc 5, 104 (2007)
- 8. Vitug A, Newman L. Complications in breast surgery. Surg Clin North Am. 2007 Apr;87(2):431-51.
- Altinyollar H, Kapucuaglu N, Pak I, Berberoglu U. Lymphatic mapping and sentinel lymphadenectomy in early stage breast carcinoma. J Experiment 9. Clin Cancer Res. 2000; 19(2):141-4.
- Bhatty I, Ibrahim M, Chaudhry ML. Complications after modified radical mastectomy in early breast cancer. Pak J Medi Sci. 2004;20(2):125-30. Dahri FJ, Awan MS, Qazi AR, Khaskheli NM, Soomro IA. Early wound
- 11. complications following modified radical mastectomy with axillary clearance. J Surg Pak (Int). 2011 Oct; 16(4).
- Kuroi K, et al. Evidence-Based Risk Factors for Seroma Formation in Breast Surgery. Jpn J Clin Oncol. 2006. 36 (4) 1 97–206 13
- Obadiel, Y.A., Al-Ba'adani, M.N. and Haidar, Q.H. (2020) Early Complications

VOLUME - 11, ISSUE - 08, AUGUST - 2022 • PRINT ISSN No. 2277 - 8160 • DOI : 10.36106/gjra

- Followin 11, 1950L '00, NOCOD' 2022 'Third ISO'NOC' 22', Following Modified Radical Mastectomy—Among Breast Cancer Patients Admitted to AL Gomhory Teaching Hospital, Sana'a, Yemen between Jan. 2019-Jan. 2020. Open Access Library Journal, 7, 1-1.
 14. Sharma R, Rajan N Singh, Singh R. Comparative Study of Complications Following Electrocautery Dissection vs Conventional Scalpel/Scissor Dissection in Modified Radical Mastectomy. Curr Trends Diagn Treat 2018;2(2):98-101.
 15. Couceiro TC, Valenca MM, Raposo MC, et al. Prevalence of post-mastectomy pain syndrome and associated risk factors: a cross-sectional cohort study.
- Concertor IC, Viterior MM, Rapos MC, et al. Prevalence of post-masteriority pain syndrome and associated risk factors: a cross-sectional cohort study. Pain Manag Nurs 2014;15:731-7.
 Chandrakar N, Shinde RK. Study the early complications of modified radical masteriory performed. Int Surg J 2019;6:239-43.
 Robertson SA, Jeevaratnam JA, Agrawal A, Cutress RI. Mastectomy skin flap
- necrosis: challenges and solutions. Breast Cancer (Dove Med Press). 2017;9:141-152. Published 2017 Mar 13.