



COMPARISON OF CAUDAL BUTORPHANOL AND KETAMINE AS ADDITIVES TO BUPIVACAINE IN PAEDIATRIC INFRAUMBILICAL SURGERY

Ritu Pauranik

Department of Anesthesia and Intensive care , MGM Medical college, Indore

Nandini Bhatt

Department of Anesthesia and Intensive care , MGM Medical college, Indore

ABSTRACT

Background: To compare the efficacy of caudal additives Butorphanol and Ketamine in pediatric age group undergoing infra umbilical surgery. **Methods:** 75 male children of ASA grade I, aged 4-10 years <25 kilograms, were randomised to one of the two groups. Group BK received 0.75 ml/kg (0.25%) bupivacaine with ketamine 0.5mg/kg and Group BB received 0.75ml/kg (0.25%) bupivacaine with butorphanol 25mcg/kg, in caudal epidural anesthesia. Hemodynamic variables and respiratory rate were monitored in all patients. Sedation score, pain score and requirement of rescue analgesia were recorded at preset time intervals along with post-operative complications. **Results:** Intra-operative hemodynamics, respiratory rate, SpO₂, EtCO₂, were maintained within normal limits in each group. Children were found to be more sedated in group BB till 2 hours post operatively as compared to group BK. There were no post operative complications in any of the groups. Mean duration of analgesia was longest in Group BB (16 hrs) as compared to Group BK (12 hrs). Minimum number of rescue analgesics were required in the BB group (1.12) as compared to group BK(1.76) in 24 hrs postoperatively. **Conclusion:** The addition of 0.5mg/kg of ketamine and 25mcg/kg of butorphanol to bupivacaine in caudal epidural space results in superior analgesia with a longer period without an increase of side effects with longest duration of analgesia achieved with butorphanol.

KEYWORDS : Pediatric surgery; Caudal epidural anesthesia; Butorphanol; Ketamine:Bupivacaine

INTRODUCTION

Caudal anesthesia is one of the most popular regional blocks in children. This technique is useful adjunct during operative and post-operative period to provide analgesia. 1,2 Caudal blocks are used worldwide to provide safe and effective perioperative analgesia for paediatric patients undergoing urological, lower abdominal, and lower limb surgery. It has gained popularity as this regional technique is easy to learn, offers a high success rate, and has a low incidence of major adverse events, including dural puncture or intravascular injections. 3 Also caudal epidural is useful as it attenuates the stress response of surgery and anesthesia in sub-umbilical surgeries 1,2.

Plain bupivacaine being an agent of shorter duration, different caudal additives have been coadministered with bupivacaine to prolong the duration of caudal epidural analgesia 4,5,6. Ketamine is an anesthetic and analgesic agent with a wide range of applications in pediatric anesthesia. [5] It exerts its effects by binding non-competitively to a subset of glutamate receptors stimulated by the excitatory amine N-methyl D-aspartate (NMDA), blockade of which leads to a decrease in the activation of dorsal horn neurons. These receptors are located throughout the central nervous system as well as in the substantia gelatinosa in the spinal cord and play an important role in central pain processing and in neural plasticity in the spinal cord. [6]

In our study we compared ketamine and butorphanol used as additives to bupivacaine for caudal epidural analgesia in children undergoing sub-umbilical surgery.

MATERIAL AND METHODS

After approval from the hospital ethical committee, a study was conducted on 60 male children of ASA grade I, aged 4-10 years, <25 kilograms, scheduled for sub-umbilical elective surgery including inguinal and genito-urinary operations at department of anaesthesia & intensive care , Sri Aurobindo Institute of Medical Sciences, Indore. Patients with airway problems, tonsillitis, upper respiratory tract infections, huge hernial sacs, allergies to local anesthetics, spinal deformities, local infection and bleeding tendencies were excluded from the study. After appropriate fasting all patients were pre medicated with syrup promethazine (0.5mg/kg) body weight

two hours prior to surgery. All procedures were carried out under general anesthesia in supine position.

Patients were shifted to operation theatre and all monitors were attached including SpO₂, noninvasive B.P, ECG. Basal parameters (pulse rate, blood pressure, respiratory rate, SpO₂) were recorded and anesthesia was induced with O₂/N₂O (50:50%) with high concentration of Sevoflurane (3-4 MAC%) on spontaneous respiration (tidal volume breathing) using Ayre's T-piece circuit via face mask. Intravenous access made and Ringer's lactate as per fasting period, started simultaneously. Injection ranitidine 1mg/kg and injection ondansetron 0.1mg/kg was added to the pediatric set in all patients. All patients were randomly allocated in two groups using table of random numbers:

Group BK (n=30): 0.75ml/kg (0.25%) bupivacaine with ketamine 0.5mg/kg
Group BB (n=30): 0.75ml/kg (0.25%) bupivacaine with butorphanol 20mcg/kg

Caudal block was given in the left lateral position. Cases with bloody tap were excluded from the study. Patients were then turned supine and an LMA (proseal / classic) of appropriate size was placed and secured. No analgesia was given during the operation.

All monitoring (pulse rate, blood pressure, respiratory rate, SpO₂, EtCO₂) was carried out every 10 minutes throughout the surgery. Anaesthesia was maintained on O₂ / N₂O (33:66%) with sevo flurane (1-2) MAC% on assisted respiration using Ayre's T-piece circuit with fresh gas flow of 2-3 times the minute volume. After the surgery was over LMA was removed after proper oxygenation and suctioning in an awake patient.

Postoperatively the patients were scored for pain and sedation along with monitoring of vitals (pulse rate, blood pressure, respiratory rate) by a blinded observer who was not aware about the group to which the patient belong. A modified objective pain score, given a maximum score of 10 (table 1) was used to assess pain over a 5-minute period. Given below in the table 2 are listed the demographic data of the patients. Sedation scoring was done according to the four point sedation scores (0 = eyes open spontaneously, 1 = eyes open

on speech, 2 = eyes open when shaken, 3= unarousable). Rescue analgesia (with syrup paracetamol 20 mg/kg) was given only on demand or when the pain score was ≥ 4. Total doses of rescue analgesic were noted for 24 hours. Patients were evaluated for any postoperative complications –pruritis, nausea, vomiting, urinary retention, local hematomas and headache. Time of passing urine and ambulation (taken as independent sitting) was inquired and assessed at 8hrs postoperatively.

Descriptive statistics i.e. mean and standard deviation have been used for continuous variables like age, weight, duration of surgery, duration of anaesthesia, heart rate, blood pressure (systolic, diastolic and mean), duration of post operative analgesia, total number of doses of rescue analgesic intake in 24 hours. Statistical methods used are ANOVA, followed by POST HOC test for comparison of the continuous variables. Frequency distributions and their percentage were applied for the categorical variables (pain score, sedation score). CHI SQUARE test was applied for the comparison of categorical data. P value < 0.05 is taken as statistically significant.

RESULTS

Demographic variables (table 2) were comparable in the two groups, including age, weight, basal heart rate and blood pressure (systolic, diastolic and mean), duration of surgery and the duration of anaesthesia with p>0.05. There was a symmetrical decrease in heart rate with time from basal to post induction to post caudal and up to 40 minutes in each group (Fig. 1) (p value<0.05). Mean Arterial Pressure showed a fall after induction of anesthesia, but was stable after the administration of caudal block (Fig.2). Respiratory rate was maintained at 24-26 breaths per minute throughout the intra operative period and there was no respiratory depression, post operatively. Supplementary analgesia was not needed intra operatively in any of the groups after the effect of caudal block was established.

There was some sedation seen in group BB, with 32% children remaining sedated (sedation score= 2) till half hour post operatively. Only 8% in group BB were sedated at 2 hour postoperatively with sedation score= 1 (p value=0.00). Pain score ≥ 4 was found in only 4% children in group BB and 8% children in group BK at 12 hours post operatively. The pain scores were recorded at regular intervals post operatively as shown in Fig.3. The time to first rescue analgesic was 18.18 hours in group BB and 14.16 hours in group BK (P<0.05). Mean rescue analgesic requirement was 1.12 and 1.76 in groups BB and BK, respectively (P<0.05)

There were no post operative complications like nausea, vomiting, pruritis, urinary retention, respiratory depression, hypotension, headache, local hematoma and no motor blockade noted in any of the groups.

Mean time of passage of urine was comparable in both groups was 3.96hrs in group BB and 3.94 hrs in group BK (p value=0.88) and time to ambulation in groups BB and BK were 3.96hrs and 2.97hrs, respectively (p=0.001).

DISCUSSION

The present study results show that both ketamine(0.5mg/kg) and butorphanol(20mcg/kg) when added to bupivacaine (0.25%) in caudal epidural space prolong the duration of analgesia with a longer duration achieved with butorphanol.

Butorphanol is a totally synthetic compound of the nalorphine cyclazocine series. It is a mixed agonist – antagonist with intrinsic activity at receptors of the muopioid type (morphine like). It is also an agonist at kappa opioid receptors. Its interactions with these receptors in the CNS apparently mediate most of its pharmacological effects, including analgesia. Butorphanol has an analgesic action similar to

that of morphine, with less respiratory depression, less nausea and vomiting, no undesirable psychomimetic effects and the provision of perioperative amnesia.⁷

In a study conducted by Singh V, Knaujia A, et al in 2006 demonstrated that butorphanol 25 g/kg added to caudally administered local anesthetics significantly increased the duration of postoperative analgesia in patients undergoing elective infraumbilical surgery with a mean duration of analgesia being 14.5±3.5 hours group BS (butorphanol with bupivacaine), (P value<0.001) without any significant side effects⁸. Similarly Lawhorn CD, et al (1997) in their study found that significantly fewer patients in the butorphanol group required rescue morphine sulfate in the postanesthesia care unit (p ≤0.001) and were much less likely to develop pruritis or nausea and vomiting or to require supplemental O₂ to maintain SpO₂ above 90%⁹. Bailey AG (1994)¹⁰, William MS(1995)¹¹, and Lawhorn CD(1995)¹² all observed that addition of a narcotic agonist antagonist butorphanol in the epidural space virtually eliminated side effects of nausea, vomiting, pruritis, urinary retention or respiratory depression without causing undue sedation.

Ketamine is chemically related to phencyclidine and acts by binding non-competitively to N-Methyl D-Aspartate receptors in the CNS and the substantia gelatinosa in the spinal cord.

A. Schnabel et al (2011) conducted a meta-analysis of 13 RCTs (published between 1991 and 2008) and found that there was a significantly longer time to first analgesic requirements in children receiving a caudal regional anaesthesia with ketamine in addition to local anaesthetics which indicates that adding ketamine to caudal local anaesthetics provides a prolonged and improved postoperative analgesia with few adverse effects compared with local anaesthetics with no major neuropathological events in the early postoperative follow-up (6–8 weeks after surgery) of four trials.¹³

Caudal additives ketamine (0.5mg/kg) and butorphanol (20mcg/kg) when added to bupivacaine (0.25%) prolong the duration of block without any side effects. Children were found to be more sedated in group BB till 2 hours post operatively with longest duration of analgesia in Butorphanol group.

CONCLUSION

We can now conclude that both ketamine and butorphanol are useful additives to bupivacaine in caudal block to prolong the duration and improve the quality of analgesia in paediatric infra umbilical surgery without any undue side effects.. The haemodynamics are stable intra-operatively. There is no motor blockade and urinary retention with any of these additives. Duration of analgesia up to 20 hours has been achieved with butorphanol which is excellent for the pediatric population as no supplementation of analgesia is required for a long time and child can enjoy an active life.

Table 1. MODIFIED OBJECTIVE PAIN SCORE:

Variable	Score 0	Score 1	Score 2
Crying	None	Consolable	Not consolable
Movement	None	Restless	Thrashing
Agitation	Asleep/calm	Mild	Hysterical
Posture	Normal	Flexed	Holds injury site
Verbal	Asleep/No complaint	Complains but cannot localize	Complains and can localize

Table 2. PATIENT DEMOGRAPHICS:

Demographic Data	Group BB (n=30)	Group BK (n=30)	P Value
Age(in years): range and Mean ±SD	5-7 6.4±2.12	5-7 6.10±1.74	0.49
Weight(in kg): range and Mean ±SD	15-18 16.36±3.71	15-17 16.44±2.63	0.42

Heart Rate(beats/min): range and Mean ±SD	107-114 110.56±8.66	107-112 109.56±6.39	0.07
Systolic BP(mm of Hg): range and Mean ±SD	106-111 108.52±5.42	103-108 105.20±5.57	0.77
MAP(mm of Hg): range and Mean ±SD	74-80 76.73-7.36	74-81 77.60±7.94	0.27
Diastolic BP(mm of Hg): range and Mean ±SD	57-65 60.84±9.59	60-68 63.80±10.24	0.89
Duration of surgery (in minutes): range and Mean ±SD	35-49 42.00±16.58	35-43 39.20±9.54	0.73
Duration of anaesthesia (in minutes): range and Mean ±SD	55-69 62.00±16.58	55-63 59.20±9.54	0.73

TABLE 3. Sedation Score, Mean Pain score, Duration of Analgesia, Total no. of rescue analgesics, time to passage of urine, time to ambulation in the Postoperative Room in Different Groups

Group	Group BB Mean±S.D.	Group BK Mean±S.D.	P value
Sedation score	0.30±0.32	0.04±0.58	0.001
Mean Pain score	0.44±0.63	0.55±0.58	0.001
Duration of analgesia(hrs)	18.85±2.05	14.16±1.65	<0.05
Total no. of rescue analgesics	1.12±0.33	1.76±0.52	<0.05
Time to passage of urine(hrs)	3.94±0.21	3.96±0.17	0.88
Time to ambulation(hrs)	3.96±0.22	2.97± 0.16	0.001




Fig 1. Intraoperative variation in Heart rate in the two groups..




Fig 2. Intraoperative variation in Mean Arterial Pressure in the two groups.




Fig.3 Mean Pain Score in the Post operative period

REFERENCES

1. Khalil SN, Hanna E, Farag A, Govindaraj R, Vije H, Kee S, Chuang AZ.. Pre surgical caudal block attenuates stress response in children. Middle East J Anesthesiol 2005; June 18(2): 391-400.
2. Silvani P, Comporesi A, Agostino MR, Salvo I. Caudal anaesthesia in paediatrics: an update. Minerva Anesthesiol 2006; Jun 72(6): 453-9.
3. A. Schnabel, D. M. Poepping, Kranke P, Zahn PK, Pogatzki-Zahn EM. Efficacy and adverse effects of ketamine as an additive for paediatric caudal anaesthesia: a quantitative systematic review of randomized controlled trials. British Journal of Anaesthesia 16 August 2011; 107 (4): 601-11.
4. Cook B, Grubb DJ, Albridge LA, Doyle E. Comparison of the effects of adrenaline, clonidine and ketamine on the duration of caudal analgesia produced by bupivacaine in children. British Journal of Anaesthesia 1995; 75(6): 698-701.
5. Kumar P, Rudra A, Pan AK, Acharya A. Caudal additives in Paed: A Comparison Among Midazolam, Ketamine and Neostigmine coadministered with Bupivacaine. Anaesth Analg 2005; 101: 69-73.
6. Reves JG, Peter SA. Intravenous Non-opioid Anaesthesia. In: Ronald D. Miller (ed), Anesthesia. 2005; vol (1) ch 10: 345-50.
7. Sung Y-F, Weinstein MS, Ghani GA. Balanced anesthesia : A comparison of butorphanol and morphine. Southern Medical Journal 1984 ; 77 : 180-182.
8. Singh V, Knaajia A, Singh GP. Efficacy of caudal Butorphanol. Indian J Paediatr 2006; 73(2): 147-150.
9. Lawhorn CD, Stoner JM, Schmitz ML, Brown RE Jr, Stewart FW, Volpe P, Shirey R. Caudal epidural butorphanol plus bupivacaine v/s bupivacaine in paediatric outpatient genitourinary procedures. J Clin Anaesth 1997; Mar 9(2): 103-8.
10. Bailey AG, Valley RD, Freid EB, Calhoun P. Epidural morphine combined with epidural or intravenous butorphanol for postoperative analgesia in pediatric patients. Anesth Analg 1994; Aug 79(2): 340-4
11. Splinter WM, O'Brien HV, Komocar L. Butorphanol: an opioid for day care paediatric surgery. Can J Anaesth 1995; 42(6): 483-6.
12. Lawhorn CD, Boop FA, Brown RE Jr, Andelman PD, Schmitz ML, Kymer PJ, Shirey R. Childs Nerv Syst. 1995 Nov;11(11):621-4. Continuous epidural morphine/ butorphanol infusion following selective dorsal rhizotomy in children. Childs Nerv syst 1995 ; Nov (11): 621-4.