



## OVARIAN TORSION IN PRE-PUBERTAL GIRL:A CASE REPORT

**Dr. Shivangi Marakana**

Resident Doctor At GMERS Medical College And General Hospital, Junagadh, Gujarat.

**Dr. Rajankumar Padasala**

Assistant Professor At P.D.U. Medical College,rajkot,gujarat.

**ABSTRACT**

**Background:** Ovarian torsion(adnexaltorsion) is a rare event in pediatric patients which is primarily managed by pediatric general surgeons. **Case Presentation:** This study presents a case of ovarian torsion in a 8 yr old girl with a history of episodic lower abdominal pain,nausea,and vomiting for 2 days.Her physical examination was normal except for mild tenderness in the lower abdomen with no palpable mass. A color Doppler ultrasound was performed for further investigation, and an ovarian torsion was reported without sonographic sign so intussusceptions and acute appendicitis, so she underwent laparotomy. A relatively complete torsion was observed in the right ovarian pedicle. Detorsion of ovary was done and it was decided to preserve the ovary.The patient was discharged and 1 months later follow-up colorDoppler ultrasound showed normal ovarian blood flow. **Conclusion:** The possibility of ovarian torsion must be considered in all female infants with suspicious abdominal pain.

**KEYWORDS :** Ovarian torsion, Child Laparotomy.

**INTRODUCTION**


Fallopian tube and ovary's abnormal twisting around the ovarian pedicle called as ovarian torsion (1). In most cases, ovarian torsion occurs during the reproductive years, while idiopathic torsion is a rare event with in the pediatric age group(2).In children, adnexal torsion can occur at different ages, but >52% of cases occur between the ages of 9 and 14, with an average of 11 yr. ovarian torsion is also rare in neonates. It occurs in16% of girls <1yr (3,4). The signs and symptoms of ovarian torsion, like lower abdominal pain, fever, nausea, and vomiting, are usually nonspecific. Tenderness in the lower abdomen and sometimes a mass may be revealed by clinical examination. Differential diagnosis includes intussusception, infantile colic pain, appendicitis, acute gastroenteritis, constipation, mesenteric lymphadenitis, ureteric colic, and urinary tract infection (5). Adnexal torsion is a clinical diagnosis. When the clinical features, patient's history, and/or imaging are suspected of adnexal torsion, immediate exploratory surgery makes the final diagnosis. In the pediatric population, the best diagnostic and therapeutic approach is laparoscopic surgery.The persistence of pain for more than 10 hr is linked to an increased rate of tissue necrosis. So, fast diagnosis and operation can prevent irreversible adnexal damage, salvaging the adnexal torsion and increasing the success of ovarian preservation(6-9). This clinical case report presented a 8-yr-old girl with right ovarian torsion as a rare case.

**Case Presentation**


The case is a 8 yr old girl complaining of episodic lower abdominal pain, nausea, and vomiting for 1 days. Her parents visited emergency room at GMERS medical college and general hospital,Junagadh .They were advised for basic blood ,urine investigation with USG-abdomen+pelvis & further observation at hospital till reports available.However, since abdominal pain,nausea,and vomiting(6-8 times/day) continued from yesterday,she looked pale and lethargic with vital signs as follows: BP: 90/60, PR: 100, T: 36.5, RR:24, O<sub>2</sub> sat: 96%. Conservative treatment given.

Her physical examination was normal except for mild to moderate tenderness in the right lower abdomen with no palpable mass. Moreover, leukocytosis was noted in complete blood count. In abdominopelvic ultrasound was performed, demonstrating a prominent right ovary size of 46×39 mm with no definite vascularity that was highly suspicious for ovarian torsion (Figure 1). No sonographic sign of intussusception and

acute appendicitis were observed. Eventually, she underwent laparotomy due to ovarian torsion. A relatively complete torsion was observed in the right ovarian pedicle. Ovarian detorsion was done and decided to preserve the ovary even though the right ovary and fallopian tube had a dark appearance (Figure 2).Ovary,fallopian tube and round ligament fixation done to prevent re-torsion episode.The patient was discharged in stable condition.1 month after the surgery the color Doppler ultrasound showed normal ovarian blood flow.



**Figure 1:** Right Ovarian torsion in USG



**Figure 2:** Right ovarian torsion in Laparotomy.

**Ethical Considerations**

Ethical considerations were maintained during this study, and written consent was taken from the patient's parents for this presentation.

**DISCUSSION**

Ovarian torsion commonly occurs during childbearing age and postmenopausal age (5). It is estimated to be about 5 per 100,000 in females between 1 and 20 yr old (1).There are several risk factors associated with ovarian torsion, such as ovarian mass (especially dermoid) or cyst, previous torsion or surgery, polycystic ovary, tubal sterilization, and pregnancy (10, 11). In the pediatric patient, about 51-84% of adnexal

torsions occur due to adnexal pathology, which includes follicular or hemorrhagic ovarian cysts, cystic teratomas or dermoid, and less frequently cystadenomas, para ovariancysts and hydrosalpinx(8,12). Physical examination findings, such as normal temperature, low-grade fever, and/or mild tachycardia, are typically nonspecific to adnexal torsion. Acute pelvic or abdominal pain with variable duration (days to months) is the most common symptom, and it is often isolated to one side. Also, it can be mild or intense, non-radiating, constant, or intermittent (depending on whether the torsion is partial or complete). In addition, symptoms such as flank pain, nausea and vomiting due to peritoneal reflexes, and anorexia are usually associated with adnexal torsion. In patients with adnexal torsion, most laboratory findings are normal. However, a slight leukocytosis can sometimes occur (13). Lower abdominal pain, nausea, vomiting, mild tenderness in the lower abdomen with no palpable mass, and leukocytosis were diagnosed in this patient. Clinical diagnosis of ovarian torsion must be differentiated from hemorrhagic ovarian cyst rupture, appendicitis, gastroenteritis, and kidney stone (8, 13, 14). Moreover, adnexal torsion is often difficult to diagnose because of the vague and variable clinical signs and nonspecific imaging findings. To evaluate blood flow to the ovaries in pediatric and adolescent patients, pelvic ultrasonography with color Doppler is the common and accurate diagnostic imaging of adnexal torsion (14, 15). Therefore, ultrasonography is usually enough for diagnosis(5). Computed tomography and magnetic resonance imaging are not the first-line imaging modalities in adnexal torsion, but they can be used to assess the structure if ultrasound is not available or if the differential diagnosis remains with other cases such as appendicitis (14). Delay in diagnosis of ovarian torsion commonly leads to loss of the involve do vary and other complications (5). Ovarian torsion can be treated with detorsion or oophorectomy. Detorsion is a safe and effective treatment for ovarian torsion and it may be more optimal for all premenopausal women because of the high rates of ovarian preservation and low risk of complications (16). Recent evaluation of pediatric patients in the national in patient sample showed that 15%, 6% and 78% of cases underwent detorsion, Detorsion with oophoropexy, and oophorectomy alone, respectively (17). Another case report presented ovarian torsion and histopathological disorders, including cystic lesions with extensive calcification in a 4-month-old baby undergoing oophorectomy (5). Although ovarian torsion is rare in children, early diagnosis is critical because it can save the ovary and prevent psychological trauma to the parents and child. Therefore, the possibility of ovarian torsion must be considered in all female infants with suspicious abdominal pain.

### Conflict Of Interest

The authors have no conflict of interest relevant to this article.

### REFERENCES

1. Tsai J, Lai J-Y, Lin Y-H, Tsai M-H, Yeh P-J, Chen C-L, et al. Characteristics and risk factors for ischemic ovary torsion in children. *Children* 2022;9:206.
2. Tielli A, Scala A, Alison M, Vo Chieu VD, Farkas N, Titomanlio L, et al. Ovarian torsion: Diagnosis, surgery, and fertility preservation in the pediatric population. *Eur J Pediatr* 2022;181:1405–1411.
3. Sriram R, Zameer MM, Vinay C, Giridhar BS. Black ovary: Our experience with oophoropexy in all cases of pediatric ovarian torsion and review of relevant literature. *J Indian Assoc Pediatr Surg* 2022;27:558–560.
4. Sosnowska-Sienkiewicz P, Mankowski P. Profile of girls with adnexal torsion: Single center experience. *Indian Pediatr* 2022;59:293–295.
5. Karambelkar RP, Shah S, Deshpande N, Singh D. Ovarian torsion in a 4-month-old baby. *Pediatric Oncall* 2013; 10: 60.
6. Sheizaf B, Ohana E, Weintraub AY. "Habitual Adnexal Torsions"-recurrence after two oophoropexies in a prepubertal girl: A case report and review of the literature. *J Pediatr Adolesc Gynecol* 2013;26:e81–e84.
7. Yildiz A, Erginel B, Akin M, Karadağ CA, Sever N, Tanik C, et al. A retrospective review of the adnexal outcome after detorsion in premenarchal girls. *Afr J Paediatr Surg* 2014; 11:304–307.
8. Geimanaitė L, Trainavičius K. Ovarian torsion in children: Management and

outcomes. *J Pediatr Surg* 2013; 48: 1946–1953.

9. Casey RK, Damle LF, Gomez-Lobo V. Isolated fallopian tube torsion in pediatric and adolescent females: A retrospective review of 15 cases at a single institution. *J Pediatr Adolesc Gynecol* 2013;26:189–192.
10. Bridwell RE, Koyfman A, Long B. High risk and low prevalence diseases: Ovarian torsion. *Am J Emerg Med* 2022;56:145–150.
11. Gupta A, Gadipudi A, Nayak D. A five-year review of ovarian torsion cases: Lessons learnt. *J Obstet Gynaecol India* 2020;70:220–224.
12. Focseneanu MA, Omurtag K, Ratts VS, Merritt DF. The auto-amputated adnexa: Are view of findings in a pediatric population. *J Pediatr Adolesc Gynecol* 2013;26:305–313.
13. Sasaki KJ, Miller CE. Adnexal torsion: Review of the literature. *J Minim Invasive Gynecol* 2014;21:196–202.
14. Gounder S, Strudwick M. Multimodality imaging review for suspected ovarian torsion cases in children. *Radiography* 2021;27:236–242.
15. Naiditch JA, Barsness KA. The positive and negative predictive value of transabdominal color Doppler ultrasound for diagnosing ovarian torsion in pediatric patients. *J Pediatr Surg* 2013;48:1283–1287.
16. Ke K, Conrad DH, Cario GM. Conservative management of ovarian torsion. *Gynecol Obstet* 2020;10:543.
17. Sola R, Wormer BA, Walters AL, Heniford BT, Schulman AM. National trends in the surgical treatment of ovarian torsion in children: An analysis of 2041 pediatric patients utilizing the nationwide inpatient sample. *Am Surg* 2015; 81:844–848.