



A CROSS-SECTIONAL OBSERVATIONAL STUDY ON CLINICOPATHOLOGICAL FEATURES OF OBSTRUCTIVE JAUNDICE

Dr. Animesh Choudhary

Professor, Department of Medicine, Shri Balaji Institute of Medical Sciences, Raipur

ABSTRACT

Background: Jaundice is the yellowish discoloration of skin and mucous membranes due to high levels of bilirubin. Amongst the different types of jaundice, obstructive/surgical jaundice poses a great challenge due to varied presentation and high morbidity and mortality. Therefore, the present study was conducted to assess the same. **Method:** This cross-sectional, observational study was conducted on patients aged more than 18 years and diagnosed of obstructive jaundice. Demographic characteristics and relevant histories were noted. The findings of the laboratory investigations were recorded. Data was analysed. **Results:** It was noted that majority of the patients belonged to the age group of 61-70 years (44.40%). Females (52.78%) were affected more than males. Yellowish discoloration of the skin and sclera and abnormal/clay coloured stool were the commonest presenting symptoms reported by 91.67% and 83.33% patients, respectively. All patients had raised serum bilirubin levels and decreased serum proteins. Liver enzymes were increased in majority of the patients. **Conclusion:** Obstructive jaundice generally affects the elderly population with a female preponderance. Yellowish discoloration of skin and sclera and abnormal/clay coloured stools are the most commonly reported presenting symptoms with laboratory profile showing deranged liver function tests and decreased serum protein levels.

KEYWORDS : Obstructive jaundice, SGOT, SGPT, alkaline phosphatase, pruritus.

INTRODUCTION

Jaundice is derived from the French word 'Jaunisse'; Jaune in French meaning Yellow.¹ Jaundice, medically termed as 'Icterus', refers to the yellowish discoloration of the skin and mucous membranes. It is generally due to high levels of bilirubin in the body.²

Bilirubin, which is the major catabolic end product of heme after removal of its iron component, is primarily metabolized by the liver and excreted in the urine and stool.³ Accordingly, the causes for hyperbilirubinemia include pre-hepatic jaundice (due to haemolysis of red blood cells), hepatic jaundice (due to defect in capture, conjugation and excretion of bilirubin by liver) and post hepatic jaundice (due to the obstruction of extra hepatobiliary system). Depending upon the aetiology, jaundice may be either acquired or congenital.⁴ Though the cause of jaundice can often be correctly anticipated clinically but biochemical and radiological investigations are often required for confirmation of diagnosis prior to planning surgical management.

Depending upon the cause, jaundice may broadly be categorized as either obstructive and non-obstructive jaundice. While non-obstructive jaundice can be managed medically, obstructive jaundice often requires surgical management. It may occur due to either intra or extrahepatic organic obstruction to biliary outflow, i.e., a block in the pathway between the site of conjugation of bile in liver cells and the entry of bile into the duodenum through the ampulla. The obstruction may result from gall stones, strictures or malignant diseases, like cholangiocarcinoma, carcinoma of gall bladder or head of pancreas and periaampullary carcinoma. Rare causes include choledochal cyst, Caroli's syndrome and primary and metastatic liver tumours.⁵

Apart from the varied aetiology, the high morbidity and mortality associated with obstructive jaundice are also of concern.⁶ For planning surgical management, exact aetiology needs to be determined which may require an array of tests, both invasive and non-invasive, are often time consuming, expensive and have limited availability. Hence, maintaining a high index of suspicion is imperative for early diagnosis of cases of obstructive jaundice and prompt initiation of treatment to reduce the morbidity and mortality. Therefore, the present study was conducted to assess the clinicopathological spectrum of obstructive jaundice.

MATERIALS AND METHOD

This cross-sectional, observational study was conducted on a total of 36 patients aged more than 18 years, diagnosed with obstructive jaundice who were admitted during the study period. Patients aged less than 18 years or who were diagnosed with non-obstructive jaundice or who did not consent to participate in the study, were excluded from the study.

The study objectives and procedures were clearly explained to the patient in their vernacular language. A voluntary, written informed consent was obtained from all the patients. Demographic details were recorded. Detailed relevant clinical history was recorded. The presence of obstructive jaundice was confirmed by ultrasonography (USG) and contrast enhanced computed tomography (CECT) scan, wherever required. Routine investigations (complete blood count, liver and kidney function tests) and other special tests were conducted and recorded. The cut-offs for normal ranges were taken as:

1. Haemoglobin: More than 10 gm/dL
2. Total Leucocyte count: 4000 to 11000 cells/mm³
3. International Normalized Ratio (INR): 0.8 to 1.5 seconds
4. Random blood sugar: 80 to 130 mg/dL
5. Urea: Less than 20 mg/dL
6. Creatinine: Up to 1.2 mg/dL
7. Total bilirubin: Up to 1.2 mg/dL
8. Direct bilirubin: Up to 0.3 mg/dL
9. Serum Glutamic Oxaloacetic Transaminase (SGOT): Up to 40 U/L
10. Serum Glutamic Pyruvic Transaminase (SGPT): Up to 56 U/L
11. Alkaline Phosphatase (ALP): Up to 140 IU/L
12. Total protein: 6 to 8.3 g/dL
13. Albumin: 3.5 to 5.4 g/dL
14. Sodium: 135 to 145 mEq/L
15. Potassium: 3.5 to 5.5 mEq/L

Data was analysed using the SPSS software version 22.0. The qualitative parameters were expressed as number and percentages while quantitative parameters were described in terms of mean and standard deviation.

RESULTS

In the present study, it was noted that obstructive jaundice predominantly affects population in the age group of 61 to 70 years followed by the age group of 51 to 60 years. A slight female predilection was also noted (Table 1).

Table 1: Age Wise Gender Distribution of the Study Population

Age Group (in years)	Males		Females		Total	
	Number	%	Number	%	Number	%
21-30	0	0.00%	0	0.00%	0	0.00%
31-40	1	2.78%	0	0.00%	1	2.78%
41-50	1	2.78%	2	5.56%	3	8.34%
51-60	6	16.66%	6	16.66%	12	33.32%
61-70	7	19.44%	9	25.00%	16	44.44%
71 and above	2	5.56%	2	5.56%	4	11.12%
Total	17	47.22%	19	52.78%	36	100.00%

When assessed for presenting symptoms, yellowish discoloration of skin and sclera was the commonest symptom (91.67%), followed by abnormal/clay coloured stool (83.33%), pain in abdomen and pruritus being reported by 80.56% and 75% patients, respectively. Diarrhea (13.89%) and fever with rigors (19.44%) were the least common symptoms (Table 2).

Table 2: Distribution of Presenting Symptoms

Presenting Symptom	Number of cases	%
Yellowish discoloration of skin and sclera	33	91.67%
Abnormal/clay colored stool	30	83.33%
Pain in abdomen	29	80.56%
Pruritus	27	75.00%
Loss of weight	20	55.56%
Nausea/vomiting	14	38.89%
Lump	11	30.56%
Steatorrhoea	11	30.56%
Fever with rigors	7	19.44%
Diarrhoea	5	13.89%

When assessed for laboratory findings, it was noted that bilirubin (total and direct) and total proteins were deranged in all the patients, followed by urea (97.2%), ALP (91.7%) and albumin (91.7%). SGOT and SGPT were found to be elevated in 75% and 55.6% patients, respectively. INR was the least affected parameter being deranged in 22% patients only (Table 3).

Table 3: Distribution of the Findings of Laboratory Investigations

Parameter	Number	%	Mean	SD
Haemoglobin (g/dL)	29	80.6%	9.16	1.11
Total Leucocyte count (cells/mm ³)	17	47.2%	6211.28	4280.14
International Normalized Ratio (seconds)	8	22%	1.43	0.19
Random Blood Sugar (mg/dL)	24	66.7%	138.89	24.02
Urea (mg/dL)	35	97.2%	37.03	13.67
Creatinine (mg/dL)	14	38.9%	1.22	0.52
Total Bilirubin (mg/dL)	36	100.0%	15.51	5.48
Direct Bilirubin (mg/dL)	36	100.0%	10.87	4.10
SGOT (U/L)	27	75.0%	60.92	38.04
SGPT (U/L)	20	55.6%	62.25	21.48
Alkaline phosphatase (U/L)	33	91.7%	452.86	246.02
Total protein (g/dL)	36	100.0%	5.01	0.36
Albumin (g/dL)	33	91.7%	2.72	0.52
Sodium (mEq/L)	32	88.9%	131.67	3.45
Potassium (mEq/L)	24	66.7%	3.36	0.38

DISCUSSION

Obstructive jaundice is a challenging condition for the Surgeons in limited resource settings due to its varied presentation and high morbidity and mortality. Limited diagnostic facilities, in turn, affect the proper management of

the condition leading to further morbidity. The exact diagnosis of the underlying condition is crucial to the planning of surgical management, which may be time consuming. Hence, a high index of suspicion is required not only for early institution of treatment but also for protecting patients of non-obstructive jaundice from the exposure to the unnecessary plethora of tests. Therefore, this study was conducted to assess the clinicopathological features of obstructive jaundice.

The findings of the present study indicate 51 to 70 years being the most affected age group with female preponderance. Yellowish discoloration of skin and sclera, abnormal/clay coloured stools, pain in abdomen and pruritus were the commonest presenting symptoms being reported in more than 75% patients, each. Liver function tests are deranged in majority of the patients with significant derangement of liver enzymes.

Verma et al⁷ included 110 patients in their study. They reported the mean age of the patients to be 50.4 years. However, they reported a male preponderance in their study (56%). They further reported that jaundice (88%) and pain in the right upper quadrant (66.2%) were the commonest presenting symptoms. They assessed the biochemical profile and noted that total bilirubin concentration was elevated in more than 90% of the patients with 31% patients having concentrations more than 15 mg/dL. SGPT was elevated in 72% patients while SGOT was elevated in 60% patients. ALP was reported to be elevated in 70% patients. These findings were almost similar to the present study.

Irabor D.⁸ included 30 patients with obstructive jaundice in their study. They noted a male to female ratio of 1:1.4 with a significant female preponderance. The mean age reported in their study was 52.8 years. They observed that 66.7% patients with obstructive jaundice belonged to the age group of above 50 years. INR was reported to be deranged in 20% patients. Amongst the liver function tests, the serum levels of total bilirubin ranged from 7.1 to 24.2 mg% (mean level: 14.2 mg%), conjugated bilirubin levels ranged between 3 to 17.6 mg% (mean level: 10.2 mg%) and ALP levels ranged from 200 to 2850 IU/L (mean level: 900 IU/L). They hypothesized that deranged INR and increased risk of bleeding may be because of decreased production of Vitamin K dependent clotting factors due to back pressure and endotoxemia. They also noted that the risk increased with age and the decline in the synthesis of the clotting factors and clearance of gut derived endotoxins may be due to the ageing liver as well.

Mangam et al⁹ conducted their study on 108 patients of obstructive jaundice and reported that the most affected age group was 51 to 60 years with a female to male ratio of 1.2:1. Jaundice (100%) and pain in abdomen (68.86%) were the commonest symptoms.

In the study by Shukla et al,¹⁰ obstructive jaundice was found to predominantly affect patients aged more than 40 years (77%) with a significant female preponderance (61%). Pain in abdomen (97%), jaundice (81%) and itching (47%) were the commonly reported presenting symptoms while fever (15%), vomiting (13%) and ascites (7%) were the least common presenting symptoms. These findings were almost similar to the present study.

Sri et al¹¹ included 67 patients of obstructive jaundice to assess the clinical profile. They reported that patients aged more than 55 years were the most affected constituting 56.7% of the patients. However, they reported a slight male preponderance in their study (55.20%). Jaundice (97.01%) was the commonest symptom. These findings were almost similar to the present study.

The female preponderance may be attributed to their higher susceptibility to gall stones which are strong risk factors for the development of benign and malignant diseases in addition to being an independent contributor to obstruction.¹²

Amongst the liver enzymes, serum levels of SGOT change in response to the ongoing inflammation and hepatocellular necrosis while SGPT levels change according to the synthesis pattern. Therefore, SGOT changes reach peak faster and then decline once the condition is resolved while SGPT changes are comparatively slower. The study by Hayat JO et al has shown that in patients diagnosed with obstructive jaundice, especially due to gall stones, the rise in SGOT may be equal or even higher than SGPT.¹³

The decrease in mean levels of protein and albumin found in the present study may be because liver is the primary organ for synthesis of most of the plasma proteins except immunoglobulins. Albumin levels are an indicator of the synthetic function of liver. They are decreased in hepatocellular jaundice due to decreased hepatocyte function which may be due to hepatocyte necrosis in viral hepatitis. They are also decreased in obstructive jaundice due to the outflow obstruction for the synthesized product.¹⁴

Limitations

The present single centre study was limited by the availability of records of patients diagnosed with non-diabetic metabolic syndrome. Hence, the results may not be generalised.

CONCLUSION

Obstructive jaundice may have a varied aetiological spectrum but assessment of the clinicopathological spectrum may aid in maintain a high index of suspicion and exclusion of other differential diagnosis. The findings of the present study suggest that obstructive jaundice generally presents in the elderly age group of 61 to 70 years and has a female preponderance. Yellowish discoloration of skin and sclera, abnormal/clay coloured stools, pain in abdomen and pruritus are the most common presenting symptoms. The liver enzymes (SGOT, SGPT and ALP) show multi-fold increase in the cases of obstructive jaundice while serum proteins are invariably decreased. Thus, it is recommended that patients presenting with this profile be assessed for the presence of obstructive jaundice and workup for surgical treatment instituted at the earliest to decrease the associated morbidity and mortality.

REFERENCES

1. Srinidhi M, Hosmani R. A study of obstructive jaundice with focus on predictive factors for outcome. *Journal of Evolution of Medical and Dental Sciences*. 2014;3(29):8040-7.
2. Fargo M, Grogan S, Sagul A. Evaluation of Jaundice in Adults. *American Family Physician*. 2017;95(3):164-8.
3. Sticova E, Jirsa M. New insights in bilirubin metabolism and their clinical implications. *World Journal of Gastroenterology*. 2013;19(38):6398-407.
4. Abbas M, Shamshad T, Ashraf M, Javaid R. Jaundice: a basic review. *International Journal of Research in Medical Sciences*. 2016;4(5):1313-9.
5. Roche S, Kobos R. Jaundice in the Adult Patient. *Am Fam Physician*. 2004;69(2):299-304.
6. Ahmad I, Jan AU, Ahmad R. Obstructive Jaundice. *J Postgrad Med Inst*. 2001;15(2):194-8.
7. Verma S, Sahai S, Gupta P, Munshi A, Verma S, Goyal P. Obstructive jaundice-aetiological spectrum, clinical, biochemical and radiological evaluation at a tertiary care teaching hospital. *Int J Trop Med*. 2010;7(2):1-14.
8. Irabor D. The Risk of Impaired Coagulation in Surgical Jaundice: An Analysis of Routine Parameters. *Surgical Science*. 2012;03(03):116-9.
9. Mangam N, Dakhore S, Bodade R, Dhurve A, Dhoran A. Presentation, Etiology and Management of Obstructive Jaundice: A Prospective Study. *Journal of Medical Science And clinical Research*. 2018;6(7):207-15.
10. Shukla S, Kharat P, Kumar K. Clinicopathological study on patients presenting with obstructive jaundice. *International Surgery Journal*. 2018;5(2):705-10.
11. Sri TL, Chaitanya K. Clinical profile of patients with obstructive jaundice: A Surgeon's perspective. *International Journal of Scientific Research*. 2023;12(04):46-8.
12. Chalya PL, Kanumba ES, McHembe M. Etiological spectrum and treatment outcome of Obstructive jaundice at a University teaching Hospital in northwestern Tanzania: A diagnostic and therapeutic challenges. *BMC Res Notes*. 2011 May 23;4:147.

13. Hayat JO, Loew CJ, Asress KN, McIntyre AS, Gorard DA. Contrasting liver function test patterns in obstructive jaundice due to biliary strictures [corrected] and stones. *QJM*. 2005;98(1):35-40.
14. Carvalho JR, Verdelho Machado M. New Insights About Albumin and Liver Disease. *Ann Hepatol*. 2018;17(4):547-60.