



New Innovative Methods for Prediction of Hybridization State in a Very Short Time

KEYWORDS

Central atom, Peripheral atom, sigma bond, Lone pair or e-s, Co-ordinate bond, UG and PG Level

Dr Arijit Das

HEAD, Department of Chemistry, Govt. Degree College, Dharmanagar, Tripura(N), Tripura, India

ABSTRACT Prediction of hybridization state is a vitally important to students of chemistry in undergraduate, graduate and also in Post-graduate level. The method which is generally used for determination of hybridization state to find out the geometry is time consuming. To keep the matter in mind a new innovative method has to be introduced for calculation of hybridization state in a very simple way, which is also be a time savings one. Experiment in vitro on 100 number of students showed that for determination of Hybridization state, using old method, strike rate is 1Q/5min and by using these new innovative methods strike rate is 1Q/5secs. On the basis of this experiment I can strongly recommend that these new methods will be the very rapid one for the determination of hybridization state.

INTRODUCTION

A clear understanding and prediction of hybridization states are vitally important to students of chemistry in undergraduate, graduate and also in Postgraduate level to solve different kind of problems related hybridization state^{1,2,3}. The method which is generally used for determination of hybridization state is time consuming⁴. This new innovative method for prediction of hybridization states would go a long way to help to the students of chemistry who would choose the subject as their career.

We Know, Hybridization is nothing but the mixing of orbital's in different ratio to form some newly synthesized orbital's called hybrid orbitals. The mixing pattern is as follows:

Mixing	Hybrid orbital	Power of the Hybridization
s + p (1:1)	sp hybrid orbital	01
s + p (1:2)	sp ² hybrid orbital	02
s + p (1:3)	sp ³ hybrid orbital	03

Formula for determination of hybridization state like sp, sp², sp³ followed the following method:

Power of the Hybridization state of the centre atom = (Total no of σ bonds around each centre atom - 1)


All single (-) bonds are σ bond, in double bond (=) there is one σ and 1π , in triple bond there is one σ and 2π . In addition to these each lone pair (i.e.no of electrons in the outermost orbit which should not take part in bond formation) and Co-ordinate bond can be treated as one σ bond.

Eg:-

1. In NH₃, centre atom N is surrounded by three N-H single bond i.e. three sigma (σ) bonds and one LP i.e. one additional σ bond. So, altogether in NH₃ there is four σ bonds (3BP + 1LP) around centre atom N, So, in this case Power of the Hybridization state of N = 4-1 = 3 i.e. hybridization state = sp³.

2. In H₂O, centre atom O is surrounded by two O-H single bond i.e. two sigma (σ) bonds and two LPs i.e. two additional σ bonds. So, altogether in H₂O there is four σ bonds (2BPs + 2LPs) around centre atom O, So, in this case Power of the Hybridization state of O = 4-1 = 3 i.e. hybridization state of O in H₂O = sp³.

3. In H₃BO₃ (AIEEE-04) -



B has 3 σ bonds (3BPs but no LPs) and oxygen has 4 σ bonds (2BPs & 2LPs)

so, in this case power of the hybridization state of B = 3-1 = 2 i.e. B is sp² hybridized in H₃BO₃. on the other hand, power of the hybridization state of O = 4-1 = 3 i.e. hybridization state of O in H₃BO₃ is sp³.

4. In I-Cl-I and Cl both have 4 σ bonds (1BP and 3LPs), So, in this case power of the hybridization state of both I and Cl = 4-1 = 3 i.e. hybridization state of I and Cl both are sp³.

5. In CH₂=CH₂ - Each carbon is attached with 02 C-H single bonds (02 σ bonds) and one C=C bond (01 σ bond), so, altogether there is 03 sigma bonds. So, in this case power of the hybridization state of both C = 3-1 = 2 i.e. hybridization state of both C's are sp².

6. In N₂O (N≡N→O): The hybridization state of first N is sp due to one LP and one σ bond from N≡N and also the hybridization state of second N is sp due to one Co-ordinate bond and one σ bond from N≡N.

For determination of the hybridization state like sp³d, sp³d², sp³d³ followed the following method:-

In case of sp³d, sp³d² and sp³d³ hybridization state there is a common term sp³ for which 04 sigma bonds are responsible. So, in addition to 04 sigma bonds, for each additional sigma, added one d orbital gradually as follows-

5 σ bonds = 4 σ bonds + 01 additional σ bond = sp³d hybridization.

6 σ bonds = 4 σ bonds + 02 additional σ bonds = sp³d² hybridization.

7σ bonds = 4σ bonds + 03 additional σ bonds = sp^3d^3 hybridization.

Eg:- 1. IF_7 - 7 I-F single bonds i.e. 7σ bonds = 4σ bonds + 03 additional σ bonds = sp^3d^3 hybridization.

2. IF_4^+ - I has 07 e's in its outermost shell, so, in this case, subtract one e⁻ from 07 i.e. $07-1=06$. So, out of 06 electrons, 04 electrons form 04 I-F bonds i.e. 04 sigma bonds and there is one LP. So, altogether there is 05 σ bonds. So, 5σ bonds = 04σ bonds + 01 additional σ bond = sp^3d hybridization.

3. ICl_2 - I has 07 e's in its outermost shell, so, in this case, add one e⁻ with 07 i.e. $07+1=08$.

So, out of 08 electrons, 02 electrons form 02 I-Cl bonds i.e. 02 sigma bonds and there is 03 LPs.

So, altogether there is 05 σ bonds. So, 5σ bonds = 04σ bonds + 01 additional σ bond = sp^3d hybridization.

4. XeF_4 - Xe, an inert gas, consider 8 e's in its outermost shell, 04 of which form 04 Xe-F sigma bonds and there is two LPs, i.e. altogether there is 06 σ bonds = 04σ bonds + 02 additional σ bonds = sp^3d^2 hybridization.

In case of determination of the hybridization state you must have a clear idea about the outermost electrons of different family members in the periodic table as follows:

Family	Outermost electrons
Nitrogen family	05
Oxygen family	06
Halogen family	07
Inert gas family	08

And in case of cationic species you must remove one electron from the outermost orbit of the central atom and in case of anionic species you must add one electron with the outermost electrons of the central atom.

CONCLUSIONS: This article is very helpful to students in chemistry of undergraduate,

graduate and also in Postgraduate level. This one be the very time savings method. By using this method student can predict hybridization state in a very simple way.

ACKNOWLEDGEMENT: Author would be grateful to Prof. P. K. Chattaraj, Convenor, centre for Theoretical studies, Deptt. of Chemistry, IIT Kharagpur, India,; Prof. R.N. Mukherjee, Head, Deptt. of Chemistry, IIT Kanpur and Director, IISER, India and Ex-Fellow of Prof. R.H. Holm, Harvard University, USA,; Prof. Md. Ali, Deptt. of Chemistry, Jadavpur University, India,; Prof. G.N. Mukherjee, Sir Rashbehary Ghose Professor of Chemistry, Calcutta University, India,; Prof. R.A. Lal, North-Eastern Hill University (NEHU), Shillong, India, Prof. A.K. Das, Ex Vice-Chancellor of Kalyani University, Prof. R.K. Nath, Head, Deptt. of Chemistry, Tripura Central University, Prof. Nilashis Nandi, Kalyani University, W.B., India, Prof. Samar Kumar Das, University of Hyderabad, Prof. Partha Sarathi Mukherjee, Indian Institute of Science, Bangalore, Prof. V. Jagannadam, Osmania University, Prof. A. T. Khan, Head, IIT Patna and Dr. Satish Nimse, Hyllym University, South Korea for their recognition in this regard.

REFERENCE

1. Lee, J.D., Concise Inorg. Chem, 5th ed.; Wiley India, (2009). | 2. Douglas, B., McDaniel, D. and Alexander, J., Concepts and Models of Inorg. Chem., 3rd ed.; Wiley India, (2007). | 3. Cotton, F.A., Wilkinson, G. and Gaus, P.L., Basic Inorg. Chem., 3rd ed.; Wiley India, (2007). | 4. Mahan, B.M. and Meyers, R.J., International Student Edition University Chemistry, 4th ed. (1998), 599-603. |