



## Spectral Characterization And Evaluation of Antimicrobial Activity of Flavonoid From The Flowers of Glyceridia Sepium

### KEYWORDS

Glyceridia sepium, human pathogens, kaempferol, pyronoside, glycoside.

**C.Jayanthi**

PG&Research Department of chemistry Government college for women (Autonomous),kumbakonam. Tamilnadu,India

**Dr.A.JohnMerina**

Principal,Government college for women (Autonomous), kumbakonam. Tamilnadu,India

**ABSTRACT** The flavonoid was isolated by extraction and fractionation with different solvents from the fresh flowers of *Glyceridia sepium* belongs to the family fabaceae and it was identified as Kaempferol 3,7-O- $\beta$ -D-diglucoPyronoside with the help of UV, <sup>1</sup>H NMR & <sup>13</sup>C NMR spectroscopic techniques, then its antimicrobial activities against some human pathogens were studied by disk diffusion method. It has been proved that, the flavonoid possess anti microbial activity against all the tested strains and its zone of inhibition nearly equal to the drug Ciproflaxacin which was used as a standard drug. Thus the compound kaempferol 3, 7 - O -  $\beta$  - D - diglucoPyronoside obtained from the flowers of *Glyceridia sepium* is one of the potential element for therapeutic use

### INTRODUCTION

It is estimated that 2% of all carbon photosynthesized by plant is converted into flavonoids or closely related compounds<sup>1,2</sup>. Flavonoid is one of the largest groups of naturally occurring phenols. Flavonoids commonly occur as flavonoid -O-glycoside or flavonoid -C-glycoside<sup>1,2</sup>. These compound show a broad spectrum of biological activities<sup>4,5</sup>. In this investigation it was aimed to elucidate the structure and antimicrobial activities of flavonoid from the flowers of *Glyceridia sepium* belongs to the family fabaceae, it is a medium size leguminous tree and can grow 10 to 12 meters height. The flowers are located on the end of branches, these flowers have a bright pink to lilac colour that is tinged with white. A pale yellow spot is usually at the flower's base

### EXPERIMENTAL

#### Extraction and Fractionation:

Fresh flowers of 2kg of *Glyceridia sepium* collected from forest area of Muthandi Kuppam Village, Panruti, Tamilnadu, India were extracted with 85% methanol (5x500ml) under reflux. The alcoholic extract was concentrated in vacuo and the aqueous concentrate successively fractionated with benzene (3x250ml) ether (3x250ml) and ethyl acetate (4x250ml). The residue from ethyl acetate fraction called flavonoid came out as yellow solid. It developed red colour with Mg-HCl, brown colour with alc. Ferric soln, developed yellow colour when viewed under uv light with and without NH<sub>3</sub>, it responded with boron boric acid, molisch's test and Gibb's test. The flavonoid which was subjected to UV, <sup>1</sup>H NMR and <sup>13</sup>C NMR spectroscopic studies<sup>6,7</sup>

#### Antimicrobial Activity

The flavonoid was tested for its antimicrobial activity<sup>8</sup> against some human pathogens like *Escherichia coli*, *klebsil-*

*la pneumonia*, *pseudomonas aeruginosa*, *proteus vulgaris*, *streptococcus mutan* and *streptococcus pyogenes* by disk diffusion method<sup>9,10</sup>. Disks measuring 6mm in diameter were punched from whatman No.1 filter paper sterilized by dry heat at 140°C for an hour. They were immersed in solution which was prepared with different concentrations by using dimethyl formamide. Disks of each concentration were placed in nutrient agar medium inoculated with fresh bacteria strains separately. The incubation was carried out at 37°C for 24 hr. Ciprofloxacin was used as a standard at a concentration of 10 $\mu$ g/ml. After the incubation period is over the zone of inhibition produced by the flavonoid with different organisms in different plates were measured and recorded immediately by using zone reader<sup>11</sup>.

### RESULTS AND DISCUSSION


The UV Spectral values of the flavonoid showed two absorption maxima at 350 nm (band I) at 255nm (band II). Indicating the presence of glycosylation at 3-OH position. A stable bathochromic shift of 47 nm (Band I) noticed in NaOMe spectrum indicated the presence of free OH at. Abathochromic shift of 48nm in band I of flavonoid in the AlCl<sub>3</sub>-HCl spectra confirmed the presence of free-OH group at C<sub>5</sub>. In the <sup>1</sup>H-NMR data the signal at 6.2 ppm and 6.4 ppm corresponds to the proton at C<sub>6</sub> and C<sub>8</sub> respectively. The doublet appearing in the region of 8.6 ppm corresponds to the proton at C<sub>2'</sub> & C<sub>6'</sub>. While the proton of C<sub>5</sub> appears at 6.8 ppm. H-1" proton resonates at 5.4 ppm which indicates glucose moiety. The remaining sugar protons appear in the range of 3.0 to 3.7 ppm. The signal at 1.6 ppm indicates glycosylation at C<sub>3</sub>. The above details were assigned based on literature survey<sup>1,7</sup>. <sup>13</sup>C NMR spectral values of flavonoid were compared with the standard values<sup>7</sup> is shown at Table 1

Table : 1 Comparison of <sup>13</sup>C NMR Spectral values of flavonoid from the flowers of *Glyceridia Sepium* with standard values

|  | C <sub>4</sub> | C <sub>2</sub> | C <sub>3</sub> | C <sub>5</sub> | C <sub>6</sub> | C <sub>7</sub> | C <sub>8</sub> | C <sub>9</sub> | C <sub>10</sub> | C <sub>1'</sub> | C <sub>2'</sub> | C <sub>3'</sub> | C <sub>4'</sub> | C <sub>5'</sub> | C <sub>6'</sub> |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Kaempferol 3,7-O- $\beta$ -D-diglucoPyronoside from literature | 177.5          | 156.4          | 133.4          | 161.1          | 98.8           | 164.2          | 93.8           | 156.4          | 104.0           | 120.9           | 131.0           | 115.1           | 159.9           | 115.1           | 131.0           |
| Flavonoid  | 177.9          | 156.7          | 133.3          | 161.7          | 98.3           | 164.4          | 94.0           | 156.8          | 104.4           | 121.3           | 131.3           | 115.6           | 160.4           | 115.5           | 131.4           |

|  | C1''  | C2'' | C3'' | C4'' | C5'' | C6'' | C1''' | C2''' | C3''' | C4''' | C5''' | C6''' |
|--|-------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|
| Kaempferol<br>3,7-O-β-D-<br>digluPyranoside<br>from literature | 101.3 | 74.4 | 76.6 | 70.2 | 77.4 | 61.0 | 100.3 | 73.3  | 76.6  | 70.0  | 77.3  | 61.0  |
| Flavonoid  | 101.3 | 74.3 | 76.9 | 70.3 | 77.2 | 61.6 | 99.1  | 73.5  | 76.6  | 70.0  | 77.2  | 61.3  |

From UV, <sup>1</sup>H NMR & <sup>13</sup>C NMR spectroscopic studies it was confirmed that the structure of flavonoid present in the flowers of *Glyceridasepium* is kaempferol 3,7-O-β-digluopyronoside, is as follows.



**Kaempferol 3,7-O-β-D-digluopyronoside**

**Table : 2 Antibacterial Activity of Flavonoid Zone of Inhibition in (Millimeters)**

| Micro Organism Used                 |    |    |    |    |    |    |
|-------------------------------------|----|----|----|----|----|----|
| Drug                                | EC | KP | PA | PV | SM | SP |
| Ciproflaxacin<br>[Standard] 10μg/ml | 19 | 23 | 20 | 22 | 20 | 22 |
| Flavonoid at 50μg/ml                | 18 | 22 | 19 | 20 | 20 | 21 |
| Flavonoid 100μg/ml                  | 18 | 23 | 20 | 21 | 20 | 21 |

|    |   |                        |
|----|---|------------------------|
| EC | - | Escherichia Coli       |
| KP | - | Klebsiella Pneumoniae  |
| PA | - | Pseudomonas Aeruginosa |
| PV | - | Proteus Vulgaris       |
| SM | - | Streptococcus Mutans   |
| SP | - | Streptococcus Pyogens  |

Antibacterial activity of flavonoid isolate is nearly equal to ciproflaxacin which was used as a standard drug against the micro organisms shown in the table 2. From this, it was concluded that, the flavonoid isolate from flowers of *glyceridia sepium* has therapeutic effects.

#### ACKNOWLEDGEMENTS:

The authors are thankful to the Indian Institute Science, Bangalore, India for providing <sup>1</sup>H & <sup>13</sup>C NMR spectral data

#### REFERENCE

1. Markham K R, Techniques of Flavonoid identification, academic press (Chap and Hall) London, 1982 | 2. Dewick P M, The flavonoids Advances in Research, London 1982, 535 | 3. Jain A C and Tuli D.K, J.Science Industrial Research 1978, 287, 37, | 4. Jain A C and Nayyar K K Indian Journal of chem. 26 B, 1987, 136 | 5. Lane G A and Newman R H, Phytochemistry, 29, 1987, 26. | 6. Harbone. J.B, The flavonoids, London 1975, 743. | 7. Agrawal P K, carbon 13 NMR of flavonoids (Elsevier science publishers) London, 1989. | 8. Damtoft S, Franzy K H and Jensen S R, Phytochemistry 45, 1997, 743. | 9. Collins A.H, Microbiological methods, 2nd Edition, London, 1976. | 10. Simmons A, Practical medical microbiology 14th Edn (Churchill Livingstone) Edin berg 11, 1996, 163. | 11. Santilli A A, Kim H & Gregoy F J, J.Pharm science, 64, 1975, 1057. |