



Bioaccumulation of Heavy Metals in Fish Species of Ratnagiri Coast, Maharashtra

KEYWORDS

Bioaccumulation, Gills, Intestine, Tissue, Toxicity

Raju M. Patil

Department of Chemistry, Institute of Science, Madam Cama Road, Mumbai, Maharashtra, India.

Sagar T. Sankpal

Department of Chemistry, Institute of Science, Madam Cama Road, Mumbai, Maharashtra, India.

Pratap V. Naikwade

Department of Botany, ASP College, Devrukh, Maharashtra, India

ABSTRACT *The presence of heavy metals in our environment has been of great concern because of their toxicity when their concentration is more than the permissible level. Exposing human beings to these heavy metals directly through the food chain causes potential danger to human health. In present study the level of heavy metals, namely Zn, Cu, Hg, Pb and Cd in different fish species were evaluated using Atomic Absorption Spectrophotometer. The samples were collected between October 2012 to May 2013 from Ratnagiri coast. The metal concentrations in different body parts (Gill, Intestine, Liver, Kidney, and Muscle) were analyzed. The means of the amounts of heavy metals were estimated follows the order Zn > Cu > Cd > Pb > Hg. Heavy metals accumulated in higher concentration in intestine and low in muscles. The concentrations of metals in all samples were within permissible range set by World Health Organization.*

Introduction:

Metals generally enter the aquatic environment through atmospheric deposition, erosion of geological matrix or due to anthropogenic activities caused by industrial effluents, domestic sewage and mining wastes (Reddy et al., 2007). In the aquatic ecosystems, fishes are considered the important indicators of heavy metal enrichment of the aquatic ecosystems (Gernhofer et al., 2001). In an aquatic ecosystem, metals are transferred to the fish through food chain that could ultimately affect the health of people consuming this fish. Heavy metals are known for their persistent toxicity and tendency to bio-accumulate in aquatic ecosystems (Miller et al., 2002). Therefore, metals eco-toxicity of the aquatic ecosystems has become a major health concern over the years (Mendilet et al., 2010).

Biomonitoring of heavy metal pollution has been given ever-increasing attention due to the ability of various aquatic organisms to accumulate these heavy metals. Most of the salts of heavy metals and pesticides produce toxic effect on fish through physical accumulation (Tilaket et al., 2005). The worldwide reports indicate relatively wide variations in metal concentrations in sea water and sediment and fishes from different oceanic areas (Chester et al., 1974). Domestic sewage and industrial effluents are discharged in the water sources in and around India in untreated or partially treated form. These, add a variety of pollutants, which include certain toxic heavy metals (Sankpal and Naikwade, 2012). The heavy metals are accumulated in the marine environment then transfer to the marine organisms e.g. fishes by different ways. When their concentrations exceed the required levels, they become toxic and cause several health problems (Goldstein, 1994; Malik, 2004).

Material and Methods:

Ratnagiri district is one of the most important maritime districts of the state with the coastal belt extending to about 200 Km. Ratnagiri is an important coastal area of Maharashtra with average rainfall about 2500 mm. Most of the activities in this area are connected with sea. Recently several chemicals, pharmaceuticals companies and some power plants are grown up along the coastal region. Developmental activities like Konkan Railway Project, Enron electricity project, proposed marine highway, Cargo Ports are attracting more tourism industries in this region, which directly or indirectly causes environmental pollution. The marine area is presently receiving water with a variety of effluents, which may be potentially contaminating, including elevated levels of pollut-

ants (Agardet et al., 1988). Very little work is carried out on the distribution of heavy metals in fish samples of Ratnagiri coast.

Fish samples were collected from selected sites of Ratnagiri coast with the help of local fisherman during the period October 2012 to May 2013. Fish samples were transported to the laboratory in a thermos flask with ice on the same day. All fish samples were kept at -4°C until analysis. Sample preparation and analysis were conducted according to the procedure described by Bernhard (1976). Before analysis, muscle, liver, gill, intestine, and kidney were removed. Fish samples were homogenized in a blender and one gram of homogenate was digested. A microwave digestion system was used to prepare samples for analysis. Atomic Absorption spectrophotometer (Shimadzu AAS-680) was used for the analysis of metal.

Table.1 Accumulation of heavy metals in different tissues of fish

Fish Species Name		Concentration (ppm wet weight) in different body parts				
		Zn	Cu	Hg	Pb	Cd
Sardinella longiceps	Gill	19.36	12.86	0.060	0.798	0.218
	Intestine	29.57	6.30	0.030	0.522	0.336
	Liver	10.10	4.77	0.064	0.065	0.389
	Kidney	8.18	1.91	0.015	0.028	0.912
	Muscles	2.19	3.83	0.040	0.370	0.078
	Mean	13.88	5.93	0.042	0.357	0.387
Lepturacanthussavala	Gill	15.29	17.69	0.056	1.20	0.182
	Intestine	33.49	11.29	0.042	0.892	0.298
	Liver	12.29	6.29	0.097	0.098	0.287
	Kidney	9.27	2.01	0.018	0.032	0.998
	Muscles	2.39	4.39	0.045	0.283	0.876
	Mean	14.18	7.02	0.046	0.422	0.451

Lactariuslactarius	Gill	27.30	6.90	0.087	0.92	0.211
	Intes-tine	31.49	10.34	0.035	1.29	0.322
	Liver	16.26	7.28	0.102	0.096	0.267
	Kidney	11.93	1.92	0.019	0.029	0.987
	Mus-cles	3.36	5.39	0.039	0.321	0.987
	Mean	15.33	6.83	0.049	0.454	0.482

Result and Discussion:

The difference in the levels of accumulation in different organs of a fish can primarily be attributed to the differences in the physiological role of each organ. Other factors such as regulatory ability, behavior and feeding habits may play a significant role in the accumulation differences in the different organs (Marzouk, 1994). Also the chemical nature of the metals ionic strength and pH tends to be a master variable in the accumulation process.

The mean concentrations of heavy metals in the Gill, Intestine, Liver, Kidney and Muscle of selected fish samples are given in Table 1. The pattern of bioaccumulation of metals follows the order Zn > Cu > Cd > Pb > Hg. The lower concentrations of copper, zinc, mercury, cadmium and lead were usually recorded in muscles rather than the other organs while the higher values were recorded in the intestine for all samples. The intestine has a tendency to accumulate heavy metals in high values in fishes. The order of bioaccumulation is as follows: Intestine > Gill > Liver > Kidney > Muscle. While comparing different fish species studied, it was observed that *Lactariuslactarius* showed highest concentrations of Zn, Hg, Pb and Cd. However Cu concentration was highest in *Lepturacanthussavala*.

Wastewater discharge from sewage and industries are major component of water pollution in Ratnagiri coast (Naikwade et al. 2012). The impact of waste from industries also plays an important role in seawater pollution (Govindraj et al., 2011).

The study clearly indicated significant accumulation of heavy metals in the organs of the fish species from Ratnagiri coast. Results revealed that heavy metal like Zn was more concentrated in the intestine, Cd and Pb were more concentrated in kidney and muscles. When compared with international standards results showed that metal concentrations did not exceed WHO (World Health Organization, 1993) limit. Proper precautions should be taken by pollution controlling authority to avoid further pollution of Ratnagiri coast. Appropriate management strategies are needed to ensure the sustainable development and management of coastal areas and their resources.

REFERENCE

- Agard, J.B.R., Boodoosingh, M., and Gobin, J. (1988). Petroleum Residues in Surficial Sediments from the Gulf of Paria, Trinidad. *Mar. Poll. Bull.* Vol. 19, No.5. | Bernhard, M., (1976). Manual of Methods in Aquatic Environment Research. FAO Publ., Roma, Italy. | Chester, R and Stoner, J. (1974). The distribution of zinc, nickel, manganese, cadmium, copper and iron in some surface waters from the world ocean, *Mar. Chem.*, 2: 17-32. | Gernhofer M, Pawert, M, Schramm, M., Muller, E. and Triebkorn, R. (2001). Ultra structural biomarkers as tools to characterize the health status of fish in contaminated streams. *J Aquat Ecosyst Stress Recov*, 8: 241-260. | Goldstein, G. W. (1990). Lead poisoning and brain cell function. *Environ. Health Perspect*, 89: 91-94. | Govindaraju, M., Kumar, P., Selvaraj, M. and Rajina, C. (2011). Study on Physico-Chemical Parameters along the Coastal Waters around Kudankulam Nuclear Power Plant, *International Journal of Oceans and Oceanography* 5(1), 73-83. | Marzouk M., (1994). *Vet. Med. J.* 42 51. | Malik, A., (2004). Metal bioremediation through growing cells, *Environment International*, 30:261-278 | Mendil, D., Demirci, Z., Tüzen, M. and Soylak, M. (2010). Seasonal investigation of trace element contents in commercially valuable fish species from the Black Sea, Turkey. *Food and Chemical Toxicology*, 48: 865-870. | Naikwade, P., Mogle, U. and Sankpal, S., (2012). Phyloplanmycoflora associated with Mangrove plant *Ceriopstagal* (Perr.) *Science Research Reporter* 2(1): 85-87. | Reddy, MS., Mehta, B., Dave, S., Joshi, M., Karthikeyan, L., Sarma, VKS., Basha, S., Ramachandraiah, G., Bhatt, P. (2007). Bioaccumulation of heavy metals in some commercial fishes and crabs of the Gulf of Cambay, India. *Current Science*, Vol. 92, No. 11. pp. 1489-1491. | Sankpal, S. T. and Naikwade, P. V. (2012). Heavy metal concentration in effluent discharge of pharmaceutical industries, *Science Research Reporter*, 2(1), 88-90. | Tilak, K. S.; Veeraiyah, K. and Thathaji, P. B. (2005). *Journal Aqua. Biol.*, 20(1): 111-115. | WHO, (1993) Guidelines for Drinking-Water Quality, Volume 1, Recommendations. 2nd edition, World Health Organization, Geneva. |