



Effect of Furadan on Protein Content, Reduced Glutathione and Lipid Peroxidation Level on Kidney of *Bufo Melanostictus Schneider, 1799*

KEYWORDS

Bufo melanostictus, furadan, reduced glutathione, lipid peroxidation, kidney

Puspanjali Parida

P. G. Department of Zoology, North Orissa University, Baripada, Odisha, 757003 INDIA.

Nitya Sundar Pal

P. G. Department of Zoology, North Orissa University, Baripada, Odisha, 757003 INDIA.

Lakshmi Priya Mohanta

P. G. Department of Zoology, North Orissa University, Baripada, Odisha, 757003 INDIA.

ABSTRACT *Bufo melanostictus* (n=20) of various sizes (body weight: 90g- 120g) were divided into four groups of 5 animals each. Group I (control) animals received distilled water; Group II-IV (experimental) treated orally with 3µl per 1gm of body weight of furadan (0.005mg of furadan dissolved in 1ml of acetone). The treated animals were sacrificed after the time intervals of 24 hour, 48 hour and 72 hour (Group II-IV) whereas the control animal was sacrificed immediately (0h) and the kidney was dissected out and kept at 00C. The protein content, reduced glutathione level (GSH) and lipid peroxidation (LPX) level were measured in the kidney of *B. melanostictus* in both control and experimental group. Variation of different parameteres of kidney were observed at different time intervals. On the basis of the results, it is concluded that furadan even at low dose altered the biochemical parameters and induces oxidative stress.

INTRODUCTION

There is an increasing concern about the ecological effect of the use of pesticides. It has been reported that pesticide ingestion either by direct or indirect exposure may lead to generation of reactive oxygen species (ROS), which are detrimental to the health of humans and non-target organisms (Otitoju and Onwurah, 2007). A good bioindicator should have a well-known taxonomy and ecology, be distributed over a broad geographic area, have a high degree of ecological specialization as specialized species are far more vulnerable to environmental perturbations compared to generalists, be cost-effective and relatively easy to survey (Rainio and Niemela 2003; Brischoux et al. 2009).

Although pesticides have long been suggested as a possible cause of amphibian declines (Carey and Bryant 1995, Stebbins and Cohen 1995, Drost and Fellers 1996, Lips 1998), there have been few toxicological studies on declines. Carbofuran or furadan (2, 3-dihydro -2, 2-dimethyl-7-benzofuranyl-N-methyl-carbamate) has been reported to have relatively high mammalian toxicity and very toxic to invertebrates and birds and should therefore be handled with a lot of care (Hodgson et al., 1991).

The kidney plays important role as a natural detoxifying organ. In the present study generation of ROS in kidney of *Bufo melanostictus* in response to furadan were estimated by measuring reduced glutathione and lipid peroxidation level after different time intervals of 24 hour, 48 hour and 72 hour and compared against the control (0 hour).

MATERIALS AND METHODS

B. melanostictus were collected locally near the North Orissa University campus, during night and early morning time. They were acclimatized for seven days prior to the experiment. *Bufo melanostictus* (n=20) of various sizes (body weight: 90g- 120g) were divided into four groups of 5 animals each. Group I (control) animals received distilled water; Group II-IV (experimental) treated orally with 3µl per 1gm of body weight of furadan (0.005mg of furadan dissolved in 1ml of acetone). The treated animals were sacrificed after the time intervals of 24 hour, 48 hour and 72 hour (Group II-IV) whereas the control animal was sacrificed immediately (0h).

The liver of both control and experimental group were dissected out quickly and kept at 0°C. A 20% homogenate was prepared with phosphate buffer (pH 7.4). The tissue homogenate was centrifuged at 4000 rpm for 10 minutes.

Protein Estimation

Protein estimation of the sample were made according to the method of Lowry et. al.,(1961). To 0.1ml suitably homogenate of tissue 0.4ml of distilled water was added. Then 5 ml of biuret reagent (containing alkaline Na₂CO₃, 0.5% CuSO₄ solution and 1% Sodium potassium tartarate solution in the ratio 100:2:2) was added and properly mixed up. After 10 minutes of incubation at room temperature 0.5ml of Folin Ciocalteu phenol reagent (the commercial reagent diluted three times with distilled water) was added and incubated at 37°C for 30 minutes at room temperature. Absorbance was measured at 700 nm against an appropriate blank

Reduced Glutathione Assay

Reduced glutathione of the sample were estimated by Ellman (1959) method. 0.7ml of the tissue homogenate was added to 0.7ml of TCA. Then the substances in the tubes were centrifuged at 4000 rpm for 10 minutes. 0.5ml supernatant was added to 2.5ml of DNTB (DNTB 30 mM) was diluted in PO₄ buffer 100 times. The absorbance was taken at 412 nm with in between 5 to 30 minutes against a appropriate blank.

Lipid Peroxidation Assay

Lipid peroxidation of the sample is estimated as thiobarbiturate acid reacting substance (TBA-RS) by thiobarbituric acid (TBA) according to the method of Ohkawa et. al. (1979). 3.8ml of TBA reagent contain (2ml of 8.1% SDS, 1.5ml of 20% acetic acid of pH 3.5, 1.5ml of 0.8% aqueous solution of TBA, 5ml of distilled water and 1ml of BHT) was added to 0.2ml of suitably diluted post nuclear supernatant. After mixing thoroughly, the test tube's substance was boiled in water bath for 1 hour. The tubes were cooled down to the room temperature. Then the tube substances were centrifuged at 4000 rpm for 10 minutes. The absorbance of the supernatant was measured at 532 nm against a appropriate blank.

Table 1: Comparison of protein content (mg/g tissue), GSH level (µ mol/g tissue), LPX level(n mol TBARS/mg protein)of kidney in *Bufo melanostictus* after treatment of furadan (3µl/g body weight) at different time interval. The value are expressed in Mean± S.D.

Duration after treatment with furadan	Protein content (mg/g tissue)	GSH level (µ mol/g tissue)	LPX level (n mol TBARS/mg protein)
0h (Group-I)	74.09±0.729	0.16±0.001	12.82±0.191
24h (Group-II)	43.60±0.484	0.09±0.001	15.90±0.222

48h (Group-III)	41.44±0.434	0.093±0.001	33.94±0.323
72h (Group-IV)	55.63±0.298	0.176±0.002	19.08±0.252




Fig : 1 Comparison of protein content (mg/g tissue) of kidney in *Bufo melanostictus* treated with furadan (3µl/g body weight) at different time interval.




Fig : 2 Comparison of GSH level (µ mol/g tissue) of kidney in *Bufo melanostictus* treated with furadan (3µl/g body weight) at different time interval.




Fig : 3 Comparison of LPX level (n mol TBARS/mg protein) of kidney in *Bufo melanostictus* treated with furadan (3µl/g body weight) at different time interval.

Duration after treatment with furadan	Protein content (mg/g tissue)	GSH level (µ mol/g tissue)	LPX level (n mol TBARS/mg protein)
0h (Group-I)	74.09±0.729	0.16±0.001	12.82±0.191
24h (Group-II)	43.60±0.484	0.09±0.001	15.90±0.222
48h (Group-III)	41.44±0.434	0.093±0.001	33.94±0.323
72h (Group-IV)	55.63±0.298	0.176±0.002	19.08±0.253

RESULTS AND DISCUSSION

Protein content (mg/g tissue) in the kidney of *B. melanostictus* treated with furadan were 74.09 ± 0.729 mg/g tissue , 43.60 ± 0.484 mg/g tissue , 41.44 ± 0.434 mg/g tissue and 55.63 ± 0.298mg/g tissue at 0hr,24hr,48hr and 72 hr respectively (Table 1, Fig1). The protein content of kidney tissue of *B. melanostictus* treated with furadan is highest at 0 hour (control) and then decline at 24h and 48h, but then increase drastically. It indicate, the animal require some time(more than 48h) to produce more protein.

GSH content (µmol/g) tissue in kidney tissue of *B. melanostictus* treated with furadan were 0.16 ± 0.001 µ mol/g tissue , 0.09 ± 0.001 µ mol/g tissue, 0.093± 0.001 µ mol/g tissue and 0.176 ± 0.002 µ mol/g tissue at 0hr,24hr,48hr and 72 hr respectively (Table 1, Fig 2). The GSH level decreases gradually at 24 hr and 48 hr but increases significantly at 72 hr like that of protein content means the animal is able to compensate oxidative stress at 72 hr.

The LPX level (n mol TBARS/mg protein) in kidney tissue of *B. melanostictus* treated with furadan is 12.82 ±0.191 n mol TBARS/mg protein , 15.90 ± 0.222 n mol TBARS/mg protein, 33.94 ± 0.323 n mol TBARS/mg protein, 19.08 ± 0.253 n mol TBARS/mg protein at 0hr,24hr,48hr and 72 hr respectively. The LPX level of kidney tissue of *B. melanostictus* treated with furadan increases gradually upto 48 hr but drastically decreases at 72 hr may be due to the increase of GSH at 72hr which reduce the lipid peroxidation.

In otherword, kidney of *B. melanostictus* have well developed GSH system to reduce oxidative stress (LPX) and it require more than 48 hr or 72 hr to activate the antioxidant system.

REFERENCE

1. Brischox F., Bonnet X. & Legagneux P. 2009. Are sea snakes pertinent bio-indicators for coral reefs? A comparison between species and sites. *Marine Biology* 156: 1985-1992. | 2. Carey, C., and C. J. Bryant. 1995. Possible interrelations among environmental toxicants, amphibian development, and decline of amphibian populations. *Environmental Health Perspectives* 103:13-17. | 3. Drost, C. A., and G. M. Fellers. 1996. Collapse of a regional frog fauna in the Yosemite area of the California Sierra Nevada, USA. *Conservation Biology* 10:414-425. | 4. Ellman, G.L. 1959. Tissue sulphhydryl groups. *Arch. Biochem. Biophys.* pp 82:70-77. | 5. Hodgson, E.; Roe, R.M. and Motoyama, N. 1991. Pesticides and the Future: Toxicological studies of risks and benefits. *Rev. Pestic. Toxicol.* 1: 3-12. | 6. Lips, K. R. 1998. Decline of a tropical montane amphibian fauna. *Conservation Biology* 12:106-117. | 7. Lowry, O.H., Resbrough, N.J., Farr, A.L. Randall, R.J. 1951. Protein measurement with the Folin-phenol reagent. *J.Biol. Chem.* 19: 265-275. | 8. Ohkawa H., Ohishi, N. and Yagi, K. 1979. Assay of LPX in animal tissues by thiobarbituric acid reaction. *Anal Biochem. Physiol.* 118C (1):33-37. | 9. Ootokujo O, Onwurah I.N.E. 2007. Glutathione S-transferase (GST) activity as a biomarker in ecological risk assessment of pesticide contaminated environment. *Afr. J. Biotechnol.*, 6: 1455-1459 | 10. Rainio J. & Niemela J. (2003). Ground beetles (Coleoptera: Carabidae) as bioindicators. *Biodiversity and Conservation* 12: 487-506. | 11. Stebbins, R. C., and N. W. Cohen. 1995. *A natural history of amphibians*. Princeton University Press, Princeton, New Jersey, USA. |