



***Carica Papaya* - A Potential Host Of *Philosamia Ricini* Eri Silkworm Under Western Maharashtra Condition**

KEYWORDS

Eri silkworm, *Philosamia ricini*, *Carica papaya*, western Maharashtra

KAVANE R.P.

Department of zoology, Y.C.W.MAHAVIDYALAYA., WARNANAGAR

ABSTRACT

An attempt was made to rear the eri silkworm, *Philosamia ricini* on the leaves of *Carica papaya* as a primary food for the first time in India and reared under laboratory condition. It was observed that the papaya is a potential host of eri silkworm. The growth parameters and cocoon characters of the eri silkworm reared on papaya leaves were normal. The eri silkworm successfully lasted its life cycle ranged from 47 to 53 days. The effective rate of rearing, weight of pre-spinning larvae, cocoon, shell and pupae ranged from 45 to 55 per cent, 6.10 to 7 g, 2.49 to 2.70 g, 0.30 to 0.32 g, 2.10 to 2.60g, respectively. Parameters such as length of cocoon shell, width of shell, shell thickness etc. The fecundity ranged from 350 to 415 eggs.

Introduction

Eri silkworm, *P.ricini* is a multivoltine sericigenous insect and largely reared by the farmers of North-eastern part of India, particularly in Assam (Sahu et al., 2006). The eri culture being carried out throughout the year in traditional areas because of the abundant availability of castor plants in the rural areas (Rao et al., 2005; Siddique 2009). The silk of the castor silkworm is produced by *Philosamia ricini*. Assam in India is the home of the eri silk industry (FAO, Manual, 1987). Sericulture being an important agro based industry provides employment at various levels i.e. host plant cultivation, silkworm rearing, reeling, spinning and weaving have much impact on the improvement of rural economy. The ambient temperature during rearing affects larval growth, survival rate, cocoon parameters and silk quality (Gumma, 1972). The quality of feed plays a remarkable role for growth and development of the silkworm and ultimately on the economic traits of cocoons (Hazarika et al., 2005).the eri silkworm feeds on many food plants like primarily castor, secondary food plants like kessaree, tapioca, payam, jatropa, papaya etc. All the food plants are not equally good for eri silkworm rearing and eri silkworm show different behavior, when reared on different food plants (Rajesh Kumar and Gangwar, 2010). Based on the above cited literature and information the present study was conducted to suitable food for *P.ricini* under western Maharashtra condition. Therefore, an attempt has been made to rear the insect from 1st instar to 5th instar & allow for cocoon and adult formation.

Materials and methods

The eggs of *P.ricini* were brought from eri research station, CSB, Assam. The eggs were incubated at room temperature 28 ± 2°C, 80 ± 5% RH and 10 hrs photoperiod. For this study the rearing methodology developed for wild silkworm was followed (Kavane, 2010). Experiment was conducted from newly hatched larvae to adult formation. Sum of 500 worms were maintained in plastic boxes and G.I. tray are specially prepared for new technique to wild silkworm rearing. The host plants tender and soft leaves were fed one time a day up to 2nd instar and late age silkworms one times a day. Cloth boxes specially prepared montages by the ripening larva. Cocoon harvesting was carried out after fifth day of spinning. The data was recorded for the study of biology, which includes larval duration, adult longevity and fecundity, effective rate of rearing (ERR), cocoon characters, colour, shape, single cocoon weight, shell

weight etc. the above species of silkworm sufficient number of silkworm (500) were used for confirming results.

Results and discussion

The biology of saturniid, *P.ricini* was studied under laboratory condition. The descriptions were made from live specimens reared during 2012- sep.2014. It was observed that there was less mortality of 1st instar worms when fed with tender, soft papaya leaves. The larvae of all the five instars fed upon tender, soft matured leaves and successfully complicated the life cycle. The result indicated that the rearing success of *P.ricini* on papaya under indoor rearing conditions (28 ± 2 °C and RH of 80 ± 5% and 10 hrs photoperiod) was 45 %. The moths mated successfully, laid fertilized eggs in the mud pot and covered with nylon mesh in rearing house. On papaya leaves, the incubation period, larval and pupal duration ranged from 10 days, 24 to 25 days, and 14 to 15 days respectively. The *P.ricini* silkworm molted four times and there were five instars each lasted 3 to 3.5, 2 to 2.1, 4.5 to 4.7, 6.4 to 6.6 and 8 to 9 days, respectively. The weight of matured tasar silkworm (vth instar) was 6.22 to 6.50 g and measured 6.35 to 6.90 cm in length, 1.87 to 1.95 cm in circum. The weight of cocoon including pupa 2.85 to 3.15 g, the shell weights of cocoon 0.30 to 0.40 g and the shell lengths were 4.85 to 4.87 cm, the shell width of cocoon in indoor method were 1.90 to 2.20cm , silk ratio 14.90 to 15.00% (Table.1).

The effective rate of rearing (ERR), adult longevity and fecundity ranged from 45 to 55 per cent, 3 to 5 days and 350 to 415 eggs, respectively. The eggs hatchability was 97 to 100 percent and it was maximum on first day between 8 to 10 am. Further the life cycle of insect ranged from 49 to 52 days. The results of the study like performance of *P.ricini* silkworms in (Table. 2) and the width of the different life stages are presented (Figs.1 to 8).

Newly hatched larvae were released on the leaves of the host plants badam with the help of soft camel hairbrush. 4 - 5 such leaves with mounted larvae were placed in the plastic box size 27 cm x 6 cm x 7 cm in length, width, and height respectively. The maximum portion of edge of each leaf was available to the larvae for feeding; the box was perforated with numerous exits for aeration and with covered lid to prevent escape of the larvae. Next day, the larvae were transferred to new box containing fresh leaf diet. The molt-

ing larvae transferred along with their support leaves. The old leaves were removed from the boxes at 12 hrs interval. Rearing of 1st and 2nd instars for used boxes were then washed, disinfected and dried for re-use (Kavane,2010).


Rearing trays of size (3'x 2.5'x 6'in length x width x height respectively) were used for rearing of 3rd, 4th and 5th instar larvae. A twig having 12 to 15 leaves was used as leaf diet 8 – 9 leaf twigs were introduced in the tray at a time. Larvae were transferred to a new tray along with help of new food. The trays were cleaned after 24 hr interval. Molting larvae were transferred along with the left over parts of the food plant.

The full grown 5th instar larvae wandering for cocoon construction were sorted out and transferred to a box (4'X 2.5'in height X width) made up of hard card sheet provided them opportunity to form the cocoons (Kavane,2010). The data was recorded for the study of biology, which includes larval duration, adult longevity and fecundity.

Eggs (Fig.1)



The eggs deposited in the mud pot were collected, washed with soap water and kept for incubation. The eggs are rounded; newly laid eggs look like brownish but become creamy white colour. An individual egg measured 0.16 to 0.17 mm and breadth 0.12 to 0.13 mm, hatching percentage is 100 percent and incubation period of 10 days. The emergence of the larva from the egg commenced from morning and continued till 10 a.m.



Figs.(1-8). Rearing of *Philosamia ricini* on *Caraca papaya*
Fig.1.Fertilised eggs, Fig.2.1st instar, Fig.3. IInd instar, Fig.4. IIIrd instar, Fig.5.IVth instar, Fig.6.Vth instar, Fig.7.Cocoon, Fig.8.Mating of moth.

Larva: (Fig.2-8)

First instar larva head black, body colour yellow with black lining and hairs, tubercles conical, legs black, black band present on the dorsal side. The length of first instar ranged from 0.74 mm. the duration of first instar ranged from 3.5 to 3.7 days with a mean of 3.5 days. The second instar head black, body colour yellow without black lining and withes short black tubercles with whitish hair. Body bears pairs of black spot longitudinally. The length of 2nd instar ranged from 1.65 to 1.68 cm. the duration of second instar ranged from 2 to 2.2 days with a mean of 2.1 days.

The third instar larva measured 2.34 to 2.35 cm with a mean of 2.35 cm. head black, body colour white with short white tubercles with powdery. The body bears longitudinal black spots, dorsally in pairs and dorsoventrally in irregular manner, legs, anal flap and claspers become yellow. The duration of third instar ranged from 4.5 days. The fourth instar ranged from 4.20 to 4.60 cm with a mean of 4.40 cm in length, head yellow, body colour white with short white tubercles with powdery. The duration of fourth instar ranged from 6 to 7 days with a mean of 6 days. The fifth instar larvae measured from 6.15 to 6.29 cm with a mean of 6.22 cm in length, head yellow, body colour white with short white tubercles with powdery. The duration of fifth instar ranged from 8 to 9 days with a mean of 8 days. The fifth instar single larval weight varied between 6.60 to 7.30 g. with an average of 6.6.22 g. the total larval duration ranged from 24 days. The effective rate survival was as 45 percent.

At the end of the larval period the ripe larvae were released on the montages for cocoon construction on which cocoons were constructed. Matured larvae crawl down for cocooning in afternoon to till noon. Cocoons white, elongate, spindle shaped with a thin floss layer and it can be easily distinguished. It measures 2.49 to 2.60 g., 4.85 cm, 1.90 cm weight, length, width respectively. The open types of cocoons are compact and hard without peduncle. The pupa is dark brown or radish brown. Female pupa is bigger and heavier than male. The male moth has pointed wings while females has broad wing. The wing span of the male and female moth is 115 to 125 mm and 135 to 145mm. respectively. Wing colors dark brown. There have been no reports on the primary feeding habits of eri silk worm on papaya, a potential alternate host plant. The present finding has firmly established that the eri silkworm *P.ricini* could be reared on papaya leaves.

The effect of host plant species on the growth and development in the insects has been reported (Reddy et al., 1989). Impact of varietal feeding on samia ricini inspring and autumn season of Utter Pradesh studied (Rajesh Kumar and Gangwar, 2010). Hazarika et al. (2003) studies that castor was found best in terms of different parameters viz, larval wt, ERR, cocoon weight, shell weight etc. Performance of promising ecoraces of eri in agro climatic condi-

tions of western Odisha region studied (Ray et al., 2010). Rearing technology of eri silkworm under varied seasonal and host plant conditions in Tamilnadu reported (Subramanian et al., 2013). In the present study, some of the characters like hatching percentage, larval duration, larval weight, cocoon yield, single cocoon weight, shell weight, shell ratio, etc on *C. papaya* food plant is new investigations under western Maharashtra climatic condition.

Table 1: Cocoon characters of *P.ricini* silkworm on papaya

Fecundity no/worm	Hatch %	Larval wt(g)	ERR%	Cocoon wt(g)	Shell wt(g)	S.R%	Larval period
425	100	6to7	45%	2.49	0.30	12.04	24-25

Table 2. Rearing performance *P.ricini* silkworm on papaya

Life stages	Duration (days)	Feeding time per day	Leaf number on food plant twig	Leaf size	No of boxes/ cages	No of trays	Box/tray cleaning time	Duration of shedding Cuticle(hrs)	Humidity %	Temp. °c
Eggs	10	---	---	-----	-----	-----	-----	-----	80 – 85	28-30
1 st instar	3.5	one	Tender	Whole	2	-----	1	12	80 – 85	28-30
2 nd instar	2	Two / June	Tender	Whole	4	-----	1	18	80 – 85	28-30
3 rd instar	4.5	Two	June	Whole	-----	2	2	18	80 – 85	28-30
4 th instar	6	Two	June	Whole	-----	3	3	24	80 – 85	28-30
5 th instar	8	Two	June	Whole	-----	4	4	-----	80 – 85	28-30
pupal	15	---	-----	-----	-----	-----	-----	-----	80 – 85	28-30
Adult longevity	3-4	---	-----	-----	-----	-----	-----	-----	80 – 85	28-30

Acknowledgement

Author is thankful to Principal Yashwantrao chavan warana Mahavidyilia, Warananagar, Kolhapur for providing Facilities

REFERENCE

- Sahu,M., Bhuyan,N&Das,P.K.(2006) Eri silkworm, *Samia ricini* Donovan, seed production during summer in Assam. In: proceeding of Regional seminar on "Prospects and problems of sericulture as an economic enterprise in north West India, dehradune, India, 490-493. | Siddique, A.A.(2009). Agricultural activity and castor plantation for eri silk production. *Indian silk*, (6):37-40. | Rao, R.M., Prasad, R.N & Suryannrayan, N.(2005). *Eri culture: an additional income for tapica growers and good nourishment to the man- nutitions tribal populace*. In:proceedings of 20 th congress of the international sericulture commission volume II,sce,3 non-mulberry silkworm, ,Bangalore india,94-98. | Gomma,A.Ahmad.(1972) . Biological study on the eri silkworm, *Attacus ricini* bosid. *Indian J.seric.*,(1):81-88 | Hazarika,U.,Barah,A& chakravorty ,R.(2005). Physiological and biological response of castor to application of NPK and their correlation with economic parameters of eri silkworm. In:proceedings of 20 th congress of the international sericulture commission volume II,sce,3 non-mulberry silkworm, Bangalore, india,94-98. | Peigler, R.S., Naumann, S. (2003). A Revision of the Silkmoth Genus *Samia*. University of Incarnate Word,San Antonio, Texas. pp: 227-228. | Rajesh Kumar, Elangovan, V. (2010). Assessment of the volumetric attributes of eri silkworm (*Philosamia ricini*) reared on different host plants. *I.J.S.N.*, VOL. 1(2): 156-160. | Rajesh Kumar, Gangwar, S.K. (2010). Impact of varietal feeding on *Samia ricini* Donovan in spring and autumn season of Uttar Pradesh. *ARPN Journal of Agricultural and Biological Science.*, 5 (3). | Deka, mukal., D. saranga and Dilali devi (2011). Impact of feeding of *Samia Cynthia ricini* boisduval (red variety) in respect of larval growth and spinning. *Int.J.Pure.Appl.sci.technol.*, 5(2).131-140. | Ray, P.P., T. V. Rao and Purushattam Dash.(2010). Performance of promising ecoraces of eri in agroclimatic conditions western Odisha. *The Bioscan* .5(2):201-205. | FAO,(1979) *Manuals on sericulture*, central silk board,Bangalore. | Subramanian, K., N. Sakthivel and S.M.H.Qadri.(2013).Rearing technology of eri silkworm (*Samia Cynthia Ricini*) under varied seasonal and host plant conditions in Tamilnadu, *Int. J. Life sc. Bt & Pharm. Res*, 2(2).130-141. | Kavane , R.P.(2010). *Biosystematics of wild silkmoths from western Maharashtra.. Ph.D. thesis , shivaji university, Kolhapur*.1-236. | Kavane , R.P., and T.V.sathe.(2011). *Wild silk technology*. D.P.H, Newdelhi.1-236 |