



Chylolymphatic cyst - a case report'

KEYWORDS

Mesenteric Cyst, Chylolymphatic Cyst,

Dr. Vaibhav Mane

Vaishnavi Pathology Laboratory Shop No.2 Kore House, Vishrambag, Sangli

ABSTRACT *Cysts of the mesentery are among surgical rarities and of varied aetiology with variable presentations and this has surgical implications in the pediatric age group.[1,2,35,6,7,]*

They may be derived from the gastrointestinal tract, the genitourinary system, previous inflammation (pseudocysts) or malignant cystic tumours, but the commonest cause is generally considered to be a congenital lymphatic cyst. [1,4,6,7,8,]The clinical presentation is not characteristic and in addition, the preoperative imaging although suggestive is not diagnostic.

In most cases, the diagnosis is confirmed after surgical exploration and removal of the cyst.[1,2,3,4,]

A case report of a baby aged 1 day is being reported

Introduction

A chylolymphatic cyst is a rare variant of a mesenteric cyst [1,2,3,4,5,6,]. The mean age of children affected is 4.9 years [1,2,6,8,] These cysts present within the mesentery, lined with a thin endothelium or mesothelium and filled with chylous and lymphatic fluid [1,2,3,6,7,]. Although mesenteric cysts in general have been reported in the literature fairly frequently, chylolymphatic cysts in the pediatric age group are extremely rare in the modern medical literature [1,2,] therefore very little information is available regarding their presentation and complications.

Case Report

A 1 day old male child, with a full term normal hospital delivery, presented with complains of excessive crying, diarrhoea mixed with blood and mucus and abdominal pain. The pain was intermittent, colicky, poorly localized and was mild to moderate in intensity. On clinical examination, there was no palpable abdominal mass. Plain abdominal radiography was normal.

Abdominal computed tomography was done which showed well defined small hypodense cystic lesions noted in infraumbilical region intraperitoneally in right paramedian location, just beneath the rectus abdominis muscle. Patient underwent diagnostic laparoscopy which showed a single chylous looking mesenteric cyst adherent to appendix.. There was a single cyst of size approximately 2 cm in diameter. Postoperative period was uneventful and the patient recovered well.

The specimen was sent for histopathological examination, which revealed multiloculated cysts lined with endothelium and filled with chylous fluid and lymph. Thus, the case was confirmed to be chylolymphatic cyst by histopathology.

Discussion

Chylous cysts are rare variants of mesenteric lesions and constitute 7.3% to 9.5% of all abdominal cysts [1,2,3,5,7,8,]. There are very few cases of pediatric chylolymphatic cysts reported in the literature. Behrs et al. [4] classified mesenteric cysts into four groups based on etiology: embryonic or developmental; traumatic or acquired; neoplastic; and infective or degenerative.

Recently, a pathologic classification system has been proposed [3,4,5]. Types 1 (pedicled) and 2 (sessile) are limited to the mesentery and can be excised completely with or without resection of the involved gut. Types 3 and 4 (multicentric) extend into the retroperitoneum and require complex operations and often sclerotherapy as well. Based on the contents of the cyst, the mesenteric cyst can be divided into serous, chylous, hemorrhagic and chylolymphatic cyst [1,2,3,6,]. The chylolymphatic cyst, as indicated by its name, contains both chyle and lymph. [1,2,3,4,5,6,7,]The accumulation of chyle and lymph is considered to be the result of an imbalance between the inflow and outflow of fluid [1,2,3,7,8,]. This cyst may be asymptomatic, and may cause abdominal distension or an abdominal [1,2,3,4,]lump or may present with complications such as intestinal obstruction, hemorrhage, infection, rupture of the cyst, volvulus or obstruction of the urinary or biliary tract[1,2,] Radiological investigations form an integral part of the management of these lesions. [1,4,5,6,7,]

Antenatal detection of cystic abdominal lesions is possible in a fetus during antenatal ultrasound scanning. The sonologic picture may help in differentiating the lesion from many other differential diagnoses[1,3,4,]. As this usually does not alter the obstetrical management, a definite diagnosis is usually made in the postnatal period [6,7,8].

The different surgical approaches used are marsupialization, sclerotherapy, drainage, enucleation, percutaneous aspiration, and excision of the cyst with or without resection of the involved gut [1,2,5,6,7,8,]. Due to high recurrence rates associated with marsupialization and drainage, complete excision of the cyst should be attempted whenever possible [4,5,]. In adults, the cyst can often be enucleated or 'shelled out' from between the leaves of the mesentery; in children, however, bowel resection is frequently required [6,7,8]. The medical literature mentions instances where laparoscopic removal of mesenteric cysts has been tried successfully [3,4,5].

Intra-operatively, similar findings can be seen in cystic lymphangioma, retroperitoneal cystic teratoma, caseating tubercular lymph nodes, and hydatid cysts. Even lymphoma and duplication cysts may also give similar appearances[2,3,]

Excision biopsy is then recommended to differentiate

these cases. Histopathology is confirmatory and differentiates chylolymphatic cysts from all these lesions. Cystic lymphangioma has a striking resemblance to chylolymphatic mesenteric cysts both grossly and microscopically. Some authors consider chylolymphatic mesenteric cysts to be a type of cystic lymphangioma, but the medical literature also shows some authors describing chylolymphatic cysts as a variant of mesenteric cysts [2,3,4,5]. The absence of smooth muscle and lymphatic spaces in the wall of the cyst differentiates mesenteric cysts from cystic lymphangioma.

Conclusion:-


Cysts of the mesentery are among surgical rarities. [1,2,3] In most of the cases the diagnosis is confirmed after surgical exploration and removal of the cyst. We would like to emphasize the importance of successful enucleation of the cyst irrespective of its size due to its independent blood supply as opposed to enterogenous cyst which requires bowel resection and anastomosis

Although very rare, chylolymphatic mesenteric cyst should be kept in mind as one of the differential diagnoses of cystic masses of the abdomen including cystic lymphangioma. [1,2,4,5,6,] Ultrasonography and computed tomography suggest the diagnosis but histopathological examination is required for confirmation. Complete excision of the cyst yields excellent results.

Figure 1. Gross : Appendix showing cyst attached to the wall on either sides



Figure 2) Microphotograph showing cysts lined with endothelium and filled with chylous fluid and lymph.. (H & E stain , 10 x).



REFERENCE

- [1] Engel S, Clagett OT, Harrison EG Jr: Chylous cysts of the abdomen. *Surgery* 1961, 50:593-599. | [2] Gupta AR, Nanavati RN, Fernandez AR, Kalgutkar A, Nathani R, Deshmukh SS: Chylous mesenteric cyst: an unusual cause of neonatal intestinal obstruction. *Indian Pediatrics* 1992, 29(4):511-513. | [3] Takiff H, Calabria R, Yin L, Stabile BE: Mesenteric cysts and intra-abdominal cystic lymphangiomas. *Archives of Surgery* 1985, 120(11):1266-1269. | [4] Beahrs OH, Judd ES Jr, Dockerty MB: Chylous cysts of the abdomen. *The Surgical Clinics of North America* 1950, 30(4):1081-1096. | [5] Losanoff JE, Richman BW, El-Sherif A, Rider KD, Jones JW: Mesenteric cystic lymphangioma. *Journal of American College of Surgeons* 2003, 196(4):598-603. | [6] Hebra A, Brown MF, McGeehin KM, Ross AJ: Mesenteric, omental, and retroperitoneal cysts in children: a clinical study of 22 cases. *South Medical Journal* 1993, 86(2):173-176. | [7] Kurtz RJ, Heimann TM, Holt J, Beck AR: Mesenteric and retroperitoneal cysts. *Annals of Surgery* 1986, 203(1):109-112. | [8] Bliss DP Jr, Coffin CM, Bower RJ, Stockmann PT, Ternberg JL: Mesenteric cysts in children. *Surgery* 1994, 115(5):517-577. |