Surgery



Treatment by Percutaneous Catheter Drainage in Hepatic Abscess Management-A Study

KEYWORDS	percutaneous catheter drainage, hypochondrium, hepatic abscess, conservative	
Dr. Sohail Anjum		Dr. Apurva kumar Rajan
Senior resident Department of surgery Jawaharlal Nehru Medical College, Bhagalpur, Bihar		Senior resident Department of surgery Jawaharlal Nehru Medical College, Bhagalpur, Biharc
ABSTRACT Aim and Objectives: To see the effectiveness of percutaneous catheter drainage (PCD) approach in the management of hepatic abscess. Materials & Methods: In this study, 60 cases of liver abscess were taken with age group between 25-60 years. Study was done in surgery department of JLNMCH, Bhagalpur, Bihar from October 2015 to September 2016. All cases were registered fulfilled the inclusion criteria and exclusion criteria. Group A was given conservative treatment and Group B was given percutaneous catheter drainage. Results were analysed sta-		

tistically. Results: it was shown in this study that mostly males were there, 55 out of 60 cases. The most common complain was abdominal pain and tenderness in right hypochondrium in 85% of the cases. The average hospital stay for patients in group B was 5 days as compared to 7 days in group A. Pain relieved in just 4 days in group B as compared to group A, it was 6 days. Conclusions: Percutaneous catheter drainage is a effective option of treatment in hepatic abscess as compared to conservative management

Introduction-

Liver abscess is most common in india. The patients of liver abscess are treated with antibiotics along with percutaneous needle aspiration (PNA) or percutaneous catheter drainage (PCD), with surgical drainage being used only in patients who fail to respond to such treatment. Though both PNA and PCD have been shown to be effective and safe, data comparing these procedures are limited. It usually occurs in the right lobe of the liver, and in very few cases in left lobe. Patients complain of fever, chills, pain in the right hypochondrium with tenderness. There may be weight loss and hepatomegaly. Percutaneous drainage combined with antimicrobial has become the first line and mainstay of treatment for most of the abscesses. It is suggested that percutaneous drainage (needle aspiration and catheter drainage) can improve response to antibiotic treatment, reduce hospital stay and the total cost of treatment. They are nonetheless an invasive procedure requiring the passage of a wide bore needle and pigtail catheter into a highly vascular organ, and can be associated with the risk of bleeding. Historically, morbidity and mortality rates associated with the treatment of pyogenic liver abscess has been high. This has improved significantly with the introduction of ultrasound (US) and computed tomography (CT). When detected early, effective treatment of small abscesses involves the use of diagnostic percutaneous aspiration and appropriate intravenous antibiotics. Liver abscesses larger than 5 cm in size often require prompt drainage for resolution of sepsis. PCD- A trocar with a 14 F multi-sidehole pigtail catheter was inserted into the abscess cavity under local anesthesia and US guidance. The contents of the abscess were aspirated and the catheter was fixed to the skin and connected to a bag. The first review US was done when drainage over the last 24 hours had declined to <10 mL. If the abscess had resolved, the catheter was removed. If residual cavity was present, the catheter was flushed with saline and aspirated till no material was aspirated.

Materials and methods

In this study, 60 cases of liver abscess were taken with age group between 25-60 years. Study was done in surgery department of JLNMCH, Bhagalpur, Bihar from Octo-

ber 2015 to September 2016. All cases were registered fulfilled the inclusion criteria and exclusion criteria. Group A was given conservative treatment and Group B was given percutaneous catheter drainage. Routine blood investigations were done. X-ray chest and abdomen with ultrasonography was done to diagnose liver abscess. Intravenous antibiotics in form of cephalosporins or fluroquinolones and metronidazole were given to all patients. PCD was done in Group B patients. Results were analysed statistically.

Results

In this study, it was shown that mostly males were there, 55 out of 60 cases. The most common complain was abdominal pain and tenderness in right hypochondrium in 85% of the cases. Mean duration of intravenous antibiotic and duration to clinical relief were shorter in the PCD group than in group with conservative approach. The average hospital stay for patients in group B was 5 days as compared to 7 days in group A. Pain relieved in just 4 days in group B as compared to group A, it was 6 days.

Discussion

Liver abscesses, both amebic and pyogenic, continue to be an important cause of morbidity and mortality in the tropical countries. Patients usually present late when the liver abscess attains a large size. The introduction of the percutaneous approach to liver abscesses was initially described by MaFadzean et al12 from Hong Kong in 1953. The body of literature in recent years suggests that PD of liver abscess is a safe and effective method for drainage of pyogenic liver abscess. As a result, this modality has become viewed as the treatment of choice, being superior to SD. The attraction of PD is related to its minimally invasive approach and the ability to perform the procedure without general anesthesia. Selected series of PD drainage report success rates of more than 95%. However, PD is not suitable for all patients with liver abscesses. In cases of rupture, difficult access to the abscess due to anatomic location, coexisting pathology requiring open surgery, SD becomes the only treatment option. In addition, the abscesses that fail to resolve after PD will also have to rely on an open surgical approach for successful management. Although the trend is toward PD, it is difficult to assess the true merits of each procedure to say which modality produces better clinical outcome. Singh and Kashyap also reported faster and more complete resolution of abscess cavities after PCD, while Rajak et al5 found that the time needed for total resolution is similar after PCD and PNA. Percutaneous drainage (either needle aspiration or catheter drainage) with systemic antibiotics has become the preferred treatment for the management of pyogenic liver contrast, for amoebic abscesses, the primary mode of treatment is medical. a preferred method most widely us liver abscesses, recent studies have shown PNA to be simpler, less costly, and equally effective. Usually needle aspiration is preferred for smaller abscesses and catheter drainage is done in larger ones. But no clear cut guidelines have been laid.

Conclusion

Percutaneous catheter drainage is a effective option of treatment in hepatic abscess as compared to conservative management

References

- Rajak CL, Gupta S, Jain S, Chawla Y, Gulati M, Suri S; Percutaneous treatment of liver abscess- needle aspiration versus catheter drainage. Am J roentgenology,(1998; 170(4): 1035-1039)
- Gerzof SG, Johnson WC, Robbins AH, Nabseth DC; Intrahepatic pyogenic abscesses: treatment by percutaneousdrainage. Am J Surg, (198; 149: 487494)
- Tandon A, Jain AK, Dixit VK, Agarwal AK, Gupta JP; Needle aspiration in large amoebic liver abscess. Trop Gastroenterol, (1997; 18: 19-21)
- Ch Yu S, Hg Lo R, Kan PS, Metreweli C; Pyogenic liver abscess: treatment with needle aspiration. ClinRadiol, (1997; 52: 912-916)
- Neoptolemos JP, Macpherson DS, Holm J, et al. Pyogenic liver abscess: a study of forty-four cases in two centres. Acta Chir Scand. (1982;148: 415–421)
- Northover JM, Jones BJ, Dawson JL, et al. Difficulties in the diagnosis and management of pyogenic liver abscess. Br J Surg. (1982;69:48–51)
- Miedema BW, Dineen P. The diagnosis and treatment of pyogenic liver abscesses. Ann Surg. (1984;200:328–335)
- Silver S, Weinstein AJ, Cooperman A. Changes in the pathogenesis and detection of intraphepatic abscess. Am J Surg. (1979;137:608–610)
- Herbert DA, Fogel DA, Rothman J, et al. Pyogenic liver abscesses: successful non-surgical therapy. Lancet. (1982;1:134–136)
- Bertel C K, van Heerden J A, Sheedy P Treatment of pyogenic hepatic abscesses: Surgical vs percutaneous drainage. Arch Surg. (558)
- Herman P, Pugliese V, Montagnini A et al. Pyogenic liver abscess: The surgical treatment. Int Surg. (98-10)
- Huang CJ, Pitt HA, Lipsett PA, et al. Pyogenic hepatic abscess: changing trends over 42 years. Ann Surg. (1996;223:600–607)
- Attar B, Levendoglu H, Cuasay NS. CT-guided percutaneous aspiration and catheter drainage of pyogenic liver abscesses. Am J Gastroenterol. (1986;81:550–555)
- Bertel CK, van Heerden JA, Sheedy PF 2nd. Treatment of pyogenic hepatic abscesses: surgical vs percutaneous drainage. Arch Surg. (1986; 121:554–558. 12)
- McFadzean AJS, Chang KPS, Wong CC. Solitary pyogenic abscess treated by closed aspiration and antibiotics: fourteen consecutive cases with recovery. Br J Surg. (1953;41:141–152)
- McDonald MI, Corey GR, Gallis HA, et al. Single and multiple pyogenic liver abscesses: natural history, diagnosis and treatment, with emphasis on percutaneous drainage. Medicine (Baltimore). (1984;63: 291–302)
- Miller FJ, Ahola DT, Bretzman PA, et al. Percutaneous management of hepatic abscess: a perspective by interventional radiologists. J Vasc Interv Radiol. (1997;8:241–247)
- Johnson RD, Mueller PR, Ferrucci JT Jr, et al. Percutaneous drainage of pyogenic liver abscesses. AJR Am J Roentgenol. (1985;144:463–467)