



“Evaluation of dynamic pulmonary function test in overweight under graduate medical students”

KEY WORDS

Overweight, BMI & PFT.

Dr Somanath Naik

MBBS, MD Associate professor, Department of Physiology Hi-Tech Medical College and Hospital, Rourkela, Odisha, India

ABSTRACT

Overweight and obesity has become a major health problem in India. My aim was to determine and compare the possible alterations of pulmonary function tests among healthy controls and overweight individuals irrespective of gender. These findings serve as a reaffirmation, for the proven fact that increasing BMI leads to defective lung functions. All the four parameters were found to have statistical significance, which proves overweight to be a major risk factor for defective lung functions.

Introduction:

Overweight and obesity has become a major health problem in India. In the recent few years sedentary life style and dietary habits had played an important role in the occurrence of overweight and obesity. Obesity can profoundly alter pulmonary function and diminish exercise capacity by its adverse effects on respiratory mechanics, resistance within the respiratory system, respiratory muscle function, lung volumes, control of breathing and gas exchange.¹⁻⁹ My aim was to determine and compare FVC, FEV1, FEV1/FVC and PEFR among controls and overweight individuals irrespective of gender.

Material and Methods:

The present study was conducted in the Department of physiology, Hi-Tech Medical College & Hospital Rourkela, Odisha, India, during the period from October 2015 to March 2016. The study protocol was approved by the Ethics committee of HMCH, Rourkela. The present study consists of total 50 subjects between the age group 18-26 years who are further subdivided into two groups;

- Group-A** : Includes total 25 healthy individual as controls.
- Group-B** : Consists of 25 overweight as cases. Subjects were selected based on the following inclusion and exclusion criteria.

Inclusion criteria:

- Subjects between 18 to 26 years of age including both males and females
- Subjects with willingness to participate in the study
- Subjects who falls in the category of normal and overweight according to BMI.

Exclusion criteria:

- History of smoking, alcohol intake, occupational hazards, medical(asthmatic) or surgical illness, upper respiratory tract infection in the past 4 weeks
- Physical examination findings suggestive of respiratory illness or skeletal deformity
- Intake of drugs which influences lipid metabolism, upper abdominal surgeries, Diabetes Mellitus, Cardiovascular Disorders, Endocrinology Disorders. Then subjects were made to undergo pulmonary function test using RMS Computerised (Helios 401) Spirometer.

All values were expressed as mean±sd. We used student t-test to find the statistical significance. A P-value <0.05 was to be considered statistically significant.

Results and Discussion:

The present study group consisted of 50 subjects between 18- 26 years of age of both genders. Among which 25(14M & 11F) were controls (normal weight) and 25(13M & 12F) were cases (overweight).




Fig-1 shows the Anthropometric Characteristics of Subjects in the Cases and Control Groups :

When compared between normal and overweight individuals, significant difference is observed in height, weight and BMI with p-values <0.05 and the mean+sd for age, they did not show statistically significant with a p value (0.14 >0.05). The spirometry parameters of the Cases and Control groups were compared in Table-2 When compared between Cases and Control individuals, significant difference is observed in all the four parameters (Forced vital capacity (FVC), forced expiratory volume in first second (FEV1), %FEV1, peak expiratory flow rate (PEFR)) as shown in table 2; with p value <0.05. In this study, significant difference is observed in all the four parameters (FVC, FEV1, %FEV1 & PEFR), which were used to compare the lung function in overweight and normal healthy individuals. Similar results has been observed in a study conducted by researchers during 2008, which showed a significant decrease in FVC, FEV1 and %FEV1 with p value <0.05. And these two factors are considered as the most common lung functions inversely related to BMI.¹⁰ A study conducted by Rubinstein also proved the same finding as mentioned above.¹¹

Table-2 shows the Pulmonary functions test of the Cases and Control Groups :

Parameters	Group B Mean±SD	Group A Mean±SD	P-Value
FVC	2.95 ± 0.12	3.22 ± 0.5	0.01
FEV1	2.34 ± 0.46	2.82 ± 0.24	0.01
%FEV1	0.76 ± 0.16	0.84 ± 0.02	0.01
PEFR	6.16 ± 2.43	7.2 ± 2.32	0.01

*Statistically Significant (P<0.05)

A recent research conducted by Arkanshu et al in Chennai during 2014 demonstrated the negative correlation existing between BMI and FEV1/FVC ratio.¹² And this study results supports our study findings with significant statistical difference. Our result is also supported by another study conducted by Donna Rinnie during 2007, which also has proved the inverse relationship between FEV1/FVC

ratio and BMI in overweight and obese individuals.¹³ Many other researchers including Lazarus et al¹⁴, Biring et al¹⁵ and Paralikar et al¹⁶ have also shown decrease in FEV1/FVC in overweight and proved the strong negative correlation existing between FEV1/FVC and BMI. Even though many studies supported the inverse relationship between FVC, FEV1 & FEV1/FVC ratio, research conducted by Emel et al had shown no statistical differences in FEV1, FVC, or FEV1/FVC ratio. In the same research conducted by Emel et al, PEF is reduced in overweight with significant statistical difference supporting our result.¹⁷ Results of another study conducted by Mohammed also had reduction in PEFR, in accordance with our study result. But there was no significant reduction in other three parameters (FVC, FEV1 and FEV1/FVC).¹⁸

Reason for reduction in these parameters of lung functions may be due to deposition of fat around the chest wall leading to decrease in the compliance of chest wall¹³ and lowered respiratory muscle endurance with increased work of breathing, airway resistance¹⁹ and gas transport.²⁰ Defective descent of diaphragm due to mechanical hindrance caused by deposition of fat around the chest wall also plays an important role in altered lung functions.^{21,22} Another reason has also been reported by many researchers, mentioning the role of adipose tissue in altered lung functions. Adipocytokines are released by adipose tissue as a result of hypoxemia induced by obesity. These adipocytokines affects systemic inflammation, leading to increase in inflammatory markers. Increased level of interleukins (ILs) 6 and 8, tumor necrosis factor α (TNF- α), CRP, leptin, and lower levels of adiponectin, which helps in regulating insulin sensitivity have all observed in individuals with increased BMIA.^{23,24} All these factors act mutually in altering the lung functions with increased BMI above normal range. In this study, many of the individuals in overweight group have decreased lung function when compared to the normal healthy group.

Conclusion:

In conclusion, these findings serve as a reaffirmation, for the proven fact that increasing BMI leads to defective lung functions. All the four parameters were found to have statistical significance, which proves overweight to be a major risk factor for defective lung functions. Therefore, factors which influence the increase in BMI like genetics, ethnicity, nutrition, physical activity and lifestyle can also influence the pulmonary functions.

REFERENCES

- Rio-Navarro BE, Cisneros-Rivero MG, Berber-Eslava A, Espinola-Reyna G, Sienra-Monge JLL. Exercise induced bronchospasm in asthmatic and non-asthmatic obese children. *Allergol et Immunopathol* 2000; 28: 5-11.
- Strauss R. Childhood obesity. *Curr Probl Pediatr* 1999; 29: 5-29.
- Inselma LS, Milanese A, Deurloo A. Effect of obesity on pulmonary function in children. *Pediatric Pulmonol* 1993; 16: 130-7.
- Koenig SM. Pulmonary complications of obesity. *Am J Med Sci* 2001; 321: 249-79.
- Valente AM, Strong W, Sinaiko AR. Obesity and insulin resistance in young people. *Am Heart J* 2001; 142: 440-4.
- Jones RL, Nzekwu MM. The effects of body mass index on lung volumes. *Chest* 2006; 130: 827-33.
- Parameswaran K, Todd DC, Soth M. Altered respiratory physiology in obesity. *Can Respir J* 2006; 13: 203-10.
- Ulger Z, Demir E, Tanac R, Goksen D, Gulen F, darcan S, Can D, Coker M. The effect of childhood obesity on respiratory function tests and airway hyperresponsiveness. *Turk J Pediatr* 2006; 48: 43-50.
- Li AM, Chan D, Wong E, Yin J, Nelson EA, Fok TF. The effects of obesity on pulmonary function. *Arch Dis Child* 2003; 88: 361-3.
- Anuradha R, Joshi, Ratan Singh and A. R. Joshi. Correlation of pulmonary function tests with body fat percentage in young individuals. *Indian J Physiol Pharmacol* 52 (4) : 383-388. (2008).
- Rubinstein L, Zamel N, DuBarry L, Hoffstein V. Airflow limitation in morbidly obese nonsmoking men. *Ann Intern Med*, 112:828-32, (1990).
- Arkanshu A, Sanjay A. Relationship between Certain Pulmonary Function Tests and Anthropometric Measures in Normal Male M.B.B.S Students. *International Journal of Health Information and Medical Research*, 1(3), (Apr - Jun 2014).
- Yue Chen., Donna Rennie., Yvon F Cormier., and James Dosman. Waist circumference is associated with pulmonary function in normal-weight, over- weight, and obese subjects. *Am J Clin Nutr*;85:35-9, (2007).
- Lazarus R, Sparrow D, Weiss ST. Effects of obesity and fat distribution on ventilatory function. *Chest*. 111:891-8, (1997).
- Biring MS., Lewis MI., Liu JT., Mohsenifar Z. Pulmonary physiologic changes of morbid obesity. *Am J Med Sci*. 318:293-7, (1999).
- Paralikar SJ, Kathrotia RG., Pathak NR., Jani MB. Assessment of pulmonary functions

in obese adolescent boys. *Lung India*, 29:236-40, (2012).

- Turkey Emel Torun., Erkan Cakir., Fatma Özgüç., Iker Tolga Özgen. The Effect of Obesity Degree on Childhood Pulmonary Function Tests. *Balkan Med J*, 31(3), (2014).
- Al Ghobain M. The effect of obesity on spirometry tests among healthy non-smoking adults. *BMC Pulmonary Medicine*, 12:10, (2012).
- McClellan KM., Kee F., Young IS., Elborn JS. Obesity and the lung: 1. *Epidemiology. Thorax*. 63:649-654, (2008).
- Santiago SQ, Silva ML., Davidson J., Aristóteles LR. Evaluation of respiratory muscle strength in overweight/obese children and adolescents. *Rev Paul Pediatr* 26:146-50, (2008).
- Lazarus R., Gore CJ., Booth M., et al. Effects of body composition and fat distribution on ventilatory function in adults. *Am J Clin Nutr* 68:35-41, (1998).
- Obesity and respiratory diseases. *International Journal of General Medicine*, 3/20, (oct 2010).
- Leone N, Courbon D, Thomas F, et al. Lung function impairment and metabolic syndrome. The critical role of abdominal obesity. *Am J Respir Crit Care Med*, 179:509-516, (2009).
- Franssen FME, O'Donnell DE, Blaak EE, Schols AMWJ. Obesity and the lung: 5. Obesity and COPD. *Thorax*, 63:1110-1117, (2008).