



A study of anatomy of middle cluneal nerve entrapment

Dr. Shruti Mamidwar

Associate Professor, Department of Anatomy, GMC, Chandrapur.

Dr. Abhilasha M

Vahane, Associate Professor, Department of Anatomy, GMC, Nagpur.

ABSTRACT

The purpose of this study was to ascertain, using cadavers, the relationship between the MCN and LPSL and to investigate MCN entrapment.

Methodology: A total of 30 hemipelves from 20 cadaveric donors (15 female, 5 male) designated for education or research, were studied by gross anatomical dissection. The age range of the donors at death was 71–101 years with a mean of 88 years.

Results: A total of 64 MCN branches were identified in the 30 hemipelves. Of 64 branches, 10 (16%) penetrated the LPSL. The average cephalocaudal distance from the PSIS to where the MCN penetrated the LPSL was 28.5 ± 11.2 mm (9.1–53.7 mm). The distance from the midline was 36.0 ± 6.4 mm (23.5–45.2 mm). The diameter of the MCN branch traversing the LPSL averaged 1.6 ± 0.5 mm (0.5–3.1 mm). Four of the 10 branches penetrating the LPSL had obvious constriction under the ligament.

KEYWORDS : middle cluneal nerve, sacroiliac joint.

Introduction:

In 1957, Strong and Davila attempted deafferentation of the superior cluneal nerve (SCN) and/or middle cluneal nerve (MCN) in 30 LBP patients. 1 Five of these 30 patients had referred pain in a leg in the S1 or S2 area; deafferentation of the MCN yielded favorable outcomes. Strong and Davila stated the MCNs were thin and difficult to identify during surgery, but did not describe the relationship between the MCN and long posterior sacroiliac ligament (LPSL).

Following anatomical reports by Maigne et al² and by Lu et al³ that described entrapment of the most medial branch of the SCN where the nerve passes through the fascia over the iliac crest, 4 successful surgical techniques were developed to open the fascial orifice for relief of this entrapment neuropathy.^{5–9} Trescot¹⁰ and Kuniya et al¹¹ stated that cluneal neuralgia is not a rare clinical entity and may be underdiagnosed and should be considered as a differential diagnosis for chronic LBP or leg pain.

No reports of MCN entrapment have been available until a recent case report that described severe LBP completely alleviated by release of the MCN.⁴ In this case, the MCN was entrapped where this nerve passed under the LPSL.

The MCN comprises sensory branches of the dorsal rami of S1–S3 foramina. It travels below the posterior superior iliac spine (PSIS) in an approximately horizontal course to supply the skin overlying the posteromedial area of the buttock.^{12–14} Controversy exists regarding a relationship between the MCN and LPSL. Tubbs et al¹⁴ reported that the MCN would be less likely to become entrapped because the MCN travels superficially to the LPSL. However, Horwitz,¹⁵ Grob et al,¹² and McGrath and Zhang¹⁶ reported that the primary and secondary loops of the posterior sacral nerve plexus passed through or beneath the LPSL. In view of the paucity of literature on this subject, we performed an anatomical study of the MCN around the LPSL with the objective of providing an accurate anatomical basis for clinical conditions involving entrapment of the nerve.

Materials and methods:

This anatomical study was conducted in the Department of Anatomy of a Medical College in Central India. A total of 40 usable hemipelves were obtained from 20 formalin-preserved Japanese cadavers (5 male and 15 female). The average age at death was 88 years and the age range was 71–101 years. All the cadavers were routinely fixed in formalin solution. Bilateral branches of the MCN were macroscopically explored. None of the cadavers showed evidence of previous surgical procedures or traumatic lesions to the pelvis. Cadavers were placed in the prone position. Branches of the MCN were identified under or over the gluteus maximus fascia on the caudal side of the PSIS and traced laterally as far as the finest visible ramification. Special attention was paid to the relationship between the

MCN and LPSL. Because lateral branches of the dorsal L5–S4 rami anastomose to form loops dorsal to the sacrum, with each branch containing nerve fibers from adjacent dorsal rami,^{15–17} it was impossible to trace them individually. Therefore, the major dorsal sacral rami were dissected and traced medially to the dorsal sacral foramina to identify the level of origin. MCN branches were counted where they traversed over and under the LPSL.

Results:

A complete exploration of the MCN failed in the initial 10 hemipelves, therefore, data from the remaining 30 hemipelves were analyzed for this study. A total of 64 MCN branches were identified in these 30 hemipelves (Table 1). MCN branches were composed of S1–S4 dorsal rami. The distances from these anatomical landmarks are shown in Table 2 relative to origin. The distances from the PSIS to dorsal rami traversing over or under the LPSL were ~20 mm for S1, 23 mm for S2, 34 mm for S3, and 41 mm for S4 (Table 2).

Discussion:

The LPSL is a significant posterior SIJ ligamentous structure that resists shearing of the SIJ.^{17,18} SIJ pain has been a controversial and ill-defined subject. SIJ disorders have an imprecise etiology and are thought to cause 15%–30% of LBP and are often associated with buttock to lower extremity symptoms.¹⁹ There are no medical history, physical examination, or radiological findings consistently capable of identifying SIJ pain.²⁰ The current gold standard for diagnosis of SIJ pain is fluoroscopically guided SIJ blocks.¹⁹ Radiofrequency ablation or blocking of the lateral branches of the dorsal sacral rami that supply the SIJ is a treatment option gaining considerable attention.^{21,22}

Several researchers consider the LPSL to be a major pain generator of SIJ pain.^{16,18,23–25} Fortin and Falco²⁰ stated that SIJ patients could localize their pain with one finger and the area pointed to was within 1 cm inferomedial to the PSIS. Murakami et al²⁴ observed positive effects from a periarticular SIJ block in 18 of 25 patients who located the primary site of their pain to within 2 cm of the PSIS. Murakami et al²⁵ compared the effect of blocking injections into the intraarticular space and around the LPSL in patients fulfilling definite criteria for SIJ pain. Blocking injections around the LPSL were effective in all 25 patients, whereas intraarticular blocking injections were effective in only 9 out of 25 patients (36%). In addition, all 16 patients without pain relief after an intraarticular blocking injection reported almost complete pain relief after a blocking injection around the LPSL. In a recent anatomical report by Cox and Fortin,²¹ which attempted to clarify innervation of the SIJ by the lateral branches, the authors stated that the most lateral portion of the lateral branch of S1 was traced after it passed through a fibro-osseous tunnel in the LPSL.²¹ In our study, 10 of 64 MCN branches passed under the LPSL. It is likely that blocks around the LPSL may infiltrate around the dorsal sacral rami passing over or under the LPSL.

Tables:

TABLE 1: Spinal levels of sacral nerve roots originating MCN branches.

Specimen no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
S1			○	○	○	○																									△
S2		●					●														△		□	□	□	□	□	□	□	□	△
S3		□	□																		△	□								△	△
S4																															

Notes: ○, Cephalad branch; △, Middle branch; □, Caudal branch. Under bar () represents branch passing under LPSL, Black marks represent a branch with macroscopic indentation by the LPSL.


Abbreviations: MCN, middle cluneal nerve; LPSL, long posterior sacroiliac ligament.

Table 2: Measurements of MCN branches relative to origin.

Nerve roots	Distance from midline (mean±SD) (mm)	Distance from PSIS (mean±SD) (mm)	Diameter of MCN (mean±SD) (mm)
S1 (n=15)	33.1±7.2	20.7±5.0	1.7±0.6
S2 (n=27)	37.0±6.1	23.3±7.9	1.7±0.6
S3 (n=17)	37.9±4.9	33.9±9.3	1.4±0.3
S4 (n=12)	34.6±6.7	41.0±10.4	1.3±0.3

Abbreviations: MCN, middle cluneal nerve; PSIS, posterior superior iliac spine; SD, standard deviation

FIGURE 1: Schematic illustration of measurements of linear distances from the posterior superior iliac spine (distance a) and midline (distance b) to a branch of the MCN traversing over or under the LPSL.



References:

- Strong EK, Davila JC. The cluneal nerve syndrome; a distinct type of low back pain. *Ind Med Surg.* 1957;26(9):417–429.
- Maigne JY, Lazareth JP, Guerin Surville H, Maigne R. The lateral cutaneous branches of the dorsal rami of the thoraco-lumbar junction. An anatomical study on 37 dissections. *Surg Radiol Anat.* 1989;11:289–293.
- Lu J, Ebraheim NA, Hunton M, Heck BE, Yeasting RA. Anatomic considerations of superior cluneal nerve at posterior iliac crest region. *Clin Orthop Relat Res.* 1998;(347):224–228.
- Aota Y. Entrapment of middle cluneal nerves as an unknown cause of low back pain. *World J Orthop.* 2016;7(3):167–170.
- Berthelot JM, Delecrin J, Maugars Y, Caillon F, Prost A. A potentially underrecognized and treatable cause of chronic back pain: entrapment neuropathy of the cluneal nerves. *J Rheumatol.* 1996;23(12):2179–2181.
- Kim K, Isu T, Chiba Y, et al. The usefulness of ICG video angiography in the surgical treatment of superior cluneal nerve entrapment neuropathy: technical note. *J Neurosurg Spine.* 2013;19(5):624–628.
- Maigne JY, Doursounian L. Entrapment neuropathy of the medial superior cluneal nerve. Nineteen cases surgically treated, with a minimum of 2 years' follow-up. *Spine (Phila Pa 1976)* 1997;22(10):1156–1159.
- Morimoto D, Isu T, Kim K, et al. Surgical treatment of superior cluneal nerve entrapment neuropathy. *J Neurosurg Spine.* 2013;19(1):71–75.
- Speed S, Sims K, Weirauch P. Entrapment of the medial branch of the superior cluneal nerve a previously unrecognized cause of lower back pain in cricket fast bowlers. *J Med Cases.* 2001;2:101–103.
- Trescot AM. Cryoanalgesia in interventional pain management. *Pain Physician.* 2003;6(3):345–360.

- Kuniya H, Aota Y, Kawai T, Kaneko K, Konno T, Saito T. Prospective study of superior cluneal nerve disorder as a potential cause of low back pain and leg symptoms. *J Orthop Surg Res.* 2014;9:139.
- Grob KR, Neuhuber WL, Kissling RO. Die innervation des sacroiliaclenkeim beim menschen. [Innervation of the sacroiliac joint of the human] *Z Rheumatol.* 1995;54(2):117–122.
- Sittitavornwong S, Falconer DS, Shah R, Brown N, Tubbs RS. Anatomic considerations for posterior iliac crest bone procurement. *J Oral Maxillofac Surg.* 2013;71(10):1777–1788.
- Tubbs RS, Levin MR, Loukas M, Potts EA, Cohen-Gadol AA. Anatomy and landmarks for the superior and middle cluneal nerves: application to posterior iliac crest harvest and entrapment syndromes. *J Neurosurg Spine.* 2010;13(3):356–359.
- Horwitz TM. The anatomy of (A) the lumbosacral nerve plexus – its relation to variations of vertebral segmentation, and (B) the posterior sacral nerve plexus. *Anat Rec.* 1939;74(1):91–107.
- McGrath MC, Zhang M. Lateral branches of dorsal sacral nerve plexus and the long posterior sacroiliac ligament. *Surg Radiol Anat.* 2005;27(4):327–330.
- Standring S, editor. *Gray's Anatomy The Anatomical Basis of Medicine and Surgery.* 38th ed. Edinburgh: Churchill Livingstone; 2008.
- Vleeming A, Pool-Goudzwaard AL, Hammudoghlu D, Stoelckart R, Snijders CJ, Mens JM. The function of the long dorsal sacroiliac ligament: its implication for understanding low back pain. *Spine (Phila Pa 1976)* 1996;21(5):556–562.
- Vanelderen P, Szadek K, Cohen SP, et al. 13. Sacroiliac joint pain. *Pain Pract.* 2010;10(5):470–478.
- Fortin JD, Falco FJ. The Fortin finger test: an indicator of sacroiliac pain. *Am J Orthop (Belle Mead NJ)* 1997;26(7):477–480.
- Cox RC, Fortin JD. The anatomy of the lateral branches of the sacral dorsal rami: implications for radiofrequency ablation. *Pain Physician.* 2014;17(5):459–464.
- Fortin JD, Washington WJ, Falco FJ. Three pathways between the sacroiliac joint and neural structures. *AJNR Am J Neuroradiol.* 1999;20(8):1429–1434.
- Cusi M, Van der Wall H, Saunders J, Wong L, Pearson M, Fogelman I. Sacroiliac steroid injections do not predict ablation relief-not a surprise. *Pain Med.* 2013;14(1):163–164.
- Murakami E, Aizawa T, Noguchi K, Kanno H, Okuno H, Uozumi H. Diagram specific to sacroiliac joint pain site indicated by one-finger test. *J Orthop Sci.* 2008;13(6):492–497.
- Murakami E, Tanaka Y, Aizawa T, Ishizuka M, Kokubun S. Effect of periarticular and intraarticular lidocaine injections for sacroiliac joint pain: prospective comparative study. *J Orthop Sci.* 2007;12(3):274–280.
- Beith ID, Kemp A, Kenyon J, Prout M, Chestnut TJ. Identifying neuropathic back and leg pain: a cross-sectional study. *Pain.* 2011;152(7):1511–1516.