



PANCREATIC STUMP MANAGEMENT FOLLOWING PANCREATICO DUODENECTOMY- OUTCOME ANALYSIS OF VARIOUS RECONSTRUCTION TECHNIQUES

Prabhakaran Raju

KEYWORDS :

Background:

Pancreatic stump anastomosis is the Achilles heel after Whipple's Procedure. The morbidity (40-60%) and mortality (1-5%) following Whipple Procedure is related to the outcome of anastomosis. The aftermath of a pancreatic leak can be devastating, particularly when it results in retroperitoneal sepsis. This is found to be a major cause of mortality in Whipple's procedure [1]. Effects to improve the anastomotic techniques and thereby outcome of Whipple procedure is still evolving. Though many randomized and prospective studies are available till date no simple best technique had been recommended.

Aim of the study:

To analyse the outcome of pancreatic stump anastomosis of various types in relation to major and minor morbidities and mortality in relation to individual type of anastomosis.

Materials and methods:

Retrospective analysis of prospectively collected data from 2010 to 2014 March on patients underwent Whipple procedure done – 138 patients have undergone Whipple procedure. Preoperative, Intraoperative and postoperative variables were taken for this study. All patients admitted with a diagnosis of periampullary carcinoma or carcinoma head of pancreas were evaluated by imaging studies and those patients found to have resectable disease were selected for study. All data were collected prospectively and the clinical parameters were noted in a proforma. Details noted included age, gender, chief complaints, co-morbid illness, nature of diet, habit of smoking and alcohol consumption were also noted. Findings on physical examination such as jaundice, pallor, pedal edema and other signs of liver failure if present were noted. Clinical examination of the abdomen was done to look for a palpable gallbladder, hepatomegaly and free fluid. Rectal examination to rule out any possibility of rectal deposits. Basic biochemical and hematologic investigations including a complete blood count, Renal function tests and Liver function tests were noted. Coagulation profile and serum tumour marker study was done for all patients. After an initial ultrasonogram of abdomen, an upper GI endoscopy and contrast enhanced computerised tomography was done for all patients.

Reconstruction pancreaticoenteric anastomosis was done either in the form of a pancreaticogastrostomy, pancreaticojejunostomy or isolated loop pancreaticojejunostomy as per the choice of operating surgeon. Patients underwent Pancreatic stump anastomosis have been categorised into three groups.

- A- Pancreatico Gastrostomy (PG)
- B- Pancreatico Jejunostomy (PJ)
- C- Isolated Pancreatico jejunostomy (IPJ)

C group later categorised into Dunking type (C1) and Duct to mucosa (C2) type. Major complications like leak (Major/Minor), Hemorrhage (Early/late), DGE (Primary and secondary), Intra-abdominal abscess have been taken in relation to anastomotic techniques. Minor morbidities like Pneumonitis, UTI, wound infection also taken into account. Mortality also related to type of anastomosis.

Techniques used in Pancreatic stump management:




As long as the basic rules of a safe anastomosis are followed, including careful handling of the pancreatic tissues, a tension-free approximation, ensuring good blood supply, and no distal obstruction, any pancreaticoenteric anastomotic technique can have a good outcome. One of the most commonly employed techniques is a

pancreaticojejunal anastomosis. This anastomosis is done by invaginating the transected pancreas into the end of the jejunum, also known as dunking method; another variation is to anastomose the pancreatic duct directly to an opening in the jejunum, called the duct-to-mucosa technique. Another technique is to anastomose the pancreatic stump to the stomach by invagination to the posterior wall of stomach.

A 40-cm long isolated loop of jejunum is fashioned and passed in the retrocolic plane through the mesocolon for pancreaticojejunal anastomosis. The anastomosis is done by a duct to mucosa technique or a dunking technique using 3.0/4.0 prolene interrupted sutures for the anastomosis.

Isolated Loop PJ

Single loop PJ



Pancreatico gastrostomy



pancreaticojejunosomy

Statistical analysis:

The data collected in the proforma were entered in an excel sheet of Microsoft Office software and inference obtained after statistical analysis. The mean and standard deviation were reported for continuous variables and for categorical variables proportions were computed. To compare and find the statistical significance between the two group proportions chi square test was used and to compare between the two group means independent t-test was used. The P-values <0.05 were considered to indicate statistical significance.

Results

Among the one hundred and thirty eight patients included in the study 62% were male and 38% were female patients. The minimum age was 30 and maximum age was 72 with a mean age of 51.7. On clinical presentation 90% had jaundice, 86% had abdominal pain, 84% had weight loss, 56% had pruritus, 11% had fever, 12% had cholangitis and 28% had other symptoms such as nausea, vomiting, loss of appetite and constipation.

On examination, 81.15% were icteric and 27.53% had pallor. Gallbladder was palpable in 71.01% of patients and liver was palpable

in 40.57% of patients. Liver echoes were found to be normal in 92% of patients. Intrahepatic biliary radical dilatation was found in 96% and Common bile duct was dilated in 92% of the patients.

Findings	Frequency	Percentage
Icterus	112	81.15
Pallor	38	27.53
Palpable gallbladder	98	71.01
Hepatomegally	56	40.57

Periampullary 102 (79.68%), Pancreatic cancer 15 (11.7%) Distal CBD growth 6 (6%) and duodenal growth 5 cases were analysed.

Among them after pancreatico dudoenectomy- PG (A)-done for 40 cases. PJ(B)- done for 60 cases and Isolated PJ (C) done for 38 cases.

DGE(delayedgastricemptying) is the most common complication 44% (57). Overall complications include- pancreatic leak-30.96%, haemorrhage- 5.4%, Intra abdominal collection-5%.

Minor complications are 31% collectively.


When comparing major complications between the three groups undergoing pancreaticogastrostomy, pancreaticojejunostomy and isolated loop pancreaticojejunostomy, the incidence of delayed gastric emptying in the PG group was 38.46% , the incidence in the PJ group was 40.98% and in the isolated loop pancreaticojejunostomy group was 44.73%.

The incidence of haemorrhage was 7.6% in the PG group, 6.5% in the PJ group and nil in the isolated PJ group .

When comparing the incidence of leak between the three groups it was about 33% in the PG and 29.5% in the PJ group and 15.78% in isolated PJ group.

The incidence of intra abdominal collection in the PG group was 7[17.9%], in the PJ group it was7[11.4%] and in the isolated PJ group was 5 [13.15%].The mean duration of nasogastric tube removal was 7.5 days in the PG group and 7.8 days in the PJ group and 7.0 in Isolated PJ group. The mean postoperative hospital stay was 12.6 days in the PG group and 13.1 days in the PJ group and 11.2 in isolated PJ group. The mortality in the patients who underwent pancreaticogastrostomy was 5.1% ,in the pancreaticojejunostomy group was 4.9 %and 4.8 %in isolated loop PJ .The overall mortality rate was 5.79%.

Procedure	PG group (A)	PJ group (B)	Isolated PJ Group
Total number of surgeries	39	61	38
Haemorrhage	7.6%	6.5%	Nil
Pancreatic leak	33%	29.6%	15.78%
Delayed gastric emptying (DGE)	38.46%	29.5%	15.78%
Intra-abdominal collection	7 (17.95%)	7 (11.4%)	5 (13.5%)
Mortality	5.1%	4.9%	4.8%
Post op hospital stay	12.6 days	13.1 days	11.2 days



Discussion:

Mere occlusion of the duct has Mere occlusion of the duct has been shown to result in higher fistula rates, along with increasing the risk of pancreatic exocrine and endocrine insufficiency. Drainage of the pancreatic remnant to the gastrointestinal tract is an important step, but

it runs the risk of anastomotic breakdown.. Several techniques have been described, and the literature will continue to report novel techniques promising to be even safer.

An ideal reconstructive technique should not only minimize the risk of Pancreatic fistula formation, but should also ensure that, should a pancreatic fistula form, its complications are prevented or minimized. More than the choice of anastomotic technique, however, the successful management of the pancreatic anastomosis depends more on the surgeon’s meticulous execution of the technique with which he or she is familiar [2]

Proponents of the pancreaticogastrostomy cite various reasons[3] First, it is easier to perform, because of the close proximity of the stomach to the pancreas. Second, rich gastric blood supply makes this anastomosis less prone to ischemia. Third, because the exocrine enzymes encounter an acidic environment, the leak rate is theoretically lower as the enzymes do not get activated. The last statement has been disproved, however.

In a prospective randomized trial comparing pancreaticojejunostomy with pancreaticogastrostomy, the leak rates were not significantly different [pancreaticojejunostomy 11%; pancreaticogastrostomy 12%][4,5].Yeo et al has concluded that pancreatic fistula is a common complication after pancreaticoduodenectomy, with an incidence most strongly associated with surgical volume and underlying disease and the data do not support the hypothesis that pancreaticogastrostomy is safer than pancreaticojejunostomy or is associated with a lower incidence of pancreatic fistula.

In a meta analysis by Wente MN and Shrikande SV et al [6],they concluded that all non randomized observational clinical studies have reported superiority of pancreaticogastrostomy over pancreaticojejunostomy but all randomized controlled studies has shown equally good results. In a study by H Ramesh et al results suggested that pancreaticogastrostomy deserves wider application [7]. An isolated jejunal loop for Pancreatico enteric anastomosis is theoretically expected to achieve these desired endpoints. Previous studies, using an isolated jejunal loop for pancreatoenteric anastomosis can minimize the risk of Pancreatic Fistula, although its effect in terms of reducing pancreatic fistula related morbidity is not clear.[8-14] Advocates of this technique believe that diverting bile away from the pancreaticojejunostomy site minimizes the pancreatic enzyme activation and hence reduces the risk of pancreato enteric anastomotic fistula[15]

In another prospective randomized trial Bassi et al has showed that both type of anastomosis does not influence significantly the risk of overall complications or the incidence of pancreatic fistula. However, significant decreases in the risk of associated complications, biliary fistulas, postoperative collections and DGE were observed using pancreatico gastrostomy. A Chinese meta analysis of all four randomized controlled trials has evidence suggesting that pancreaticogastrostomy is better than pancreaticojejunostomy after pancreaticoduodenectomy.

In our study though we found no overall difference in the morbidities between the techniques, severity of complications is lesser with isolated loop technique like Grade A leak. Pancreatic leak occurred in 39 patients with grade A leak in 20(15.62%), grade B leak in 12(9.37%) and grade C leak in 7(5.46%) patients. All patients with pancreatic leak were managed by non-operative means. Grade A leaks were managed conservatively and grade B leaks required supportive care in the postoperative ward with drainage tube retained for a prolonged period and grade C leaks were managed aggressively in the ICU with one or more image guided percutaneous drainage tubes and nutritional support .We have not reoperated for a suspected leak. We also observed that it has demerits like long operating hours and increased incidence of DGE . In the subgroup analysis between Dunking method(C1)and Duct to mucosa(C2) anastomosis technique there is no difference between the techniques. Mortality is comparatively lesser than other methods but it has no statistical difference.

Conclusion:

Among various techniques of pancreas stump reconstruction (PG/PJ /Isolated PJ) none of them showed statistical significant morbidity or mortality of the existing standard. But isolated loop PJ has had statistically significant lower grade leak and increased DGE.

Subgroup analysis within thepancreaticojejunostomy has no difference in outcome. Pancreatic stump management has to be individualised. Surgeon should be familiar with all techniques.

References:

1. Berberat, 1999b. Berberat PO, et al: Prevention and treatment of complications in pancreatic cancer surgery. *Dig Surg* 1999; 16: 327-336.
2. Trede, 2001. Trede M, et al: Personal observations, opinions, and approaches to cancer of the pancreas and periampullary area. *Surg Clin North Am* 2001; 81:595-610.
3. Zenilman, 2000. Zenilman ME: Use of pancreaticogastrostomy for pancreatic reconstruction after pancreaticoduodenectomy. *J Clin Gastroenterol* 2000; 31:11-18.
4. eo, 1995a. Yeo CJ, et al: Pancreaticoduodenectomy for cancer of the head of the pancreas: 201 patients. *Ann Surg* 1995; 221:721-731.
5. Yeo, 1995b. Yeo CJ, et al: A prospective randomized trial of pancreaticogastrostomy and pancreaticojejunostomy after pancreaticoduodenectomy. *Ann Surg* 1995; 222:580-588.
6. Wente MN, Shrikande et al .Pancreaticojejunostomy versus pancre aticogastro stomy: systematic review and meta-analysis
7. Ramesh H, Pancreaticojejunostomy versus pancreaticogastrostomy in reconstruction following pancreaticoduodenectomy
8. Khan AW, Aggarwal AK, Davidson BR. (2002) Isolated Roux loop duct to mucosa pancreaticojejunostomy avoids pancreatic leaks in pancreaticoduodenectomy. *Dig Surg* 19:199–204.
9. Sutton CD, Garcea G, White SA, O'Leary E, Marshall LJ, Berry DP et al. (2004) Isolated Roux loop pancreaticojejunostomy: a series of 10 patients with zero postoperative pancreaticoenteric leaks. *J Gastrointest Surg* 8: 701–705.
11. Funovics JM, Zöch G, Wenzl E, Schulz F. (1987) Progress in reconstruction after resection of the head of the pancreas. *Surg Gynecol Obstet* 164:545–548.
12. ingsnorth AN. (1994) Safety and function of isolated Roux looppancreaticojejunostomy after Whipple's pancreaticoduodenectomy. *Ann R Coll Surg Engl* 76:175–179.
13. Albertson DA. (1994) Pancreaticoduodenectomy with reconstruction by Roux en Y pancreaticojejunostomy: no operative mortality in a series of 25 cases. *South Med J* 87:197–201.
14. Papadimitriou JD, Fotopoulos CA, Smyrniotis B, Prahalias AA, Kostopanagiotou G, Papadimitriou LJ. (1999) Subtotal pancreatoduodenectomy. Use of a defunctionalized loop for pancreatic stump drainage. *Arch Surg* 134:135–139.
15. Jover JM, Carabias A, Feurte S, Rios R, Ortega I, Limones M. (2006) Results of defunctionalized jejunal loop after pancreaticoduodenectomy. *Cir Esp* 80:373–377.
16. Machado MC, Monterio da Cunha JE, Bachella T, Bove P. (1976) A modified technique for the reconstruction of the alimentary tract after pancreatoduodenectomy. *Surg Gynecol Obstet* 143:271–273.
17. Funovics JF, Zoch G, Wenzel G, Schulz M. Progress in reconstruction after resection of the head of the pancreas. *Surg Gynecol Obstet* 1987; 164: 545-8.
18. Isolated Roux loop pancreaticojejunostomy versus pancreatico gastrostomy after pancreaticoduodenectomy: a prospective randomized study. El Nakeeb A, Hamdy E, Sultan AM, Salah T, Askr W, Ezzat H, Said M, Zeied MA, Abdallah T. *HPB (Oxford)*. 2014 Jan 28