



RELATIONSHIP OF BMI, VITAMIN D AND BMD IN AGE AND GENDER LINKED POPULATION

Priyanka Pahari

M.Sc (Medical Physiology), Assistant Professor, Department of Physiology
KPC Medical College and Hospital, Jadavpur, Kolkata

Ritwik Ganguli

MBBS, MS (orth), DNB (orth), Fellowship arthroscopy & sports medicine(Japan).
Assistant Professor, Department of Orthopaedics. KPC Medical College and Hospital,
Jadavpur, Kolkata

ABSTRACT Vitamin D insufficiency prevalence has been related to low bone mineral density (BMD). However, controversial results have been reported for the relationship between serum 25-hydroxyvitamin D [25(OH)D] levels and BMD. This study was done to investigate whether serum 25(OH)D levels were associated with BMD in different age group and sex link population. This study involved, aged 40-70 yr, who are consecutively selected from KPCMCH, BMD camp. BMD was measured at the lumbar spine and femoral neck. The correlation between serum 25(OH)D levels and BMD was investigated. Vitamin D Levels for Healthy and patients Individuals at Hospital. Subjects. The age of 40 healthy subjects ranged from 40 to 70 years with the average of 55.30±10.30 years and body mass index (BMI) ranged from 18 to 37 kg/m², with the of average of 28.90 ± 5.20 kg/m² Comparison between healthy and patients based on BMI and vitamin D level for the overweight BMI healthy individuals was 29.78 ± 9.40 ng/mL, and that of hyperlipidemic patients was 24.47 ± 8.78 ng/mL. In this study, there is significant different between healthy and patients group in vitamin D3 level. BMD significantly decreased in patients group more elderly.

KEYWORDS : Vitamin D, gender, BMI, BMD

INTRODUCTION

Vitamin D is considered essential for bone health. In some studies, vitamin D insufficiency has been reported to be associated with low bone mineral density (BMD) and increased bone loss (1,2). However, the results reported so far have been controversial (3, 4) Now Obesity is another rapidly growing health problem in most developed countries (2). During the last decade, the prevalence of obesity (body mass index (BMI) ≥30) increased dramatically. Vitamin D low levels negatively affect bone mineralization causing rickets in children and osteomalacia in adults [3,5]. In addition, vitamin D insufficiency is associated with other diseases; chronic kidney disease (CKD) gives rise to secondary hyperparathyroidism (SHPT) which can lead to loss of bone density and elevated rates of fracture in renal patients [6], common cancers [7], autoimmune disorders [8,9], multiple sclerosis [10], Cardiovascular disease [11], lung function, and asthma [12]

MATERIALS AND METHODS:

Study Population. In this cross-sectional study 100 subject have been enrolled which was conducted in March 2016 in BMD camp of KPCMCH. Out of hundred subject: 40 subjects were apparently healthy and 60 subjects were selected as a patients after taking the history of inclusion criteria included: Multiple joint pain for prolong period in adult, Low back pain with kyphotic deformity in elderly age group, young patients with pain in long bones of lower limb, History of fracture with insignificant trauma.

Serum collection and analysis: Blood samples (6 ml) were collected from each subject in the morning after an overnight fast. The blood was centrifuged for 10 min at 1000 x g to obtain serum. The serum was placed in Eppendorf tubes and stored at -80°C until used.

Measurement of Vitamin D. Quantitative colorimetric immunoenzymatic determination of 25(OH) vitamin D concentrations in human plasma level was developed by using vitamin D ELISA kit (Diametra, Milano, Italy). The kit is a competitive solid phase enzyme-linked immunosorbent assay (ELISA). Samples were analyzed according to the manufacturer guidelines.

Vitamin D levels were classified into 3 major groups [13, 14] as follows:

- (1) sufficient (>30 ng/mL);
- (2) insufficient (20–30 ng/mL);
- (3) deficient (<20 ng/mL).

BMI measurements: women were divided into six and men into five BMI groups: i) BMI <20 – underweight (only for women); ii) BMI 20-24.9 – normal weight; iii) BMI 25-29.9 –overweight; iv) BMI 30-34.9 – obesity, degree I; v) BMI 35-39.9 –obesity, degree II; vi) BMI ≥40 – super obese, obesity degree III (15).

Bone mineral density examination: BMD was determined using Dual Energy X-ray Absorptiometry (DEXA). Both spine region including lumbar vertebrae 1-4 and femoral neck area I BMD were obtained. To eliminate operator differences, all women were tested by the same operator during the study

Statistical methods: Data from 100 subjects were expressed as mean ± SD and statistically analyzed using SPSS v. Linear regression analysis was performed to assess correlations between BMI, serum 25(OH)D3 and 1,25(OH)2D3 levels, age and gender. P-values <0.05 were considered as indicating statistical significance.

RESULTS:

Table 1: Demographics data (age and body mass index) of all participants (n = 100).

| Parameter | Healthy individuals (N=40) Mean±SD | Patients (N= 60) Mean±SD |
|-------------------------|--|--------------------------------|
| Total | | |
| Age(years) | 55.30±10.30 | 55.88±9.38 |
| BMI(kg/m ²) | 28.90±5.20 | 28.82±5.49 |
| Males | | |
| Age(years) | 55.90±10.24 | 54.80±9.86 |
| BMI(kg/m ²) | 29.10±5.21 | 27.98±4.90 |
| Females | | |
| Age(years) | 54.10±10.50 | 58.20±10.04 |
| BMI(kg/m ²) | 28.40±4.50 | 28.74±5.21 |

Table 2: Mean value of vitamin D levels in ng/mL for 40 to 70 years of age in healthy and Patients at hospital

| category | Healthy Vitamin D(ng/ml) | Patient Vitamin D(ng/ml) | P value |
|----------------------|-----------------------------|-----------------------------|---------|
| Total | 30.78±10.50(N=40) | 25.98±8.83(N=60) | <0.0001 |
| Total males | 33.18±11.22(N=16) | 25.04±8.02(N=45) | <0.0001 |
| Total females | 29.55±9.98(N=24) | 23.54±7.74(N=15) | 0.0225 |
| Normal BMI Weight | 31.30±9.42(N=6) | 27.67±10.10(N=14) | 0.2209 |

| | | | |
|---------------|------------------|------------------|---------|
| Overweight | 29.78±9.40(N=33) | 24.47±8.78(N=21) | 0.0087 |
| obese | 26.51±6.90(N=14) | 20.76±6.21(N=25) | 0.0002 |
| 40-50 years | 37.42±9.08(N=20) | 26.88±9.88(N=30) | <0.0001 |
| 51-60 years | 31.07±4.65(N=12) | 26.35±6.34(N=18) | 0.0027 |
| Over 60 years | 22.52±3.56(N=08) | 17.97±6.98(N=12) | 0.0126 |

significant (P value< 0.050).

Table 1 summarizes the age and the body mass index (BMI) of the first group subjects (100 subjects (40–70 years)

Vitamin D Levels for Healthy and patients Individuals at Hospital. Subjects. The age of 40 healthy subjects ranged from 40 to 70 years with the average of 55.30±10.30 years and body mass index (BMI) ranged from 18

to 37 kg/m², with the of average of 28.90 ± 5.20 kg/m² (Table 1). Besides, the age of 60 patients ranged from 40 to 70years, with the average of 55.89 ± 9.39 years, and body mass index (BMI) ranged from 17 to 37 kg/m², with the average of 28.82 ± 4.94 kg/m² (Table 1). Further classification is also found in Table 1.

Furthermore, Table 2 showed the differences in vitamin D levels between the healthy and patients subjects. A significant difference (< 0.050) was detected between healthy males (25.04±08.02 ng/mL) and females (23.55±7.74 ng/mL).

comparison between healthy and patients based on BMI was done and vitamin D mean level for the normal BMI healthy individuals was 31.30 ±9.42 ng/mL, while that of patients was 27.67±10.10 ng/mL. Also, vitamin D level for the overweight BMI healthy individuals was 29.78 ± 9.40 ng/mL, and that of hyperlipidemic patients was 24.47 ± 8.78 ng/mL (Table 2).




Figure 1. Serum 25(OH)D3 concentrations (nmol/l) in women (A) and men (B) versus BMI.

DISCUSSION:

The finding of this study revealed that vitamin D levels are affected by many factors such as nationality, gender, sex, BMI, physical activity, and lifestyle and this was reported previously [16–21]. Obviously,

factors other than BMI are of significance for vitamin D status, as can be assumed due to the large scattering of the data points in Figure 1. The effect of BMI on serum 25(OH) D3 may be explained by the fact that persons with high BMI usually have a high content of body fat, acting as a reservoir for lipid-soluble vitamin D. It has previously been shown in animal models that body adipose tissue can accumulate about 10-12% of a supplemented dose of vitamin D (22). At the same time, the release of vitamin D from the fat is extremely slow and proportional to the concentration of the vitamin in the adipose tissue (22). This biological mechanism may have the purpose of protecting the body from toxic effects of active forms of vitamin D and maintaining an optimal level in the blood. However, excess body fat results in its increased sequestration and low availability and, as a consequence, low serum 25(OH)D levels (22, 23). A large fraction of severely obese patients undergoing surgical treatment for obesity have hypovitaminosis D before surgery (24, 25). Surprisingly, however, serum levels of 25(OH)D do not increase significantly after surgery and weight loss, even if vitamin D supplements are administered (24). Nevertheless, the magnitude of weight loss is negatively correlated with serum 25(OH) D (24).

CONCLUSION:

In this study there is significant difference between healthy and patients group in vitamin D3 level. BMD significantly decreased in patients group more elderly.

There is significant correlation between vitamin D3 level and BMD at hip and spine. Male gender, BMI and age are significant predictor of BMD. Patients with higher BMI have significantly lower BMD. So, vitamin D3 level is adversely related with BMI. It suggests that Obesity adversely affects bone health and prone to bone fragility, bone pain and fractures.

REFERENCES:

- Narula R, Tauseef M, Ahmad IA, Agarwal K, Ashok A, Anjana A. Vitamin D deficiency among postmenopausal women with osteoporosis. *J Clin Diagn Res* 2013; 7: 336-8.
- Sadat-Ali M, Al Elq AH, Al-Turki HA, Al-Mulhim FA, Al-Ali AK. Influence of vitamin D levels on bone mineral density and osteoporosis. *Ann Saudi Med* 2011; 31: 602-8.
- Wat WZ, Leung JY, Tam S, Kung AW. Prevalence and impact of vitamin D insufficiency in southern Chinese adults. *Ann Nutr Metab* 2007; 51: 59-64.
- von Mühlen DG, Greendale GA, Garland CF, Wan L, Barrett-Connor E. Vitamin D parathyroid hormone levels and bone mineral density in community-dwelling older women: the Rancho Bernardo Study. *Osteoporos Int* 2005; 16: 1721-6.
- MF. Holick and T. C. Chen. "Vitamin D deficiency: a worldwide problem with health consequences." *The American Journal of Clinical Nutrition*, vol. 87, no. 4, pp. 1080S–1086S, 2008.
- AS. Dusso, "Kidney disease and vitamin D levels: 25-hydroxyvitamin D, 1,25-dihydroxyvitamin D, and VDR activation," *Kidney International Supplements*, vol. 1, no. 4, pp. 136–141, 2011.
- KC. Chiang, C.-N. Yeh, M.-F. Chen, and T. C. Chen, "Hepatocellular carcinoma and vitamin D: a review," *Journal of Gastroenterology and Hepatology*, vol. 26, no. 11, pp. 1597–1603, 2011.
- JJ. Cannell, R. Vieth, J. C. Umhau et al., "Epidemic influenza and vitamin D," *Epidemiology and Infection*, vol. 134, no. 6, pp. 1129–1140, 2006.
- PT. Liu, S. Stenger, H. Li et al., "Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response," *Science*, vol. 311, no. 5768, pp. 1770–1773, 2006.
- G. Giovannoni and G. Ebers, "Multiple sclerosis: the environment and causation," *Current Opinion in Neurology*, vol. 20, no. 3, pp. 261–268, 2007.
- T. L. Larose, Y. Chen, C. A. Camargo Jr., A. Langhammer, P. Romundstad, and X.-M. Mai, "Factors associated with vitamin D deficiency in a Norwegian population: the HUNT Study," *Journal of Epidemiology and Community Health*, vol. 68, no. 2, pp. 165–170, 2014.
- T. L. Larose, A. Langhammer, Y. Chen, C. A. Camargo Jr., P. Romundstad, and X. M. Mai, "Serum 25-hydroxyvitamin D levels and lung function in adults with asthma: the HUNT Study," *The European Respiratory Journal*, vol. 45, pp. 1019–1026, 2015.
- MF. Holick, N. C. Binkley, H. A. Bischoff-Ferrari et al., "Evaluation, treatment, and prevention of vitamin D deficiency: an endocrine society clinical practice guideline," *Journal of Clinical Endocrinology and Metabolism*, vol. 96, no. 7, pp. 1911–1930, 2011.
- C. Annweiler, G. Allali, P. Allain et al., "Vitamin D and cognitive performance in adults: a systematic review," *European Journal of Neurology*, vol. 16, no. 10, pp. 1083–1089, 2009.
- AA Ginde, M. C. Liu, and C. A. Camargo Jr., "Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004," *Archives of Internal Medicine*, vol. 169, no. 6, pp. 626–632, 2009.
- N. M. Al-Daghri, K. M. Alkharfy, A. Al-Othman et al., "Effect of gender, season, and vitamin D status on bone biochemical markers in Saudi diabetes patients," *Molecules*, vol. 17, no. 7, pp. 8408–8418, 2012.
- A. Al-Othman, S. Al-Musharaf, N. M. Al-Daghri et al., "Effect of physical activity and sun exposure on vitamin D status of Saudi children and adolescents," *BMC Pediatrics*, vol. 12, article 92, 2012.
- A. Al-Othman, S. Al-Musharaf, N. M. Al-Daghri et al., "Tea and coffee consumption in relation to vitamin D and calcium levels in Saudi adolescents," *Nutrition Journal*, vol. 11, no. 1, article 56, 2012.
- B. Hamilton, J. Grantham, S. Racinais, and H. Chalabi, "Vitamin D deficiency is endemic in Middle Eastern sportsmen," *Public Health Nutrition*, vol. 13, no. 10, pp. 1528–1534, 2010.
- NA Meguid, A. F. Hashish, M. Anwar, and G. Sidhom, "Reduced serum levels of 25-hydroxy and 1,25-dihydroxy vitamin D in Egyptian children with autism," *Journal of Alternative and Complementary Medicine*, vol. 16, no. 6, pp. 641–645, 2010.

21. S. Racinais, B. Hamilton, C. K. Li, and J. Grantham, "Vitamin D and physical fitness in Qatari girls," *Archives of Disease in Childhood*, vol. 95, no. 10, pp. 854-865, 2010.
22. Rosenstreich SJ, Rich C and Volwiler W: Deposition in and release of vitamin D3 from body fat: evidence for a storage site in the rat. *J Clin Invest* 50: 679-687, 1971.
23. Wortsman J, Matsuoka LY, Chen TC, Lu Z and Holick MF: Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 72: 690-693, 2000.
24. Abbasi AA, Amin M, Smiertka JK, Grunberger G, MacPherson B, Hares M, Lutrzykowski M and Najjar A: Abnormalities of vitamin D and calcium metabolism after surgical treatment of morbid obesity: a study of 136 patients. *Endocr Pract* 13: 131-136, 2007.
25. Carlin AM, Rao DS, Meslemani AM, Genaw JA, Parikh NJ, Levy S, Bhan A and Talpos GB: Prevalence of vitamin D depletion among morbidly obese patients seeking gastric bypass surgery. *Surg Obes Relat Dis* 2: 98-103, 2006.