



RELATION OF TYPE 2 DIABETES AND BMI WITH AUDITORY ACUITY

*Isaac David

P.G.Student, Department of physiology, MGM College, Aurangabad. Maharashtra, India. *Corresponding Author

Shinde P.U

Professor, Department of physiology, MGM College, Aurangabad. Maharashtra, India.

ABSTRACT**Introduction:**

Diabetes-related sensorineural hearing impairment affects people's ability to hear and understand sounds. This is a case control study with the purpose of determining the hearing loss in type II diabetes mellitus in relation to the BMI of the patients Method: 50 diabetic patients and 50 age and sex matched controls who have satisfied inclusion and exclusion criteria and have consented to participate in study were enrolled. They were evaluated by Tuning fork tests and Pure Tone Audiometry.

Result:

The hearing of diabetics was significantly impaired than the non-diabetic control group. This hearing impairment was noted in all the frequencies tested. The hearing acuity was compared with BMI in diabetics. The hearing acuity was not influenced by normal BMI levels. However at higher values of BMI, the hearing loss was significant, particularly at the right side (right ear). The significance was more pronounced in case of obese individuals with BMI > 30.

Conclusion:

Type 2 diabetes causes significant hearing loss in the patients, and it is affected by the higher BMI of the patient particularly BMI>30.

KEYWORDS : Audiometry, Diabetes, Sensorineural hearing loss, BMI**Introduction:**

Obesity is a disorder of energy balance affecting wide range of people belonging to diverse ethnic groups, age and socioeconomic status. Obesity a major risk factor for many acute and chronic disorders/ including cardiovascular and cerebrovascular disease and diabetes. Diabetes Mellitus is a metabolic disorder, due to relative or absolute lack of insulin resulting in elevated blood glucose levels associated with long term vascular and neurological complications.¹ Among glucose metabolism disorders, diabetes mellitus is the one most commonly related with auditory disorders. Type-2 diabetes mellitus (T2DM) is the predominant form of diabetes worldwide accounting for 90% of cases globally. One of the worst affected nations in the South East Asia region is India.² The most common complications of diabetes are diabetic ketoacidosis, hyperglycemic hyperosmolar coma, diabetic retinopathy, macular oedema, neuropathy, coronary artery disease, peripheral vascular disease and cerebrovascular disease.³ Hearing impairment is one of the under recognized complications of diabetes. The characteristic finding in diabetes mellitus is a progressive, gradual, bilateral symmetrical sensorineural hearing loss (SNHL) particularly in higher frequencies.^{4,5} It would be similar to presbycusis, but with more severe losses than those expected by ageing.⁶ Pure tone Audiometry is a simple, non-invasive test and can detect type and degree of the hearing loss.⁷ Pure tone Audiometry involves the estimation of the threshold of hearing for certain standardized stimuli via the air and bone conduction routes.⁸ Since many studies have reported contradicting results regarding hearing impairment in diabetic patients and only a few studies are done in India hearing impairment in diabetic patients in relation with BMI level, the present study is undertaken to determine the incidence of auditory dysfunction in type-2 diabetes persons and whether or not such auditory dysfunction could be correlated with the BMI.

MATERIAL AND METHOD:**study Design:**

Present study is a case control study involving 100 subjects, divided into two groups. Group-1 comprised 50 voluntarily willing normal, healthy subjects Group-2 comprised 50 voluntarily willing diabetic patients.

Method:

Group 1: Inclusion Criteria included normal, healthy subjects of either sex between 20 and 50 years. Exclusion Criteria 1) Hypertension 2) Diabetes mellitus, 3) History of consumption of ototoxic drugs in past three months, 4) History of ear surgeries, 5) History of recent infections in ear, nose or throat.

Group2: Inclusion Criteria included Type 2 diabetic patients of either

sex between 20 and 50 years. Exclusion Criteria 1) Hypertension 2) History of consumption of ototoxic drugs in past three months. 3) History of ear surgeries, 4) History of recent infections in ear, nose or throat. Informed and written consent, questionnaires including personal information (age, H/o of surgery, drugs, occupation) was obtained. Institutional ethical committee had cleared the project. Group2 subjects were screened to identify the obesity by grouping: BMI<20Kg/m², BMI 20-25 Kg/m², BMI25- 30 Kg/m², BMI>30Kg/m².

Tests for Hearing:

The tuning fork tests- Weber test, Rinne test and Absolute bone conduction tests were done for both the ears of all subjects.

Pure Tone Audiometry - is the most routine audiometric evaluation. It is based on the measurement of hearing thresholds for a range of pure tones presented through earphones according to the ascending method (Hughson - Westlake, up 5, down 10 method) Audiological examination was performed using a Pure Tone Audiometer model(EDA Giga 3 of ELKON). The audiometer [ELKON EDA Giga 3] is an electronic device that produces pure tones, the intensity of which can be increased or decreased in 5-dB steps. It was performed in a sound proof room in the ENT department, MGM'S Medical College and Hospital. The patient was described what will happen during the test and the purpose of the test. Ear phones were used to test hearing by air conduction and a small vibrator placed over the mastoid was used to test hearing by bone conduction. All audiometers incorporate a calibration circuit, which allows the output sound level to be set at each frequency. The signals presented to the subject by an audiometer were characterized by its frequency, sound pressure level and wave form which were all controlled. Biological calibration was done every day before starting the test. Both air and bone conduction were tested for each ear. Air conduction thresholds were measured for tones of 250, 500, 1000, 2000, 4000 6000 and 8000 Hertz. Bone conduction thresholds were measured for 250, 500, 1000, 2000, 4000 Hz.

Body mass index was calculated as: Body mass index = Weight (Kilograms)/ Height (Meter)².

Statistical analysis:

The hearing loss was evaluated on the basis of the values of PTA obtained for right and left side. PTA values exceeding 25 indicated hearing loss. On the basis of PTA > 25, the comparisons within and among various groups were evaluated by students t test. 'p' value > 0.05 was considered non significant, 'p' value < 0.05 was considered significant, 'p' value < 0.01 was considered highly significant,

RESULTS AND DISCUSSION:

100 subjects (**group 1**, normal, healthy n=50) and (**group 2**, diabetic n=50) that have satisfied the inclusion and exclusion criteria were selected.

Table 1: Comparative statistics: of age, blood sugar level (fasting and post meal) Pure Tone Audiometry (right & left ear)

	Parameter	Control (n=50)	Diabetic (n=50)
1.	Age	39.3 ± 6.42	40.34 ± 6.13
2.	BSL – F	85.46 ± 6.34	190 ± 53.23
3.	BSL – PP	112.4 ± 6.43	285.06 ± 74.42
4.	PTA – R	17.86 ± 6.58	25.04 ± 16.32
5.	PTA – L	17.74 ± 5.43	26.58 ± 16.02
6.	SNHL Total	6	37
	- Mild	4	20
	- Moderate	2	10
	- Mod-Severe	-	3
	- Severe	-	3
	- Profound	-	1

Table – 1: Shows the comparative data. There were 100 subjects, controlled, normoglycemic n =50 and hyperglycemic n = 50 in the age range of 28 to 50 years. In the control group the fasting sugar and post meal glucose ranged from 76 to 102 mg/dl. and 98 to 124 mg/dl. The fasting blood glucose and post meal glucose in hyperglycemic group ranged from 96 to 368 mg/dl and 120 to 556 mg/dl. Based on pure tone average a total of 6 subjects in controlled group were found to have SNHL (4 mild and 2 moderate) and 37 subjects in hyperglycemic group (20 mild, 10 mod, 3 mod-severe, 3 severe and 1 profound) were found to have SNHL.

Table 2: Classification of obesity based on measurement of BMI according to WHO⁽¹⁰⁾

Normal --	18.5-22.9
Average Overweight	-- > 23
At risk --- Increased	23-24.9
Obese – I – Moderate	---25-29.9
Obese – II Severe	>30

Table-3 Comparison of Hearing Loss between Control and Diabetic Group

Hearing Loss	Control (n=50)	Diabetic (n=50)	t Value	p Value
SNHL – R	17.86 ± 6.57	25.04 ± 16.31	2.885963	0.004799
SNHL – L	17.74 ± 5.43	26.58 ± 16.01	3.69526	0.000362

Table – 3: showed highly significant increase in PTA levels in both ears in diabetic patients where compared with controls. Based on this observation, it was found that diabetic subjects were more prone to developing SNHL when compared with controls.

Table 4: Comparison of BMI and hearing loss

Hearing loss	BMI < 20 (n=1)	t value	p value
SNHL-Right ear	31 ± 0	-	-
SNHL-Left ear	38 ± 0	-	-

Hearing loss	BMI 20.1 to 24.9 (n=17)	t value	p value
SNHL-Right ear	24.70 ± 15.55	0.393224	0.6994
SNHL-Left ear	28.05 ± 15.72	0.614432	0.5476

Hearing loss	BMI 25 to 29.9 (n=28)	t value	p value
SNHL-Right ear	30.84 ± 8.86	2.659928	0.0205
SNHL-Left ear	11.51 ± 13.51	1.902974	0.0897

Hearing loss	BMI > 30 (n=4)	t value	p value
SNHL-Right ear	14.75 ± 4.43	3.275692	0.0466
SNHL-Left ear	16.75 ± 3.03	6.270544	0.0082

Table 4 shows that there was no significant difference between the hearing loss under the influence of Body Mass Index (BMI) when the BMI was normal i.e. below 25. However at higher values of BMI, the hearing loss was significant, particularly at the right side (right ear). The significance was more pronounced in case of obese individuals with BMI > 30.

Discussion:

Most of the recent studies show an association of SNHL with diabetes. This study also supports the association of SNHL with diabetes with an

incidence of 74% as compared to 12% among non-diabetics. All diabetic patients who reported hearing loss had slow progressive hearing loss. In this study diabetic patients had a higher threshold for high frequency there was significant hearing loss higher values of BMI, particularly at the right side (right ear). The significance was more pronounced in case of obese individuals with BMI > 30.

Mechanism that may underlie the relation between high BMI and hearing function include hypoxia and ischemic damage, oxidative stress and formation of reactive oxygen species, and resultant death of cochlear and spiral ganglion cells that leads to hearing loss.⁹ Obesity related atherosclerosis may lead to stiffening and constriction of the internal auditory artery and reduction in cochlear blood flow.¹⁰ Reduced blood supply to cochlea, whether due to micro-vascular or macro-vascular compromise, can lead to capillary constriction within the stria vascularis, cell death, and poorer hearing sensitivity.^{11,12}

The tissue effects of diabetes are thought to be related to the polyol pathway, where glucose is reduced to sorbitol. Sorbitol accumulation is implicated in neuropathy by causing a decrease in myoinositol content, abnormal phosphoinositide metabolism, and a decrease in Na⁺/K⁺ ATPase activity.¹³ They also observed that the VIIIth nerve showed signs of myelin degeneration, with fibrosis of the perineurium.

Conclusions:

In present case controlled study There was a significant increase in incidence of sensorineural hearing loss in the diabetic group as compared to the control group. Significant difference in hearing loss was observed when the BMI level was high particularly in subjects having BMI more than 30.

REFERENCES:

- Gareth Williams, John C. Pickup. Text book of diabetes mellitus. Blackwell publishing Ltd, Oxford, Third edition, 2003
- Buse JB, Polonsky KS, Brunt CF. Type 2 Diabetes Mellitus. In: Melmed S, Polonsky KS, Larsen PR, Kronenberg HM, editors. 'William's Textbook of Endocrinology'. 12th edition. Philadelphia: Elsevier / Saunders; 2011. p. 1371-1435.
- Powers AC. Diabetes Mellitus. In: Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, Loscalzo J, editors. 'Harrison's Principles of Internal Medicine'. 18th ed. New York 2012: 2968-3003
- Kalli k et al. 'Hearing disturbance in patients suffering from diabetes'. Otorhinolaryngology: Head and Neck Surgery 2012;48:22-24.
- Iranifb, Isaac David. 'Relation of type 2 Diabetics and haemoglobin with auditory acuity' IJHSR. 2017; vol 7(10)
- Taylor, I.G., Irwin, J. 'Some audiological aspects of diabetes mellitus'. J Laryngol Otol. 1978;92:99-113
- Biswas A. 'Clinical Audio-Vestibulometry for Otolologists and Neurologists'. Fourth edition. Bhalani publishing house. New Delhi. p. 100-117
- Kerr AG, Stephens D. 'Scott-Browns Otolaryngology: Adult Audiology'. Volume 2. 6th ed. Butterworth Heinemann; 1997. P. 2/1/6-9
- Curhan SG, Eavey R, Wang M, Stampfer MJ, Curhan GC. Body mass index, waist circumference, physical activity, and risk of hearing loss in women. Am J Med. Dec 2013; 126(12): 1142. doi: 10.1016/j.amjmed.2013.04.026. Epub 2013 Oct 11.
- Makishma K. Arteriolar sclerosis as a cause of presbycusis. Otolaryngology. Mar-Apr 1978; 86920: 322-326. [pubmed: 113738].
- Gates GA, Cobb JL, D'Agostine RB, Wolf PA. The relation of hearing in the elderly to the presence of cardiovascular disease and cardiovascular risk factors. Arch Otolaryngol Head Neck Surg. Feb 1993; 119(2): 156-161. [pubmed: 8427676].
- Seidman MD, Quirk WS, Shirwany NA. Mechanisms of alterations in the microcirculation of the cochlea: Ann N Y Acad Sci. Nov 28, 1999; 884: 226-232. [pubmed: 10842596].
- Makishima K, Tanaka AK. Pathological changes of the inner ear and central auditory pathway in diabetes. Ann Otol Rhinol Laryngol. 1971;80(2):218-28