



## VALUATING THE RADIOPATHOLOGIC CORRELATION OF COMMONLY OCCURRING CNS TUMOURS

**Muthuraman  
pichaiah**

Assistant Professor, Department of Neurosurgery. Thanjavur Medical College, Thanjavur.

**Senthamarai  
kannan.  
S**

Assistant professor, department of neurosurgery, Thanjavur Medical College, Thanjavur

### ABSTRACT

**AIM :** To study the correlation between the Pathological nature and the radiological appearance of primary Central Nervous System neoplastic lesions based on the two most commonly used and easily available investigations namely CT Brain plain and contrast and MRI Brain plain and contrast.

**MATERIALS AND METHOD :** The study was conducted on all the patients admitted in the concerned admitting unit with a diagnosis of a central nervous system tumour between May 2013 to June 2015 at Thanjavur Medical College Hospital, Thanjavur. Total number of 243 patients were enrolled for the study.

**CONCLUSION :** MRI had a higher or equal predictive value when compared to CT but statistically not significant. MRI had a higher predictive value for benign lesions than CT brain. Malignant lesions had more or less equal value for both CT and MRI low grade glioma is an important differential diagnosis for all the intrinsic tumours.

**AIM OF THE STUDY :** To study the correlation between the Pathological nature and the radiological appearance of primary Central Nervous System neoplastic lesions based on the two most commonly used and easily available investigations namely CT Brain plain and contrast and MRI Brain plain and contrast. To identify the specificity of CT Vs MRI in achieving the same. To assess the cost effectiveness of the imaging methods and find out which had better specificity.

### KEYWORDS :

#### INTRODUCTION

Neoplasms of the central nervous system are comparatively infrequent, though it is not as rare a condition as it was once assumed to be. From the very early days of very high mortality and morbidity rates, due to CNS tumours, substantial improvement in survival and outcome has been made because of the development of advanced, state of the art, imaging modalities which have made early diagnosis possible and improvement in neurosurgical equipments and techniques that allowed greater accessibility permitting a greater chance of gross total surgical resection and advanced radiation therapy to counter the rapid growth of the tumour. The advent of modern cross-sectional imaging techniques especially MR imaging have completely changed the method of assessment for follow up in affected patients. Towards this goal, the spectrum of cross sectional imaging manifestations of commonly occurring primary central nervous system neoplasms like astrocytoma, meningioma, medulloblastoma, ependymoma and craniopharyngioma. A comprehensive summation of the correlation between the radiological appearances of these tumours and their pathological nature is studied.

#### STUDY MATERIALS AND METHODS

The study was conducted on all the patients admitted in the concerned admitting unit with a diagnosis of a central nervous system tumour between May 2013 to June 2015 at Thanjavur Medical College Hospital, Thanjavur. Total number of 243 patients were enrolled for the study.

#### Exclusion Criteria

1. Patients who died before surgery.
2. Patients admitted with recurrence.
3. Patients who were admitted with a proved pathological diagnosis at some other institute and later referred here.
4. Tumours which were not routinely biopsied at the institute like, brain-stem gliomas.
5. Patients who presented with haemorrhage at the tumour site were not included because of their distorted radiological appearance.
6. All the patients had CT Brain plain and contrast. Patients who had only plain MRI without contrast were included in the CT Brain group, their MRI was not taken into consideration for the study.
7. Patients who had multiple lesions were excluded from the study.
8. After admission patients who were diagnosed with a primary lesion elsewhere in the body were excluded from the study.

The above 191 patients were divided into two groups.

- Group I - Only CT Brain plain and contrast.
- Group II - CT Brain and MRI Brain plain and contrast.
- Group I - Contained 191 patients
- Group II - Contained 82 patients

Radiology criteria for the 6 types of tumours to be studied were formulated.

Group I had 153 patients

Group II had 67 patients

Group I had the following number of patients in each category.

Low Grade Glioma	-	27
High grade Glioma	-	47
Medulloblastoma	-	24
Ependymoma	-	7
Craniopharyngioma	-	17
Meningioma	-	31
Total	-	153

Group II had the following number of patients in each category.

Low Grade Glioma	- 15
High grade Glioma	- 17
Medulloblastoma	- 11
Ependymoma	- 2
Craniopharyngioma	- 7
Meningioma	- 15
Total	- 67

After surgery, the histopathological diagnosis of all the selected 153 patients were entered. Based on the above data a master chart was prepared. The correlation and measure of agreement were analysed statistically and the results are discussed in the following pages.

For Histopathological examination, light microscopy with routine eosin and haematoxylin stains were used.

For the statistical analysis chi-square pearson formula was used.

#### RESULTS AND ANALYSIS

##### Table - 1

Table showing Correlation between MRI and Pathology.

MRI Diagnosis	Pathology							Total
	Low Grade Glioma	High Grade Glioma	Medulloblastoma	Ependymoma	Cranio-pharyngioma	Meningioma		
Low grade Glioma	15							15
High Grade Glioma	2	15						17
Medulloblastoma			10	1				11
Ependymoma				2				2
Cranio-pharyngioma					6			6
Meningioma						12		12
Total	17	15	10	3	6	12		63

Kappa (measure of agreement) = 94.0% (p < 0.001)

**Table-2**  
**Table showing Correlation between MRI and Pathology (Including Other Diagnosis)**

MRI Diagnosis	Pathology							Total
	Low Grade Glioma	High Grade Glioma	Medulloblastoma	Ependymoma	Cranio-pharyngioma	Meningioma	Others	
Low grade Glioma	15							15
High Grade Glioma	2	15						17
Medulloblastoma			10	1				11
Ependymoma				2				2
Cranio-pharyngioma		1			6			7
Meningioma						12	3	15
Not available	16	27	12	3	9	13	6	86
Total	33	43	22	6	15	25	9	153

**Table-3**  
**Correlation between CT and Pathology**

CT Diagnosis	Pathology							Total
	Low Grade Glioma	High Grade Glioma	Medulloblastoma	Ependymoma	Cranio-pharyngioma	Meningioma		
Low grade Glioma	25							25
High Grade Glioma	3	41						44
Medulloblastoma	2		21	1				24
Ependymoma	1		1	5				7

Cranio-pharyngioma	1				15		16
Meningioma	1	1				25	27
Total	33	42	22	6	15	25	143

Kappa (Measure of Agreement) = 90.3% (p < 0.001)

**Table-4**  
**Correlation between CT and Pathology (Including other Diagnosis)**

CT Diagnosis	Pathology							Total
	Low Grade Glioma	High Grade Glioma	Medulloblastoma	Ependymoma	Cranio-pharyngioma	Meningioma	Others	
Low grade Glioma	25						2	27
High Grade Glioma	3	41					3	47
Medulloblastoma	2		21	1				24
Ependymoma	1		1	5				7
Cranio-pharyngioma	1	1			15			17
Meningioma	1	1				25	4	31
Total	33	43	22	6	15	25	9	153

**Table-5**  
**Low Grade Glioma CT Vs Pathology**

CT Diagnosis	Pathology Confirmed	Agreement %
27	25	92.59%

**Table-6**  
**Low Grade Glioma MRI Vs Pathology**

MRI Diagnosis	Pathology Confirmed	Agreement %
15	15	100%

**Table 7**  
**Low Grade Glioma - Positive percentage**

CT Diagnosis	MRI Diagnosis
92.59%	100%

**Table-8**  
**High Grade Glioma CT Vs Pathology**

CT Diagnosis	Pathology Confirmed	Agreement %
47	41	87.23%

**Table - 9 High Grade Glioma MRI Vs Pathology**

MRI Diagnosis	Pathology Confirmed	Agreement %
17	15	88.24%

**Table-10**  
**High Grade Glioma - Positive Percentage**

CT Diagnosis	MRI Diagnosis
87.23%	88.24%

**Table-11 Medulloblastoma CT Vs Pathology**

CT Diagnosis	Pathology Confirmed	Agreement %
24	21	87.5%

**Table-12 Medulloblastoma MRI Vs Pathology**

MRI Diagnosis	Pathology Confirmed	Agreement %
11	10	90.91%

**Table-13**  
**Medulloblastoma - Positive Percentage**

CT Diagnosis	MRI Diagnosis
87.5 %	90.94%

**Table – 14 Ependymoma CT Vs Pathology**

CT Diagnosis	Pathology Confirmed	Agreement %
7	5	71.43%

**Table – 15 Ependymoma MRI Vs Pathology**

MRI Diagnosis	Pathology Confirmed	Agreement %
2	2	100%

**Table – 16 Ependymoma - Positive Percentage**

CT Diagnosis	MRI Diagnosis
71.43 %	100 %

**Table – 17 Craniopharyngioma CT Vs Pathology**

CT Diagnosis	Pathology Confirmed	Agreement %
17	15	88.24%

**Table – 18**

**Craniopharyngioma MRI Vs Pathology**

MRI Diagnosis	Pathology Confirmed	Agreement %
7	6	85.71%

**Table – 19 Craniopharyngioma – Positive Percentage**

CT Diagnosis	MRI Diagnosis
88.24 %	85.71%

**Table – 20 Meningioma**

**CT Vs Pathology**

CT Diagnosis	Pathology Confirmed	Agreement %
31	25	80.65%

**Table – 21 Meningioma**

**MRI Vs Pathology**

MRI Diagnosis	Pathology Confirmed	Agreement %
15	12	80%

**Table – 22 Meningioma – Positive Percentage**

CT Diagnosis	MRI Diagnosis
80.65%	80.00%

**Table – 23**

**MRI Correlation for the Selected Tumours**

MRI	Positive	Negative
Low Grade Glioma	100.00	0.00
High Grade Glioma	88.24	11.76
Medulloblastoma	90.91	9.09
Ependymoma	100.00	0.00
Craniopharyngioma	85.71	14.29
Meningioma	80.00	20.00

**Table – 24**

**CT Correlation for the Selected Tumours**

CT	Positive	Negative
Low Grade Glioma	92.59	7.41
High Grade Glioma	87.23	12.77
Medulloblastoma	87.50	12.50
Ependymoma	71.43	28.57
Craniopharyngioma	88.24	11.76
Meningioma	80.65	19.35

**Table – 25 Correlating CT Diagnosis with MRI Diagnosis**

CT Diagnosis	MRI Diagnosis							Total
	Low Grade Gliom	High Grade Gliom	Medulloblastoma	Ependy moma	Craniop haryng ioma	Menin gioma	Not Avail able	
Low grade Glioma	15						12	27
High Grade Glioma		16					31	47

Medulloblastoma			11				13	24
Ependymoma				2			5	7
Craniopharyngioma					7		10	17
Meningioma		1					15	31
Total	15	17	11	2	7		15	153

**Significance of the difference in measures of agreement between CT and MRI**

**Table – 26 For Low Grade Glioma**

CT	MRI
25	15
2	0
27	15

Not Significant

**Table – 27 For High Grade Glioma**

CT	MRI
41	15
6	2
47	17

Not Significant

**Table – 28 For Medulloblastoma**

CT	MRI
21	10
3	1
24	11

Not Significant

**Table – 29 For Ependymoma**

CT	MRI
5	2
2	0
7	2

Not Significant

**Table – 30 For Craniopharyngioma**

CT	MRI
15	6
2	1
17	7

Not Significant

**Table – 31 For Meningioma**

CT	MRI
25	12
6	3
31	15

Not Significant

**Radiological Wrong Diagnosis**

**Table – 32 Wrong Radiological Diagnosis for Low grade glioma Group –1 - CT**

Wrong Diagnosis	Case No.	Total
Haemangioblastoma	80 and 145	2
Total		2

**Table – 33 Group – 2 - MRI**

Wrong Diagnosis	Nil
Total	Nil

**Table – 34**  
**Wrong Radiological Diagnosis for High Grade Glioma Group – 1 - CT**

Wrong Diagnosis	Case No.	Total
Low Grade Glioma	14, 52 and 151	3
Abscess	15	1
Secondaries	127	1
Tuberculoma	151	1
Total		6

**Table– 35 Group – 2 - MRI**

Wrong Diagnosis	Case No.	Total
Low Grade Glioma	14 and 52	2
Total		2

**Table – 36**  
**Wrong Radiological Diagnosis for Medulloblastoma Group – 1 - CT**

Wrong Diagnosis	Case No.	Total
Low Grade Glioma	64 and 72	2
Ependymoma	89	1
Total		3

**Table–37 Group – 2 - MRI**

Wrong Diagnosis	Case No.	Total
Ependymoma	89	1
Total		1

**Table – 38**  
**Wrong Radiological Diagnosis for Ependymoma Group – 1 - CT**

Wrong Diagnosis	Case No.	Total
Low Grade Glioma	53	1
Medulloblastoma	60	1
Total		2

**Table–39 Group – 2 - MRI**

Wrong Diagnosis	Total
Wrong Diagnosis	Nil
Total	Nil

**Table – 40**  
**Wrong Radiological Diagnosis for Craniopharyngioma Group - 1 - CT**

Wrong Diagnosis	Case No.	Total
High Grade Glioma	21	1
Low Grade Glioma	123	1
Total		2

**Table–41 Group – 2 - MRI**

Wrong Diagnosis	Case No.	Total
High Grade Glioma	21	1
Total		1

**Table – 42**  
**Wrong Radiological Diagnosis for Meningioma Group – 1 - CT**

Wrong Diagnosis	Case No.	Total
Tuberculoma	6 and 38	2
High Grade Glioma	7	1
Secondaries	18	1
Low Grade Glioma	30	1
Schwannoma	70	1
Total		6

**Table – 43 Group – 2 - MRI**

Wrong Diagnosis	Case No.	Total
Tuberculoma	6 and 38	2
Schwannoma	70	1
Total		3

**Table 44**  
**CT Diagnosis Malignant –Vs Benign**

CT Diagnosis	Pathology				Total	
	Malignant		Benign		Count	%
	Count	%	Count	%		
Malignant	101	95.28	4	8.51	105	68.63
Benign	5	4.72	43	91.49	48	31.37
Total	106	100.00	47	100.00	153	100.00

**Table 45**  
**MRI Diagnosis Malignant –Vs Benign**

MRI	Pathology				Total	
	Malignant		Benign		Count	%
	Count	%	Count	%		
Malignant	45	97.83			45	67.16
Benign	1	2.17	21	100.00	22	32.84
Total	46	100.00	21	100.00	67	100.00

**Table 46**  
**Positivity Agreement for Malignant Lesions CT Vs MRI**

CT	MRI
95.28	97.83

**Table 47**  
**Positivity Agreement for Benign Lesions CT Vs MRI**

CT	MRI
91.49	100

**Table 48**  
**Correlation between CT and MRI for the Group II Patients**

CT	MRI	%
67	66	98.50

**DISCUSSION**  
**GROUP I PATIENTS - CT BRAIN**

The pathological evaluation for the 153 patients were as follows:

Low Grade gliomas	-	33
High grade Gliomas	-	43
Medulloblastoma	-	22
Ependymoma	-	6
Craniopharyngioma	-	15
Meningioma	-	25
Others	-	9
Tuberculoma	-	3
Abscess	-	1
Secondaries	-	2
Schwannoma	-	1
Haemangioblastoma	-	2

**GROUP II: PATIENTS - MRI BRAIN**

The pathological evaluation for the 67 patients were as follows:

Low Grade gliomas	-	17
High grade Gliomas	-	16
Medulloblastoma	-	10
Ependymoma	-	3
Craniopharyngioma	-	6
Meningioma	-	12
Others	-	3
Tuberculoma	-	2
Schwannoma	-	1

Among group II patients, of the 15 patients with a radiological diagnosis of Low grade glioma, all the 15 has been reported as low grade glioma in histopathology study giving a measure of agreement of 100% for MRI (Table 6).

Among group I patients, of the 27 patients with a radiological low grade glioma diagnosis, 25 were low grade glioma on pathological examination, and the remaining two, were reported as haemangioblastomas. The measure of agreement was 92.59%, when compared with the 100% for MRI. The difference was not statistically significant (Table 25). The only radiological wrong diagnosis for low grade glioma among Group I patients was haemangioblastomas occurring in both patients, Cases No. 80, 145 (Table 32).

For the 47 patients in Group I with a radiological appearance of high grade glioma, 41 were reported as high grade glioma with a positive measure of 87.23% (Table 8). Among the different pathological diagnosis, three were low grade gliomas, (Cases No. 14, 52, 151), one was an abscess (case no. 15), one

secondaries brain (case no.- 127) and one was a tuberculoma (case

no.151), having a common radiological wrong diagnosis of low grade glioma occurring 50% (Table 34).

As per table (9), a positive measure of 88.24 was found among Group II patients which was not significant when compared with group I patients (Table 26). Among the two radiological wrong diagnosis both were low grade gliomas cases 14 and 52 (Table 35).

So, low grade gliomas were the commonest radiological wrong diagnosis encountered in both group I and group II patients.

On evaluating the correlation for medulloblastoma, group one had 87.5% Table (11) agreement as against 90.9% for Group II patients Table (11). Among Group I, Low grade glioma was the common radiological wrong diagnosis with two out of three cases (Case No. 64 and 72), the other being an ependymoma (Case No. 89, Table 36). In group II, the only radiological wrong diagnosis was an ependymoma (Case No.89, Table 37) which occurred in Group I also. The agreement percentage between Group I and Group II patients was not significant statistically (Table 27).

The measure of agreement of Group I was 71.43% among the 7 patients (Table 14) as compared with 100% for Group II patients (Table 15) among ependymoma. But statistically considering, the difference was not significant (Table 28).

The two patients with radiological wrong diagnosis in the Group I were one low grade glioma-(Case No. 53) and one medulloblastoma-(Case No. 60, Table 38).

On considering craniopharyngioma there was a positive measure of 88.24% for Group I (Table 17) and 85.71% for Group II (Table 18) with the difference of agreement measure being not significant (Table 29).

The two radiological wrongly diagnosed lesions in Group I were one high grade astrocytoma (Case No. 21), and the other was a low grade glioma (Case No. 123 Table 40).

In group II the radiological wrongly diagnosed case was a high grade astrocytoma Case No. 21 as in Group I (Table 41).

On evaluating Meningioma, the positive measure of Group I was 80.65 (Table 20) and for Group II it was 80.00 (Table 21).

In Group I, among the six radiologically wrong diagnosis for meningiomas there were two-tuberculomas, one - high grade glioma, one- secondaries brain, one - low grade glioma and a schwannoma, resulting in a wide spectrum of varied diagnosis when compared with all other tumors which had a very frequently occurring radiological wrong diagnosis (Table 42).

In Group II, there were two tuberculomas and one schwannoma, the same cases that occurred in group I also (Table 43).

The measure of agreement between the two groups was not significant (Table 30).

Among the above results group II had higher correlation percentage when compared with Group I for all tumours except Craniopharyngioma and meningioma although the differences were marginal and not significant (Tables 22 and 23).

Overall group II had a kappa value of 94% ( $p < 0.001$ ) where kappa is a measure of agreement.

Group I had a kappa value of 90.3% ( $p < 0.001$ )

On considering meningioma and craniopharyngioma as benign lesions and the remaining four as malignant, the positive predictive value for Group II was 97.83 for malignant and 100% for Benign, slightly higher than that of Group I patients with 95.28 in malignancies and 91.49 for benign lesions. Considering these two groups statistically, it was not significant (Tables 44 and 45).

Among wrong diagnosis, Haemangioblastoma occurred in two patients when radiologically it was diagnosed as low grade gliomas and both cases on CT only.

The maximum radiological wrong diagnosis occurred in both High grade gliomas and meningioma patients among group I patients. Low grade gliomas occurring often in Group I and tuberculomas in Group II and all the frequent radiological wrong diagnosis occurred in Group II, with the same frequency as in group I.

Secondaries were reported in two cases one each in high grade glioma and Meningioma group among group I patients.

Among radiologically diagnosed meningiomas, tuberculoma occurred in two cases followed by schwannoma, secondaries, high grade glioma and low grade glioma once each. Interestingly both the tuberculomas were en plaque varieties, when both the radiological diagnosis were en plaque meningiomas.

Abscess and schwannoma occurred once each as pathological diagnosis when radiological diagnosis of the former was High grade glioma and the latter was meningioma. The latter especially belonging to Group II.

Low grade glioma occurred as pathological diagnosis for both radiologically diagnosed ependymoma and medulloblastoma with no statistical significance.

Finally considering correlation among CT and MRI, the radiological diagnosis differed only once among 67 patients, (Case No.7) when MRI diagnosed High grade glioma and CT appearance resembled a meningioma. But, Pathologically it was a high grade glioma. The measure of agreement between CT and MRI as far as radiological diagnosis was concerned was 98.50, which showed not much of difference between the two common modes of investigations available, although CT has high affordability when compared with MRI.

## CONCLUSION

In all groups MRI had a higher or equal predictive value when compared to CT but statistically not significant.

MRI had a higher predictive value for benign lesions than CT brain. Malignant lesions had more or less equal value for both CT and MRI. Haemangioblastomas occurred as a common pathological correct diagnosis for CT diagnosed low grade glioma cases whereas all MRI diagnosed low grade gliomas were pathologically correct. Hence in CT Brain suggestive of low grade glioma, Haemangioblastoma should be considered as a close differential diagnosis.

Low grade glioma was a common histological diagnosis for all the remaining tumours diagnosed radiologically as high grade glioma, medulloblastoma, ependymoma and Craniopharyngioma except meningioma in both Group I and II patients. So low grade glioma is an important differential diagnosis for all the intrinsic tumours.

Two en plaque meningiomas diagnosed radiologically, both were pathologically proved to be en plaque tuberculomas. Meningioma and high grade glioma were associated with higher number of radiological wrong diagnosis for a variety of lesions occurring in six cases each. So meningioma and high grade gliomas had the least measure of agreement in both the groups

The agreement value between CT and MRI is 98.5%. Although MRI had a higher kappa value than CT, the difference was marginal. When considering the cost, affordability and availability, though CT is slightly inferior to MRI, it is still comparable with MRI as far as pathological diagnostic aspect alone is considered.

## References

1. Adair LB, Ropper AH, Davis KR. Cerebellar hemangioblastoma : computed tomographic, angiographic and clinical correlation in seven cases. *J Comput Tomogr.* 1978 Dec; 2(4):281-94.
2. Altemus LR, Radvany J. Multifocal glioma visualized by contrast enhanced computed tomography : report of a case with pathologic correlation. *J Maine Med Assoc.* 1977 Sep;68(9):324-7.
3. Anderson DR, Falcone S, Bruce JH, Mejidas AA, Post MJ. Radiologic- pathologic correlation. Congenital choroid plexus papillomas *AJNR Am J Neuroradiol.* 1995 Nov-Dec;16(10):2072-6
4. Castillo M, Scatliff JH, Bouldin TW, Suzuki K. Radiologic-pathologic correlation: intracranial astrocytoma. *AJNR Am J Neuroradiol.* 1992 Nov-Dec; 13(6): 1609-16.
5. Centeno RS, Lee AA, Winter J, Barba D. Supratentorial ependymomas. Neuroimaging and clinicopathological correlation. *J Neurosurg.* 1986 Feb;64(2):209-15.
6. Daumas-Duport C, Monsaignon V, Blond S, Munari C, Musolino A, Chodkiewicz JP, Missir O. Serial stereotactic biopsies and CT scan in gliomas: correlative study in 100 astrocytomas, oligo-astrocytomas and oligodendrocytomas. *J Neurooncol.*

- 1987;4(4):317-28.
7. Dewulf P, Demaeral P, Wilms G, Delanote G, Defloor E, Casselman J, Baert AL. Cerebral metastatic malignant melanoma: CT and MR findings with pathological correlation. *J Belge Radiol.* 1993 Oct;76(5):318-9
  8. Earnest F 4th, Kelly PJ, Scheithauer BW, Kall BA, Cascino TL, Ehman RL, Forbes GS, Axley PL. Cerebral astrocytomas: histopathologic correlation of MR and CT contrast enhancement with stereotactic biopsy. *Radiology.* 1988 Mar;166(3):823-7.
  9. Freund M, Hahnel S, Sommer C, Martmann M, Kessling M, Tronnier V, Sartor K. CT and MRI findings in gliomatosis cerebri: a neuroradiologic and neuropathologic review of diffuse infiltrating brain neoplasms. *Eur Radiol.* 2001;11(2):309-16.
  10. Ho VB, Smirniotopoulos JG, Murphy FM, Rushing EJ. Radiologic-pathologic correlation: hemangioblastoma. *AJNR Am J Neuroradiol.* 1992 Sep-Oct;13(5):1343-52.
  11. Iwama T, Yamada H, Sakai N, Andoh T, Nakashima T, Hirata T, Funakoshi T. Correlation between magnetic resonance imaging and histopathology of intracranial glioma. *Neuro Res.* 1991 Mar;13(1):48-54.
  12. Koeller KK, Smirniotopoulos JG, Jones RV. Primary central nervous system lymphoma: radiologic-pathologic correlation. *Radiographics.* 1997 Nov-Dec;17(6):1497-526.
  13. Koeller KK, Sandberg GD; Armed Forces Institute of Pathology. From the archives of the AFIP. Cerebral intraventricular neoplasms: radiologic-pathologic correlation. *Radiographics* 2002 Nov-Dec;22(6):1473-505.
  14. Lee YY, Castillo M, Nauert C, Moser RP. Computed tomography of gliosarcoma. *AJNR Am J Neuroradiol.* 1985 Jul-Aug;6(4):527-31.
  15. Melikian AG, Golubev VI, Lobanov SK. Correlation of the histological structure of gliomas with densitometry findings during computer tomography *Zh Vopr Neurokhir Im NN Burdenko.* 1984 Mar-Apr; (2):3-9.
  16. Messina AV. Cranial computerized tomography. A radiologic-pathologic correlation. *Arch Neurol.* 1977 Oct;34(10):602-7.
  17. Morota N, Sakamoto K, Kobayashi N, Hashimoto K. Recurrent low-grade glioma in children with special reference to computed tomography findings and pathological changes. *Childs Nerv Syst.* 1990 May;6(3):155-60.
  18. Mori H, Lu CH, Chiu LC, Cancilla PA, Christie JA. Reliability of computed tomography: correlation with neuropathologic findings. *AJR Am J Roentgenol.* 1977 May;128(5):795-8.
  19. Munari C, Musolino A, Daumas-Duport C, Missir O, Brunet P, Giallonardo AT, Chodkiewicz JP, Bancaud J. Correlation between stereo-EEG, CT-scan and stereotactic biopsy data in epileptic patients with low-grade gliomas. *Appl Neurophysiol.* 1985;48(1-6):448-53.
  20. Oi S, Wetzel N, Kim KS. Gliomas in computerized axial tomography. Correlation with tumor malignancy in 100 cases. *No Shinkei Geka.* 1979 Aug;7(8):759-63.
  21. Pierallini A, Bonamini M, Pantano P, Palmeggiani F, Raguso M, Osti MF, Anaveri G, Bozzao L. Radiological assessment of necrosis in glioblastoma: variability and prognostic value. *Neuroradiology.* 1998 Mar;40(3):150-3.
  22. Pigeau, I Sigal R, Halimi P, Comoy J, Doyon D. MRI features of craniopharyngiomas at 1.5 Tesla. A series of 13 cases. *Neuroradiol* 1988;15(3):276-87.
  23. Rao KC, Levine H, Itani A, Sajor E, Robinson W. CT findings in multicentric glioblastoma: diagnostic-pathologic correlation. *J Comput Tomogr.* 1980 Sep;4(3):187-92.
  24. Rees JH, Smirniotopoulos JG, Jones RV, Wong K. Glioblastoma multiforme: radiologic-pathologic correlation. *Radiographics.* 1996 Nov;16(6):1413-38; quiz 1462-3.
  25. Russell EJ, George AE, Kricheff II, Budzilovich G. Atypical computed tomography features of intracranial meningioma: radiological-pathological correlation in a series of 131 consecutive cases. *Radiology.* 1980 Jun;135(3):673-82.
  26. Selker RG, Mendelow H, Walker M, Sheptak PE, Phillips JG. Pathological correlation of CT ring in recurrent, previously treated gliomas. *Surg Neurol.* 1982 Apr;17(4):251-4.
  27. Sundaram C, Rammurti S, Reddy JJ, Prasad SS, Purohit AK. Haemangioblastoma: a study of radiopathologic correlation. *Neurol India.* 2003 Sep;51(3):373-5.
  28. Takeda N, Tanaka R, Yamazaki K. Correlation of computed tomography with post-mortem histopathology of cerebral malignant glioma (author's transl) *Neurol Med Chir (Tokyo).* 1980 Jun;20(6):603-11.
  29. Wu RH, Lang ZJ, Du CC. Pathological studies of CT findings in supratentorial astrocytoma. Correlation between low-density lesions and changes in fine structure. *Chin Med J (Engl).* 1991 Aug;104(8):685-92.
  30. Youmans neurological surgery 5th Edition- H. Richard Winn. Saunders -2004.
  31. Neurosurgery - Robert H. Wilkins and Setti S. Rengachary II edition. Megran Hill 1996
  32. Neuroimaging William W Orrison WB Saunders Company 1998.
  33. Neuroradiology 3rd edition by Juan M. Taveras Williams and Wilkins 1996.