



A STUDY OF ANATOMICAL VARIATIONS OF THE FIRST EXTENSOR COMPARTMENT IN RELATION TO DE QUERVAIN'S DISEASE.

Dr. Tarkeshwar Golghate

Professor And HOD Of The Department Of Anatomy, Gmc, Chandrapur.

Dr. Shruti Mamidwar*

Associate Professor In The Department Of Anatomy, Gmc, Chandrapur *Corresponding Author

ABSTRACT

AIMS: To examine the incidence and length of the first extensor compartment septum.

MATERIALS AND METHODS: Forty sides of the wrists in 20 cadavers were used. The presence of a septum in the first extensor compartment was examined. The septum length was recorded with the radial styloid process as the reference point.

OBSERVATION AND RESULTS: The anatomical variations of the first extensor compartment were classified into 3 types. Type I compartment was found in 7 sides in males (29.2%) versus 6 sides in females (37.5%, $P = .733$), type II was found in 6 sides in males (25%) versus 1 side in females (6.25%, $P = .21$), and type III was found in 11 sides in males (45.8%) versus 9 sides in females (56.25%, $P = .56$). There was no significant difference in the septum length between males and females (5.3 ± 2.3 vs 4.8 ± 1.1 mm, $P = .54$).

KEYWORDS : de Quervain disease, first extensor compartment

INTRODUCTION:

De Quervain disease is the most common disease involving the hand extensor tendons, with a general incidence of 0.5% in males and 1.3% in females.^[1] The pathological examination of de Quervain disease, although also known as de Quervain tenosynovitis, shows no signs of inflammation, but signs of degenerative changes, such as mucoid degeneration, fibrocartilage metaplasia, mucopolysaccharide deposition, and neovascularization.^[2] These pathological changes restricted the sliding of APL and EPB tendons in the first extensor compartment, which results in pain.^[3]

Understanding the anatomy of the first extensor compartment is essential for successful treatment of de Quervain disease.^[4] The anatomical variations of the first extensor compartment include the number of the APL tendon slips and the presence of a septum. Multiple APL tendon slips ranging from 3 to 14 have been reported in up to 89% subjects.^[5,6] The previously reported incidence of the first extensor compartment septum ranged from 34.6% to 72%, which may completely or incompletely divide the compartment into 2 subcompartments.^[7-14] The septum has been classified into the complete type and the incomplete type.^[15] It has been shown that the septum plays an important role in the development of de Quervain disease.^[6]

Corticosteroid injection is the first line treatment for de Quervain disease with an initial response rate ranging from 50% to 83%.^[16-18] However, treatment failure may occur in 14% to 34.5% patients.^[16]

Our study aimed to examine the incidence of first extensor compartment septum and its length. The anatomical variations of the first extensor compartment were classified and its clinical implications were discussed.

MATERIALS AND METHODS:

Twenty adult cadavers were studied in the Department of Anatomy in a tertiary medical college in central India from 1st January 2017 to 31st December 2017. The cadavers were preserved in 10% formalin. There were 12 males (24 sides) and 8 females (16 sides). The wrists of all cadavers showed intact skins and no signs of previous trauma or surgery.

The cadaver was put into the supine position with both arms lying on the sides of the body. The thumb was upwards with the arm in the neutral position. The wrist skin, subcutaneous tissues, the cephalic vein and its tributaries, and the superficial branch of the radial nerve and its branches were removed. The dissection area was from the proximal interphalangeal joint of the thumb to 10 cm proximal to the wrist. Caution was used when dissecting and exposing the retinaculum of the first extensor compartment. Dissection was carried out along the APL tendon, then the APL tendon was retracted to the ulnar side, and

the presence of a septum was examined. Upon presence of a septum, the most distal end of the radial styloid process was exposed. The septum length was defined as the distance from the proximal end of the septum to the distal end of the radial styloid process. The measurement was performed using a digital caliper (accuracy 0.01 mm).

Categorical data were compared using the Fisher exact test and continuous data were compared using the Student t test. A P value less than .05 was considered statistically significant.

RESULTS:

We observed three types of anatomical variations of the first extensor compartment. There are no significant differences in the septum incidence between males and females (7/24 sides, 29.2% vs 6/16 sides, 37.5%; $P = .733$) (Table 1). The septum incidence also did not differ significantly between the left and right sides (Table 2).

There are no significant differences in the septum incidence between males and females (6/24 sides, 25% vs 1/16 sides, 6.25%; $P = .210$) (Table 1). The septum incidence also did not differ significantly between the left and right sides (Table 2).

The type III compartment contains a septum proximal to the radial styloid process, which separates the APL and the EPB tendons (Fig. 1). There are no significant differences in the septum incidence between males and females (11/24 sides, 45.8% vs 9/16 sides, 56.25%; $P = .563$) (Table 1). The septum incidence also did not differ significantly between the left and right sides (Table 2).

In our 20 cadavers, dual-sided septa were found in 11 (55%) cadavers, single-sided septum was found in 5 (25%) cadavers, and no septum was found in 4 (20%) cadavers. The overall septum incidence among the 40 wrists was 67.5% (27/40 wrists).

DISCUSSION:

Our study identified 3 types of anatomical variations of the first extensor compartment. The type I compartment has no septum and the APL and the EPB tendons are contained in the same tendon sheath. The type II compartment contains a septum and its proximal end is distal to the radial styloid process, which separates the APL and the EPB tendons. The type III compartment contains a septum and its proximal end is proximal to the radial styloid process, which separates the APL and the EPB tendons. The type I and II compartments are theoretically at a lower risk of de Quervain disease, and may have better treatment efficacy than the type III compartment. The septum has been classified into the complete type and the incomplete type.^[15] We speculate that the length of the septum is more meaningful for clinical treatment.

In our 20 cadavers, dual-sided septa were found in 11 (55%) cadavers, which is higher than that of the single-sided septum in 5 (25%)

cadavers. The overall septum incidence among the 40 wrists was 67.5% (27/40 wrists).

Our study showed that the proximal end of the septum is 5 mm proximal to the radial styloid process and has no significant differences between different sexes and sides. Therefore, we propose that injection at 5 mm from the radial styloid process may increase the injection accuracy and treatment efficacy without concerning the presence of the septum.

Failure in recognizing the subcompartments in the first extensor compartment may lead to inadequate decompression, and finally resulting in undesirable treatment effects or symptom relapse.[14]The multiple APL tendon slips may be mistakenly recognized as the EPB tendon.[6] This can result in missed decompression of the EPB tendon, which is in another subcompartment. During operation, the presence of a subcompartment can be examined by retracting a tendon and observing the movement of the corresponding finger. Retracting the EPB tendon extends the metacarpophalangeal joint of the thumb, and retracting the APL tendon abducts the first metacarpal. It is also important to note that the presence and length of the septum are not bilaterally symmetrical. Our findings suggest that exploration from 5 mm proximal to the radial styloid process is useful in identifying the septum and subcompartments. By doing so, it is possible to fully release the first extensor compartment and improve surgical efficacy while minimizing the incision.

There are some limitations in our study. First, the cadavers were fixed using formalin solution, which may cause differences in the measurement of the septum length between the cadavers and physiological conditions. Second, removal of the skin and subcutaneous tissues may also make the measurement results smaller. Finally, the medical history of previous de Quervain disease of the cadavers was not clear.

Conclusion Our study identified 3 types of the anatomical variations of the first extensor compartment. The type III variation is theoretically associated with the development of de Quervain disease. The mean septum length is 5 mm, which is a useful reference data for injection treatment and surgical exploration.


TABLE 1. Comparison of the first extensor compartment between males and females.

	Males (n=24)	Females (n=16)	P
TYPE I	7 (29.2%)	6 (37.5%)	0.733
TYPE II	6 (25%)	1 (6.25%)	0.210
TYPE III	11 (45.8%)	9 (56.25%)	0.563

2. Comparison of the first extensor compartment between the left and right sides.

	Left sides (n=20)	Right sides (n=20)	P
TYPE I	7 (35%)	6 (30%)	0.975
TYPE II	3 (15%)	4 (20%)	0.677
TYPE III	10 (50%)	10 (50%)	1.0

Fig 1. 3 types of the anatomical variations in the first extensor compartment.



REFERENCES:

- Hadianfard M, Ashraf A, Fakheri M, et al. Efficacy of acupuncture versus local methylprednisolone acetate injection in De Quervain's tenosynovitis: a randomized controlled trial. *J Acupunct Meridian Stud* 2014;7:115-21.
- Knobloch K, Gohritz A, Spies M, et al. Neovascularisation in de Quervain's disease of the wrist: novel combined therapy using sclerosing therapy with polidocanol and eccentric training of the forearms and wrists: a pilot report. *Knee Surg Sports Traumatol Arthrosc* 2008;16:803-5.
- Ilyas AM, Ast M, Schaffer AA, et al. De quervain tenosynovitis of the wrist. *J Am Acad Orthop Surg* 2007;15:757-64.
- Lee ZH, Stranix JT, Anzai L, et al. Surgical anatomy of the first extensor compartment: a systematic review and comparison of normal cadavers vs. De Quervain syndrome patients. *J Plast Reconstr Aesthet Surg* 2017;70:127-31.
- Thwin SS, Fazlin F, Than M. Multiple variations of the tendons of the anatomical

- snuffbox. *Singapore Med J* 2014;55:37-40.
- Kulthanan T, Chareonwat B. Variations in abductor pollicis longus and extensor pollicis brevis tendons in the Quervain syndrome: a surgical and anatomical study. *Scand J Plast Reconstr Surg Hand Surg* 2007;41:36-8.
- Bahm J, Szabo Z, Foucher G. The anatomy of de Quervain's disease. A study of operative findings. *Int Orthop* 1995;19:209-11.
- Kay NR. De Quervain's disease. Changing pathology or changing perception? *J Hand Surg Br* 2000;25:65-9.
- Minamikawa Y, Peimer CA, Cox WL, et al. De Quervain's syndrome: surgical and anatomical studies of the fibroosseous canal. *Orthopedics* 1991;14:545-9.
- Kang HJ, Hahn SB, Kim SH, et al. Does endoscopic release of the first extensor compartment have benefits over open release in de Quervain's disease? *J Plast Reconstr Aesthet Surg* 2011;64:1306-11.
- Gonzalez MH, Sohlberg R, Brown A, et al. The first dorsal extensor compartment: an anatomic study. *J Hand Surg Am* 1995;20:657-60.
- Mahakkanukrauh P, Mahakkanukrauh C. Incidence of a septum in the first dorsal compartment and its effects on therapy of de Quervain's disease. *Clin Anat* 2000;13:195-8.
- Leslie BM, Ericson WB, Jr, Morehead JR. Incidence of a septum within the first dorsal compartment of the wrist. *J Hand Surg Am* 1990;15:88-91.
- Nayak SR, Hussein M, Krishnamurthy A, et al. Variation and clinical significance of extensor pollicis brevis: a study in South Indian cadavers. *Chang Gung Med J* 2009;32:600-4.
- Gurses IA, Coskun O, Gayretli O, et al. The anatomy of the fibrous and osseous components of the first extensor compartment of the wrist: a cadaveric study. *Surg Radiol Anat* 2015;37:773-7.
- Richie CA3rd, Briner WW., Jr Corticosteroid injection for treatment of de Quervain's tenosynovitis: a pooled quantitative literature evaluation. *J Am Board Fam Pract* 2003;16:102-6.
- Sawaizumi T, Nanno M, Ito H. De Quervain's disease: efficacy of intra-sheath triamcinolone injection. *Int Orthop* 2007;31:265-8.
- Valen PA, Foxworth J. Evidence supporting the use of physical modalities in the treatment of upper extremity musculoskeletal conditions. *Curr Opin Rheumatol* 2010;22:194-204.