



RENAL MORPHOLOGICAL CHANGES IN HEALTHY SENIOR CITIZENS

Dr ANV Prasad	MD, DMRD, Associate Professor, NRI Institute of Medical Sciences, Visakhapatnam, Andhra Pradesh, India
Dr Bikash Kumar Singh*	MD, Assistant Professor, Radio-Diagnosis NRI Institute of Medical Sciences, Visakhapatnam, Andhra Pradesh, India *Corresponding Author
Brig Dr Hariqbal Singh	MD, DMRD Director (Hospital Administration) and Professor & HOD, Radio-Diagnosis NRI Institute of Medical Sciences, Visakhapatnam, Andhra Pradesh, India

ABSTRACT This study of 100 healthy senior citizens (SC) screened out of 10,000, who attended the free medical camps. Renal morphological parameters were assessed sonographically and correlated with age and weight of SC. The result showed significant changes after the age of 70 years.

Aim: A retrospective study to evaluate the sonographic values of renal length, parenchymal and cortical thickness in healthy senior citizens and to assess if renal size significantly decreases with age in healthy senior citizens free from life style diseases and those affecting the renals.

KEYWORDS : Healthy Senior Citizens (SC), Ultrasound, Renal Morphology

INTRODUCTION:

The renals undergo involution with the age. There is a gradual decline in the renal volume, weight and function after the age of 60 years. The most marked decrease occurring between the 7th and 8th decade. The fundamental alterations of renal morphology in the SC include size reduction, parenchymal thickness reduction and margin irregularities. In particular, the most important morphologic alteration of the aged renal is the volume reduction, approximately 20%-30% in above 80 years and a loss of weight that decreases from 250-270 gm to 180-200 gm after age of 65 years¹. The progressive loss of renal mass appears to affect the renal cortex more than the renal medulla. Due to the progressive increase in the mean age of global population it is very important to know the renal morphologic changes according to the age. The functional alterations of the aged renals are characterized principally by a progressive reduction of renal blood flow from about 600 to 300 mL/min/1.73 m², and of glomerular filtration rate (GFR) from 130 mL/min to 60-80 mL/min.²

MATERIALS AND METHODS:

Regular free medical camps for SC were conducted in the Institute of Geriatrics, a social wing of Sri Govardhanagiri Medical Foundation, Visakhapatnam established in 1998. Camps were conducted on last Sunday of every month since 2007. On average 100 SC attended the camp of which 40% were new every month. Thus the health check services were provided to more than 10,000 SC including treatment for various chronic diseases like asthma, arthritis, diabetes, hypertension, cardiac and renal.

For this study screening data of 100 healthy SC who were free from all diseases including the above mentioned entities with normal creatinine levels were analyzed. They were divided in 3 groups.

- 1). Group 1. 60-69 years (young SC)
- 2). Group 2. 70-79 years (middle SC)
- 3). Group 3. 80 years and above (grand SC)

The sonographic renal parameters in different age groups of SC were displayed in Table 1 and 2. The mean and standard deviations of renal parameters were shown in Table 2. Renal morphological changes in SC were most marked after the age of 80 years.¹² For proper understanding and interpretation of results the sonographic and pathological changes were shown in Table 3.

RESULTS:

The total number of SC studied was 100, amongst which 60 were males and 40 were females. The numbers of SC in group 1 were 48, in group 2 were 38 and in group 3 were 22. The mean renal length on right was 95.05mm and on left was 97.38mm. There was progressive reduction in renal length with increase in age with mean value of 3.9mm in group 3. The mean parenchymal thickness was 15.2mm with reduction of 2.6mm in group 3 and the mean cortical thickness was 9.02mm with

reduction of 1.8mm in group 3. The renal parameters in relation to the age of SC are shown in Figs 1 and 2, in relation to body weight of SC are shown in Figs 3 and 4. The figures showed reduced parameters in SC with low body weight and advanced age.

TABLE 1: Renal Parameters in different age groups

Parameter in mm	60-69 (n =48)	70-79 (n = 38)	80 and above (n = 22)
Right renal length	96.3	94.4	93
Left renal length	99.1	96.5	94.6
Parenchymal Thickness	15.8	15.6	13.2
Cortical Thickness	9.6	8.9	7.8
SC's Body Weight Kg	58.7	57.4	55.9

TABLE 2: Descriptive Statistics

	N	Range	Minimum	Maximum	Mean		Std. Deviation
					Statistic	Std. Error	
AGE	100	24	60	84	70.56	.716	7.161
Weight	100	25	45	70	57.76	.582	5.822
Right	100	30	80	110	95.05	.636	6.359
Left	100	30	85	115	97.38	.733	7.329
Cortical Thickness	100	8	5	13	9.02	.160	1.595
Parenchymal Thickness	100	10	9	19	15.21	.225	2.249


TABLE 3: Renal morphological changes in SC after 80 years of age¹²

a. Measurements	* Gradual decline in volume of 20-30% *Weight is reduced by 50-150 gm (Normal weight 250-270 gm).
b. Sonographic	*Length reduction is variable *Parenchymal and cortical thickness is significantly reduced, cortical echoes variable. *Renal capsule may be thickened, renal sinus fatty echo is increased. * Retention cysts and renal sinus cysts are seen frequently *Renal margin can have irregularity with pseudo lobular appearance due to cortical scars of infarcts and infections. *Doppler shows reduced sub capsular and cortical perfusion.
c. Microscopic	*Glomeruli become shrunken due to progressive sclerosis, hyalinization, hyper filtration and hyper perfusion. *Reduction in number of glomeruli is not consistent.

d. Electron microscopic	<p>*The loss of glomeruli mass is proportional to the loss of tubular mass to preserve the balance.</p> <p>*Outer cortical glomeruli are more extensively involved than the deeper.</p> <p>*Thickening of basement membrane of glomeruli and distal tubule due to type-4 collagen deposition</p> <p>Glomerular sclerosis, interstitial fibrosis and tubular atrophy</p>
e. Renal vascular	<p>* Renal arteries show increased tortuosity, tapering and obliteration of interlobular arteries, intimal thickening, shunts between afferent and efferent arterioles</p> <p>* Vascular changes due to aging are difficult to differentiate from changes due to hypertension</p>
f. Functional; GFR and renal blood flow	<p>* Progressive reduction of renal blood flow from about 600 to 300 mL/min/1.73 m²,</p> <p>* Glomerular filtration rate (GFR) from 130 mL/min to 60-80 mL/min.</p>

Society of Radiology, electronic presentation online system ECR 2010 c-1351

- LucaZanoli, et al. Renal function and ultrasound imaging in elderly subjects The scientific world journal, volume 2014, Article ID 830649
- Wael El et al Sonographic Assessment of Renal Size in Healthy Adults Med Princ Pract 2014; 23:432–436DOI:10.1159/000364876
- Glasscock RJ, Rule AD. The implication of anatomical and functional changes of the aging kidney: with an emphasis on the glomeruli .Kidney Int 2012; 82:270-7
- Libório A.B et al Quantitative Renal Echogenicity as a Tool for Diagnosis of Advanced Chronic Kidney Disease in Patients with Glomerulopathies and no Liver Disease Kidney Blood Press Res 2017;42:708–716.



DISCUSSION:

Measure what is measurable, and make measurable what is not so (Galileo 1564-1642).

Renal size was conventionally determined on X-ray KUB or on urography with various drawbacks in measuring the renal length relative to the distance of lumbar vertebrae from L1 to L3/4 as renal hilum usually corresponds to L2. With the advent of newer modalities in radiology renal morphology is effectively illustrated by ultrasonography, computed tomography and MRI. Renal volume is the most precise indicator of renal size. However volume assessment is not used clinically because its measurement is difficult due to the complex renal shape. Many factors like body mass index, height, gender, age, position of renal and renal vasculature influence the measurements.³

Renal Length: Renal length decreases with age especially after 70 years because of parenchymal reduction due to reduced perfusion related to nephroangiosclerosis.⁴ There is progressive reduction in renal length with mean value of 3.9 mm in different groups of SC in this study (Table 1).

Renal parenchymal and cortical thickness: The mean parenchymal thickness is 15.2 mm and cortical thickness is 9.02 mm. There is significant change in parenchymal and cortical thickness with age and weight of SC (Table 2). Renal cortical echogenicity is variable from hypoechoic, isoechoic and even slightly hyperechoic normally. Renal echogenicity is assessed qualitatively by operator and is unreliable. Quantitative assessment of renal cortical echo with that of liver with advanced software by calculating the pixel densities will avoid the subjective variability.⁵

CONCLUSION:

The most marked involution changes due to ageing are seen after the age of 80 years, which may not have any particular clinical impact. The data available on the relationship between renal morphology and function in SC is inadequate. Quantitative assessment of renal cortical echogenicity and renal perfusion studies by ultrasound may improve the ability of prediction to discriminate renal changes in healthy SC in future.⁵

REFERENCES

- Martingano et al. Renal imaging in the elderly. Normal findings and pathology European