



## STUDY OF PULMONARY FUNCTION TESTS IN TRAFFIC POLICEMEN EXPOSED TO AUTOMOBILE POLLUTION IN PATNA

**Manish Ranjan\***

Tutor, Department of Physiology, Nalanda Medical College, Patna \*Corresponding Author

**Kiran Narain**

Professor & HOD, Department of physiology, Nalanda Medical College, Patna

### ABSTRACT

**BACKGROUND:** This study was undertaken with a view to investigate the effect of automobile pollution on pulmonary function parameters in Traffic policemen, so that the potential health risks in these subjects can be minimised by early identification of abnormalities in PFT parameters.

**MATERIAL & METHODS:** The study involved the traffic policemen who are exposed to automobile pollution and normal healthy controls who are age & gender matched. In both the groups, PFT parameters like Forced Vital Capacity (FVC), Forced Expiratory Volume in first second (FEV<sub>1</sub>), FEV<sub>1</sub>/FVC ratio, Peak Expiratory Flow Rate (PEFR) and 1/Forced Expiratory Flow between 25-75% of Forced Vital Capacity (FEF) were recorded using Spiro meter RS 232C.

**RESULTS:** Traffic police personnel had significantly declined forced vital capacity (FVC), forced expiratory volume in one second (FEV<sub>1</sub>), slow vital capacity (SVC) and maximum voluntary ventilation (MVV) when compared with predictive normal values, which is probably due to exposure to vehicular exhaust. Traffic personnel with longer duration of exposure showed significantly reduced lung functions than those with shorter duration.

**CONCLUSION:** A significant reduction in Forced Expiratory Volume in first second, Peak Expiratory Flow Rate and FEV<sub>1</sub>/FVC ratio reflected an obstructive type of abnormality in Traffic policemen and there was a linear relationship between duration of exposure.

**KEYWORDS :** Automobile pollution, Flow parameters, Pulmonary Function Tests, Traffic policemen

### INTRODUCTION:

Patna is growing rapidly owing to the expanding economic base. This has led to an increase in the use of motor vehicles with a subsequent rise in the levels of air pollution. Among the motor vehicle generated air pollutants, highly significant percentage of the air particles are emitted by diesel exhaust. Bus drivers are consistently exposed to diesel exhaust and other atmospheric pollutants.<sup>1</sup>

The air quality index shows that Patna is moderately polluted in all the seasons.<sup>2</sup> Increased number of vehicles result in slow traffic movement which is further compounded by acceleration and de-acceleration of the vehicles on the congested roads. This leads to increased air pollution in the city. Pollutants may be changed by chemical reactions after being emitted. Oxides of nitrogen and volatile organic compounds from automobile exhaust react with sunlight and produce ozone.<sup>3</sup>

Acute effects of the exhaust exposure include irritation of eyes and nose, headache, fatigue, and nausea.<sup>4</sup> Chronic exposure is associated with cough, sputum production, backache, increased stress<sup>5</sup> and decrease in lung function.<sup>6</sup> The most important traffic-related air pollutant is particulate matter (PM). Fine particles of particulate matter (PM 2.5) can reach smaller airways and alveoli leading to decrease in the pulmonary compliance.<sup>7</sup> Pulmonary function tests (PFTs) are an important tool in the investigation and monitoring of patients with respiratory pathology. Spirometry is the most commonly used lung function screening study. It has been observed that exposure to automobile exhaust and fuel vapor impairs lung function in a time dependent manner. If changes in pulmonary functions can be detected in the earliest stages, then preventive or corrective measures are likely to be more beneficial. Thus the present study was aimed to assess ventilatory lung function status of Traffic policeman exposed to automobile pollution in Patna.

### METHODS

The current study was done by taking fifty traffic policemen who are exposed to automobile exhaust and Fifty healthy controls. The subjects were selected from the general population randomly according to following inclusion and exclusion criteria.

### INCLUSION CRITERIA:-

Healthy non smoker traffic policemen in the age group of 20–55 years who are working in traffic junctions for more than 1 year are included in the study.

Healthy non smoker control population of the same age, sex and BSA

are selected from general population and included for the study.

Informed consent is taken from all the subjects those who are willing to participate in the study.

### EXCLUSION CRITERIA:-

Any evidence of chronic obstructive pulmonary disease, asthma, musculoskeletal abnormality, heart disease, anaemia, obesity and history of smoking.

Any history of angina or chest pain, diabetes or hypertension.

Non cooperation or inability to perform pulmonary function tests.

### STUDY DESIGN:

Traffic policemen were selected as per the inclusion criteria laid down. The controls were selected as per the criteria laid down. Their written consent was taken. The screening of the subjects was done for exclusion criteria.

The Anthropometric measurements like height (cm), weight (kg) and Body surface area (square meter) were measured. PFT by computerised spirometer measuring FVC, FEV<sub>1</sub>, FEV<sub>1</sub>/FVC, PEFR and FEF were recorded. 1 | 1 25-75%

The subjects were explained about the actual procedure and purpose of the study before they were put to test. PFT measurements are done by using Kit micro RS 232 C, Computerised spirometer recording, a minimum of 3 readings on each occasion at 15 minute interval and maximum values were taken.

### STATISTICAL ANALYSIS:

Results are presented in Mean ± SD (Min-Max) for each of the parameter. The two groups were compared by using unpaired t test. Analysis of variance (ANOVA) has been used to find the significance of the study parameters between the 3 or more groups of patients.

### RESULTS

Both the subjects and controls were well matched with respect to age, weight, height and body surface area (BSA) respectively.

Table 1 shows the comparison of pulmonary function tests in cases and control groups. It can be seen that there is decline in pulmonary function parameters like FVC, FEV<sub>1</sub>, PEFR, FEV<sub>1</sub>/FVC, FEF in cases when compared to 1 | 1 25-75% controls which is statistically significant.

This study is in agreement with Wongsurakiat P,14 15 Maranetra KN et al , Kim SM, Cheon GR et al , Jafary ZA,1 16 Faridi I et al , Binawara BK, Gahlot S et al and Ingle ST,3 Pachpande BG et al who also found the same results.

Table 2 shows the comparison of pulmonary function tests with respect to duration of exposure. It shows a linear relationship in pulmonary function parameters in traffic policemen with respect to duration of exposure. This study is in agreement with Uzma NB, Khaja17 Mohinnuddin Salar BM, et al who also found the same results.

**Table 2: Comparison Of Pulmonary Function Tests According To Duration Of Exposure**

Pulmonary Function test	1-5 years	6-10 years	11-15 years	>15 years	Control Group	f value	P value
FVC	3.91±0.42	3.32±0.46	3.41±0.63	3.02±0.37	4.09±0.39	46.261	<0.001**
FEV1	3.23±0.46	2.56±0.37	2.55±0.54	2.22±0.26	3.48±0.39	79.989	<0.001**
PEFR	9.03±0.61	7.43±1.19	7.19±1.27	6.87±1.06	8.92±0.86	40.270	<0.001**
FEV1/FVC	0.83±0.05	0.77±0.07	0.74±0.06	0.72±0.06	0.84±0.06	33.088	<0.001**
FEF25-75%	4.22±0.58	3.66±0.45	3.58±0.44	3.03±0.44	4.37±0.49	51.076	<0.001**

Significant at 1% level of significance

## DISCUSSION

In the present to see the effect of automobile exhaust, PFT was recorded in the Traffic policemen. The dynamic parameters like FVC, FEV<sub>1</sub>, PEFR, FEV<sub>1</sub>/FVC, FEF<sub>25-75%</sub> were reduced in traffic policemen exposed to automobile exhaust when compared to control group.

Diesel exhaust organic extracts induce reactive oxygen species in macrophages and bronchial epithelial cells, two key cell types targeted by particulate matter in the lung. Reactive oxygen species activate the promoters of cytokines and chemokines involved in allergic inflammation through activator protein-1 and nuclear factor- kappa B signalling pathways. Organic diesel exhaust particle chemicals also induce apoptosis and necrosis in bronchial epithelial cells via a mitochondrial pathway<sup>18,19</sup>.

Laboratory studies in humans and animals have shown that particulate toxic pollutants particularly diesel exhaust particles can enhance allergic inflammation and can induce allergic immune responses. Most of these immune responses are mediated by the carbon core of diesel exhaust particle. Poly aromatic hydrocarbons are major chemical components of diesel exhaust particulates, and they have enhanced the production of immunoglobulin E<sup>20</sup>.

Diesel exhaust particles (DEP) are thought to consist of a carbon core surrounded by trace metals, such as nickel, and salts to which are adsorbed organic hydrocarbons. A number of these components have inflammatory effects in the lungs of laboratory animals. For example, intratracheal instillation of ultrafine carbon particles in rats leads to neutrophil influx into the lungs, and increase in bronchoalveolar lavage fluid (BALF) concentrations of tumour necrosis factor- (TNF)-alpha. Intratracheal instillation of nickel in rats causes severe and sustained inflammation, with generation of free radicals. Inhalation of hydrocarbons also leads to lung inflammation. The foregoing observations indicate that diesel particles themselves can induce airway inflammation<sup>21,22</sup>.

This may be the reason for decreased FVC and FEV<sub>1</sub> and PEFR in traffic policemen.

FEF<sub>25-75%</sub> indicates flow rates in small airways i.e. those with internal diameters of less than 2 mm. Decrease in FEF<sub>25-75%</sub> suggest greater involvement of small airways. Particles generated from diesel exhaust are extremely small and are present in the nuclei or accumulation modes, with diameters of 0.02 nm and 0.2 nm respectively. These small sized particles, by virtue of their greater surface area to mass ratio, can carry a much larger fraction of toxic compounds, such as hydrocarbons and metals on their surface. Hence chronic exposure to them can lead to chronic inflammation of respiratory tract and lung parenchyma<sup>23,24</sup>.

The FEV<sub>1</sub>/FVC ratio was also reduced in the current study suggesting an obstructive type of respiratory involvement.

The pulmonary function parameters recorded using Kit micro RS 232 C, Computerised spirometer reflected an obstructive type of respiratory involvement in Traffic policemen.

Recording of Slow Vital Capacity would have revealed restrictive

**Table 1: Comparison of Pulmonary Function tests in study and Control group**

Pulmonary Function test	Study group	Control group	t value	P value
FVC	-	4.09±0.39	10.159	<0.001**
FEV1	2.62±0.54	3.48±0.39	12.953	<0.001**
PEFR	7.62±1.32	8.92±0.86	8.294	<0.001**
FEV1/FVC	0.76±0.07	0.84±0.06	8.486	<0.001**
FEF25-75%	3.59±0.63	4.37±0.49	9.652	<0.001**

\*\* Significant at 1% level of significance

abnormality more accurately. Post Bronchodilator test along with FEV<sub>1</sub>/FVC would have supported obstructive type of abnormality.

The PFT parameters in the Traffic policemen showed a linear relationship with respect to duration of exposure in the present study.

The chronic inflammation of respiratory tract may be the reason for decreased pulmonary function parameters in traffic policemen with respect to duration of exposure.

## CONCLUSIONS

The results of this study have shown a significant fall in pulmonary function parameters like FVC, FEV<sub>1</sub>, PEFR, FEV<sub>1</sub>/FVC and FEF<sub>25-75%</sub> in traffic policemen and the fall in pulmonary function parameters is also associated with respect to duration of exposure indicating a reduction in lung function efficiency of the traffic policemen exposed to automobile pollution.

The significant reduction in PEFR and FEV<sub>1</sub>/FVC indicate warning symptoms of asthma among Traffic policemen. This study concludes that the Traffic policemen are highly vulnerable for respiratory impairment due to vehicular exhaust at workplace environment. The protective measures like use of mask, regular health check up and also awareness on health impacts of pollution need to be adopted for protection of Traffic policemen working on the heavy traffic roads.

## REFERENCES

- Jafary ZA, Faridi IA, Qureshi HJ. Effects of airborne dust on lung function of the exposed subjects. Pak J Physiol 2007; 3(1): 30-34.
- Zhu Y, Hinds WC, Kim S, Sioutas C. Concentration and size distribution of ultrafine particles near a major highway. J Air Waste Manag Assoc 2002; 52(9): 1032-42.
- Ingle ST, Pachpande BG, Wagh ND, Pate VS, Attarde SB. "Exposure to vehicular pollution and respiratory impairment of traffic policemen in Jalgaon city, India". Industrial health 2005; 43: 656-652
- Nielsen OR, Nielsen ML, Gehl J. Traffic-related air pollution: Exposure and health effects in Copenhagen street cleaners and cemetery workers. Arch Environ Health 1995; 50(3):207-13.
- Chattrjee BP, Alam J, Gangopadhyay PK. A study of dynamic lung function in jute mill workers. Indian J Indus Med 1989; 35, 157-65.
- Chattopadhyaya BP, Dipali S, Satipati C. Pulmonary function test and the jute mill workers. Indian J Occup Health 1994; 37, 1-10.
- Taggart SC. Asthmatic bronchial hyper responsiveness varies with ambient levels of summertime air pollution. Eur Respir J 1996; 9, 1146-54.
- Rusas I. Analysis of relationship between environmental factors and asthma emergency admissions. Allergy 1998; 53, 394-401.
- Cassino C, Ito K, Bader I, Ciotoli C, Thurston G, Reibman J. Cigarette smoking and ozone associated emergency department use for asthma by adults. Am J Res Cri Care Med 1999; 159, 1773-9.
- English P, Neutra R, Scalf R, Sullivan M, Waller L, Zhu L. Examining association between childhood asthma and traffic flow using geographic information system. Environ Health Perspect 1999; 107, 761-7.
- Stone V. Environmental air pollution. Am J Crit Care Med 2000; 162, S44-7.
- Graham NM. The epidemiology of acute respiratory infections in children and adults: a global prospective. Epidemiol Rev 1990; 12, 149-78.
- Tiittinen P, Timonen KL, Ruuskanen J, Mirme A, Pekkanen J. Fine particulate air pollution, resuspended road dust and respiratory health among symptomatic children. Eur Respir J 1999; 13, 266-73.
- Wongsurakiat P, Maranetra KN, Nana A, Naruman C, Aksornint M, Chalermpanyakorn T. "Respiratory symptoms and pulmonary function of traffic policemen in Thonburi". J Med Assoc Thai 1999 May; 82(5): 435-43.
- Kim SM, Cheon GR, Kim YW, Kim JH, Lee HH, Hong SC, et al. Comparison study of the pulmonary function and serum carboxy hemoglobin level between the traffic policemen and clerk policemen. Tuberc Respir Dis 2003 Dec; 55(6): 560-569.
- Binawara BK, Gahlot S, Mathur KC, Kakwar A, Guptar R, Rajnee. Pulmonary function tests in three wheeler diesel taxi drivers in bikaner city. Pak J Physiol 2010; 6(1):28-31.
- Uzma N, Khaja Mohinnuddin Salar BM, Kumar BS, Aziz N, David MA, Reddy VD. Impact of organic solvents and environmental pollutants on the physiological function

- in petrol filling workers. *Int J. Environ. Res. Public Health* 2008; 5(3): 139-146
18. Dickson RP, Schwartz DA. Acute and chronic responses to toxic inhalations. In: Fishman AP, Elias JA, Fishman JA, Grippi MA, Senior RM, Pack AI, editors. *Fishman's pulmonary diseases and disorders*. 4th edition. China: Mc Graw Hill companies; 2008. p.995-1002.
  19. Nel AE, Diaz-Sanchez D, Li N. The role of particulate pollutants in pulmonary inflammation and asthma: evidence for the involvement of organic chemicals and oxidative stress. *curr opin pul med* 2001 Jan; 7(1): 20-6.
  20. Polosa R, Salvi S, Di Maria GU. Allergic susceptibility associated with diesel exhaust particle exposure: clear as mud. *Arch Environ Health* 2002 May-Jun; 57(3): 188-93.
  21. Inoue KI, Takano H, Yanagisawa R, Ichinose T, Shimada A, Yoshikawa T. Pulmonary exposure to diesel exhaust particles induces airway inflammation and cytokine expression in NC/Nga mice. *Archives of Toxicology* 2006; 79(10):595-599
  22. Nightingale JA, Maggs R, Cullinan P, Donnelly, Rogers DF, Kinnersley R, et al. Airway Inflammation after Controlled Exposure to Diesel Exhaust Particulate. *Am. J. Respir. Crit. Care Med* 2000; 162(1): 161-166
  23. Singhal MK, Khaliq F, Singhal S, Tandon OP. Pulmonary functions in petrol pump workers: a preliminary study. *Indian J Physiol Pharmacol* 2007; 51(3): 244-248
  24. Levsen K. The analysis of diesel particulate. *Fresenius Z Anal Chem* 1988; 331: 467-478.