



VASCULAR COMPLICATION DUE TO THE USE OF CALCIUM HYDROXYAPATITE: CASE REPORT

| | |
|---|---|
| Dr. Gicelio Nobrega | Physician |
| Dr. Honório Sampaio Menezes | Physician, MD |
| Dr. Victor Hugo Rabello | Physician |
| Dr. Roberto Chacur* | Physician, Specialist *Corresponding Author |
| Karina Alcantra Silva | Medical Student |
| Anny Carolina de Melo Guedes | Medical Student |
| Sayonara Bruna Mesquita | Medical Student |
| Nivea Maria Bordin da Silva Chacur | Physician |
| Dr. Rodrigo Cadore Mafaldo | Physician, Specialist |
| Dra. Danuza Dias Alves | Physician |
| Dr. Leandro Dias Gomes | Physician, MD |
| Dra. Fernanda Bertolo | Physician |
| Giulia Casa Nova Barbosa | Biomedical student |

ABSTRACT

Introduction: Filler injection for augmentation and facial filling is a cosmetic procedure that has been increasingly common in recent decades.

Objective: This work aims to report a patient's case.

Methodology: Case report of a complication after an aesthetic procedure performed using calcium hydroxyapatite. Patient was a 27-year-old female who complained of ischemia in her upper lip.

Conclusion: Among the complications of filling procedures, the one presented in the case, tissue necrosis due to loss of blood circulation, is the most serious. The treatment used, which included hyperbaric chamber, reduced the ischemic area.

KEYWORDS : Hydroxyapatite, Vascular complication, Calcium Hydroxyapatite injection, necrosis, HBO².

INTRODUCTION

Filler injection for facial volume augmentation is a cosmetic procedure that has been increasingly common in recent decades.^{1,2,3}

Because these types of fillings are non-surgical, relatively easy to be carried out, and present rapid results, in recent years, an increasing number of complications have been related to them, including allergic reaction, granuloma formation, skin necrosis or cellulite, as well as arterial occlusion or embolization.^{1,4,5,6}

Calcium hydroxyapatite (CaHA) is a semi-permanent soft tissue filling substance which can last from one to two years in tissues. It is used mainly to treat facial depressions and asymmetries, in addition to correcting post-rhinoplasty deficiencies.¹

Injectable soft tissue fillers provide patients with immediate results without the downtime associated with traditional facial rejuvenation surgery.

Mild pain, swelling, and bruising at the injection site are common and expected consequences of these treatments. Real complications develop less frequently.^{2,6}

In the present case report, the results of a serious complication after an aesthetic procedure which injected calcium hydroxyapatite into a patient's upper lip is described.

CASE REPORT

A 27-year-old single female, born and resident in João Pessoa (Paraíba,

Brazil), came to the clinic with a complaint about her upper lip. The initial medical evaluation on September 09, 2019 indicated the presence of devitalized tissues, an area of necrosis, and moderate exudation in the patient's upper lip, as well as perilesional erythema, all resulting from a complication of a filling procedure with calcium hydroxyapatite.

Treatment was carried out using aseptic techniques: cleaning with polyhexamethylenebiguanide (PHMB) solution and antiseptic soap, use of non-adherent mesh embedded with petrolatum emulsion, sterile antimicrobial gauze dressing embedded with PHMB, and sticking plaster made of elastic polyester.

Treatment with hyperbaric oxygen therapy started on the same day of the first consultation, being followed by debridement between sessions. These presented significant improvement in the necrotic area. Dressings were performed daily due to the patient's need for hyperbaric oxygen therapy and to her clinical conditions. Tramadol, amoxicillin with clavulanate, and Ketorolac Trometamol were prescribed. During the medical reassessment after ten daily sessions of hyperbaric chamber, no ischemic process was noticed and the wound was healed. This case report was approved by the Research Ethics Committee of the Veiga de Almeida University (CONEP record).



Figure 1 – Images before and after treatment.

DISCUSSION

Even when administered by experienced professionals, soft tissue fillers can cause various reactions. These may range from small and self-limited responses to serious complications that require immediate treatment and follow-up.^{2,6}

Several fillings have gained popularity due to their fast and predictable results.

Over the years, as there has been an increasing demand for these types of procedures, the group of people who can provide these techniques has expanded from cosmetic surgeons and dermatologists to include health professionals with little or no training in aesthetic medicine. Even when administered by experienced professionals, the use of fillers can cause several unwanted reactions.^{2,3}

Calcium hydroxyapatite was the filling material most commonly associated with complications in rhinoplasty.^{2,3}

Cosmetic injections of calcium hydroxyapatite into the nasal bridge can result in arterial embolism of the ophthalmic system. It can also reach the optic, the retinal, and the choroidal nerves, causing severe long-term visual damage and impairment of the visual field.^{1,2,4}

Although the case presented here was not of nasal correction, the mechanism of vascular obstruction was the same as that presented in cases of nasal correction.

CONCLUSION

Tissue necrosis, as presented in this case, although rare, is a devastating accident caused by treatment using calcium hydroxyapatite filler. The proposed treatment, which included hyperbaric oxygen therapy, was effective and reduced the local ischemic damage. Doctors should be aware of this possible complication and inform their patients before offering these cosmetic treatments.

REFERENCES

- 1- Cohen, Eyal. "A case report of ophthalmic artery emboli secondary to Calcium Hydroxyapatite filler injection for nose augmentation- long-term outcome." *BMC Ophthalmology*, 2016: 1-6.
- 2- Daines, Steven M. "Complications associated with injectable soft-tissue fillers: A 5-year retrospective review." *JAMA Facial Plastic Surgery*, 2013: 226-231.
- 3- Stupak, Howard D. "Calcium hydroxyapatite gel (radiessse) injection for the correction

of post-rhinoplasty contour deficiencies and asymmetries." *Archives of Facial Plastic Surgery*, 2007: 130-136.

- 4- Ozturk, Cemile Nurdan. "Complications following injection of soft-tissue fillers." *Aesthetic Surgery Journal*, 2013: 862-877.
- 5- Barabas, Reka. "Hydroxyapatite - Carbon nanotube composites for drug delivery applications." *Brazilian Journal of Chemical Engineering*, 2019: 913-922.
- 6- Carruthers, Jean D.A. "Blindness caused by cosmetic filler injection: A review of cause and therapy." *Plastic and Reconstructive Surgery*, 2014: 1197-1201.