



## A STUDY OF PERINATAL MORBIDITY AND MORTALITY IN PRETERM DELIVERY.

**Dr. Ranoji V Shinde\***

Associate professor, Srinivasa medical college , Karnataka. \*Corresponding Author

**Dr. Roopa Shinde**

Associate professor, Srinivasa medical college , Karnataka.

### KEYWORDS :

#### INTRODUCTION

Preterm labour is labour occurring after 28 weeks and prior to 37 completed weeks. Incidence of preterm labour is 5 – 10 % of all pregnancies. Despite great deal of research and introduction of new diagnostic and therapeutic technology the rate of preterm birth has not changed over the past 40 years.

Currently it is one of the most challenging problem confronting the obstetrician and neonatologist as it takes a heavy toll of perinatal morbidity and mortality (contributing between 60 – 70 % of all perinatal deaths in most data series.

The focus of this study is on the perinatal outcome in terms of morbidity and mortality of a preterm baby. It is an important health priority to estimate the risk of adverse perinatal events in preterm births.

The value of perinatal morbidity and mortality study is that it reflects the quality and quantity of health care services available to the mother and the new-born. It reflects the result of maternity care more clearly than the neonatal death rate. Reliable knowledge of neonatal morbidity and mortality is useful antenatally not only to provide women in preterm labour with prognostic guidance but also to guide obstetrician who may be considering elective delivery of a preterm baby.

#### MATERIALS AND METHODS

A Prospective study of patients in preterm labour (after 28 weeks prior to 37 completed weeks) admitted during February 2012 to January 2014 to the obstetrics department of JSS Hospital Mysore Karnataka. The total number of deliveries during the period were 3269

The overall preterm delivery rate was determined after consideration of all causes of spontaneous preterm labour and deliberate intervention to achieve elective preterm delivery

Total number of patients in the study – who fulfilled criteria (Working Criteria)

Patients in spontaneous preterm labour and those patients in whom preterm delivery was electively achieved during gestational period from 28 to 37 weeks who were in true labour was also included in the study

#### Exclusion Criteria:

Patients who were subjected to tocolytic therapy to arrest labour have been excluded from the study and pregnancies before 28 and after 37 completed weeks gestational age were excluded from the study period.

Detailed history was taken including the age, booking, marital status, socioeconomic status, educational, employment status and parity and thorough obstetric examinations considering all the high-risk factors was carried out.

Investigations like routine, urine, vaginal culture sensitivity, C reactive protein, LFT (liver function tests), Kidney function tests), Non-stress test) Ultrasound (for gestational age, liquor, congenital anomaly etc.)

APGAR scores and birth weights of the new born were noted at the time of birth and they were followed up throughout the hospital stay. Incidence and perinatal morbidity and mortality of preterm birth were calculated.

#### RESULT

There were total number of 3269 deliveries in the period of 2 years in the study

The total number of preterm deliveries in the study was 241, the incidence rate of which was 7.4%

Out of which there were 180 cases of spontaneous labour which

**Table 1** shows the various triggering factors for spontaneous preterm labour and also the indications of elective preterm delivery

Triggering factors	Spontaneous		Elective	
	Number of cases	Percentage	Number of cases	Percentage
Preterm premature rupture of membranes	32	17.8	6	9.84
Twin pregnancy	31	17.2	-	-
Pre-eclampsia	20	11.11	19	31.14
Anaemia	15	8.33	-	-
Intra uterine growth retardation	-	-	3	4.91
Antepartum haemorrhage	7	3.9	1	1.60
Congenital anomaly of foetus	6	3.33	8	13.11
Intrauterine foetal demise	5	2.8	10	16.4
Uterine anomaly	4	2.22	-	-
Cervical incompetence	4	2.22	-	-
Cardiac disorders	4	2.22	-	-
Urinary tract infections	3	1.66	-	-
malaria	2	1.11	-	-
Typhoid	1	0.55	-	-
Eclampsia	1	0.55	14	23.0
Idiopathic	45	25	-	-
Total	180	100	61	100

accounted for 74.7% and 61 cases had elective preterm labour which accounts for 25.3%

The Multifetal pregnancies were also included. There were 31 twin pregnancies. So, the total number of babies included in the study were 272

Again, the total number of mortalities in the perinatal period were 82 which accounts for 30% out of which the total number of stillbirths were 28(34%) and early neonatal deaths were 54 (66%)

**DISTRIBUTION OF VARIOUS TRIGGERING FACTOR**

The incidence of preterm labour in our study was 7.4%.

Preterm premature rupture of membranes (17.8%) was the leading causative factor followed by twin pregnancy (17.2%). In our study we observed maximum number of preterm labours occurred in booked cases than un-booked cases, and in age group of 26 to 30 years. Also, preterm labour occurred more in primigravida cases than the multi-gravida

The total number of congenital malformations were seen in 18 babies resulting in incidence of 6.6%. out of which maximum number of cases were of neural tube defect 6 (33.0%)

Among the total number of 272 preterm babies, maximum number of babies were delivered in gestational age from 31-34 weeks were 128 (47%) out of which majority of the babies 101 (37.1%) weighed less than 1500gms the mean weight was 1943gms with standard deviation of 285.2gms

**OCCURRENCE OF PERINATAL MORBIDITY**

Maximum number of morbidity was observed 53 (63.84%) in babies born between 31 to 34 weeks of gestation mostly of which were un-booked cases 47 (56.6%)

**Occurrence of perinatal mortality.**

Total number of babies with mortality were 82 which accounted for incidence of perinatal mortality of 30 % out of which the number of stillbirths were 28 (34%) and the deaths which occurred in the first 7 days of birth were 54 (66%)

Here again Birth asphyxia accounted for maximum number of deaths .14 cases (25.92%). Followed by respiratory distress syndrome accounted for 13 cases (24.07%).

The perinatal mortality was maximum observed 26 (48%) among

babies delivered before 30 weeks of gestation and in babies weighing less than 1500 grams in un-booked cases Thus gestational age and baby birth weight are both inversely proportional to mortality.

**DISCUSSION**

A hospital-based maternity review of preterm deliveries was an appropriate study design to determine the incidence and trend of preterm births and perinatal deaths

There were 3269 deliveries in the study period. The total number of patients in our study is 241. Thus, the incidence of preterm labour in our study was 7.4%.

The occurrence of spontaneous preterm labour in our study is 74.7 % and of elective preterm labour is 25.3%. Idiopathic preterm labour accounts for 45 % of cases in our study. In our study preterm labour was more in primigravida of middle socioeconomic status and was highest in patients belonging to the age group of 20 -25 years (58.1%)

The incidence in developed countries ranges from 3.5 to 7 % . which is comparatively lower than the developing countries which may be due to various reasons like poor antenatal care, screening programmes, lack of health care services for pregnant mothers and also lack good Intensive neonatal care unit are the corner stone in the management of preterm births

**Table 4** Shows The Compares The Preterm Labour Rate With Our Study

Bhargav et al 1984	7 – 8
Kou-Huang Chen et al 2001 – 2011	8.2% to 9.1%
Chukwuemeka Anthony Iyoke, et al 2009 – 2010	16.9
Xiaoqin Zhu, et al 2015 at Huai	4.06
Stacy Beck et al 2005	9.6
Present study	7.4

Study done by Neetu Gupta et all in 2018 showed perinatal morbidity of 67.2 % and perinatal mortality of 52.8%.

Singh et al and Singh Uma et al showed incidence of perinatal mortality around 21% and 12.7% respectively, which is quite less when compared with the present study (incidence of perinatal mortality in our study was 30 %.

In comparesim, to the above studies our study had perinatal morbidity of 30.5 % and perinatal mortality of 30 % which were

**Table 2** Shows The Distribution Of Various Causes Of Perinatal Morbidity

Morbidity	Number of cases	Percentage
Birth asphyxia	29	35
Respiratory distress syndrome	19	23
Septicaemia	16	19.3
Hypothermia	5	6
Hyperbilirubinemia	3	3.6
Apnoea	3	3.6
Hypocalcaemia	2	2.4
Dislocation of hip joint	2	2.4
Meconium aspiration syndrome	1	1.2
Umbilical sepsis	1	1.2
Intraventricular haemorrhage	1	1.2
Patent ductus arteriosus	1	1.2
Total	83	100

**Table 3** Shows The Distribution Of Various Causes Of Perinatal Mortality In The First 7 Days And It Excludes The Still Births.

Causes of death	No. of deaths	Percentage
Birth asphyxia	14	25.92
Respiratory distress syndrome	13	24.07
Septicaemia	11	20.37
Intra ventricular haemorrhage	6	11.11
Meconium aspiration syndrome	3	5.55
Congenital anomaly of foetus	3	5.55
Aspiration	2	3.71
Necrotising enterocolitis	1	1.85
Meningitis	1	1.85
Total	54	100

N=272

in very close range

## CONCLUSION

To conclude it was seen that preterm labour was associated with high occurrence of perinatal morbidity and mortality. The main cause of these were found to be birth asphyxia, respiratory distress syndrome, and septicaemia

Preterm babies are born with a number of physiological handicaps and are thus predisposed to a large number of pathological conditions which needs anticipation and prompt treatment. Prevention of preterm labour is very essential to minimize a perinatal morbidity and mortality. Thus, timely identification of the risk factors and prompt treatment would help to bring down the incidence of preterm labour and hence its perinatal mishap.

And lastly but not least improvement of health care services and a good intensive neonatal care unit are the corner stone in the management of preterm births

## REFERENCES

1. Preterm labour and its effect on perinatal morbidity and mortality Neetu Gupta1, Pragati Divedi, Deepti Dwivedi Int J Reprod Contracept Obstet Gynecol. 2018 May;7(5):1993-1996
2. The trends and associated factors of preterm deliveries from 2001 to 2011 in Taiwan Kou-Huang Chen, PhD I-Chu Chen, MPH, Yi-Chieh Yang, MD, and Kow-Tong Chen, Medicine (Baltimore). 2019 Mar; 98(13): e15060. doi: 10.1097/MD.00000000000015060
3. Ross MG. Preterm labour. University of California. Preterm labour. April, 2011.
4. Pandey K, Ajay B, Neena G, Geetanjal K. Predictive value of various risk factors for preterm labour. J Obstet Gynecol India. 2010;60(2):141-5.
5. Prevalence and perinatal mortality associated with preterm births in a tertiary medical center in South East Nigeria Chukwuemeka Anthony Iyoke, Osaheni Lucky Lawani, Int J Womens Health. 2014; 6: 881–888.
6. Adimora GN, Odetunde IO. Perinatal mortality in University of Nigeria Teaching Hospital (UNTH) Enugu at the end of the last millennium. Niger J Clin Pract. 2007;10(1):19–23. [PubMed] [Google Scholar]
7. High risk pregnancy associated perinatal morbidity and mortality: a second birth population-based survey in Huai'an in 2015 Xiaoqin Zhu, Huiyuan Niu, BMC Pregnancy and Childbirth volume 19, Article number: 224 (2019)
8. Goldenberg RL, Rouse DJ. The prevention of premature birth. N Engl J Med 1998; 339:313–20. [PubMed] [Google Scholar]
9. Tracy SK, Tracy MB, Dean J, et al. Spontaneous preterm birth of liveborn infants in women at low risk in Australia over 10 years: a population-based study. BJOG 2007; 114:731–5. [PubMed] [Google Scholar]