



THYROID DYSFUNCTION AND DYSLIPIDEMIA - RISK FACTORS FOR CARDIOVASCULAR DISEASE IN PCOS WOMEN

Dr Monali Rewatkar*	Assistant Professor, Department of Biochemistry, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India. *Corresponding Author
Dr Kshama Kedar	Associate Professor OBGY Department, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India.
Dr Amrita Singam	Assistant Professor, Pharmacology Department, Indira Gandhi Government Medical College, Nagpur, Maharashtra, India.

ABSTRACT **Background** – Thyroid function regulates a wide array of metabolic parameters. Thyroid function significantly affects lipoprotein metabolism as well as some cardiovascular disease (CVD) risk factors[1,2] A linear increase in total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C) and triglycerides (TGs) and a linear decrease in high-density lipoprotein cholesterol (HDL-C) levels has been observed with increasing TSH [3]. Dyslipidemia is a very common metabolic abnormality in women with polycystic ovary syndrome (PCOS).[4] Polycystic ovary syndrome (PCOS) is widely accepted as the most common endocrine abnormality in women of childbearing age and may be accompanied by dyslipidemia, hyperandrogenism, hyperinsulinemia.[5] Lipid abnormalities, including elevated low-density lipoprotein and triglyceride levels and reduced high-density lipoprotein levels, are often found in women with PCOS and play an important role in PCOS [6] So, aim of this study is to study the correlation between Thyroid hormone dysfunction and dyslipidemia (alteration in lipid profile) which are risk factors for Cardiovascular diseases in PCOS women.

Materials & Methods – This Observational study included about 50 patients attending the outpatient department of ObstGynae department, IGGMC, Nagpur. All PCOS patients who were attending OBGY OPD were selected for study and their blood samples was collected for thyroid hormonal tests ie estimation of T3, T4 & TSH as well as Lipid profile tests. History was taken as per designed proforma and consent form was obtained. Their Blood samples was analyzed for thyroid profile as well as lipid profile test ie Serum cholesterol, Triglycerides, HDL, and LDL & VLDL in clinical Biochemistry Laboratory. Thyroid tests was run on Elisa reader & washer while Serum Cholesterol, Triglycerides, HDL, LDL & VLDL tests was done on Autoanalyser EM 460. Serum values of Thyroid stimulating hormone (TSH), thyroxine (T4) and tri-iodo thyronine (T3) were assayed by ELISA tests and compared with lipid Profile tests and values were compared & correlated. The data was analysed & Student's T-test was used for the calculation. P < 0.05 was considered significant.

Result – Our results shows positive correlation between Serum TSH (Mean=4.63) and serum cholesterol level (Mean= 182) as well as serum Triglycerides, LDL and VLDL level (Mean 163, 100, 32) while TSH shows negative correlation with HDL (Mean=48). This positive correlation was found to be statistically significant between TSH and Serum cholesterol, Triglycerides, LDL & VLDL. Similarly vice versa with T3 and T4 in PCOS women which shows high risk of Cardiovascular disease.

Conclusion – Results of our study indicates how the risk factors in PCOS women for coronary heart disease ie lipid profile affected by alteration in the level of Thyroid hormones in the body. This study showed a significant increase in cholesterol, Triglycerides, VLDL & LDL in patients with hypothyroidism PCOS women. So early screening of PCOS thyroid patients for lipid profile must be advised to prevent future cardiovascular related disorders & its further complications.

KEYWORDS : Lipid Profile(Dyslipidemia), Cardiovascular Risk ,Thyroid dysfunction , PCOS

INTRODUCTION

Polycystic ovary syndrome (PCOS) is a common endocrinological disorder in women of reproductive age characterized by excessive androgen secretion, persistent anovulation and polycystic ovarian morphology [3,4,7]. Studies have revealed that there are different degrees of obesity, dyslipidemia, insulin resistance (IR), oxidative stress and other metabolic abnormalities [4,5,8], and among these abnormalities, dyslipidemia is one of the most common phenomena observed in women with PCOS [3,4,9].

Lipid abnormalities are found in women affected by PCOS. A recent study showed that mild hypercholesterolemia is frequently encountered in women with PCOS [3,4,10]. Different lipid patterns are present in PCOS, including low levels of high-density lipoprotein cholesterol (HDL-C), high triglyceride (TG), total cholesterol (TC) and low-density lipoprotein cholesterol (LDL-C) and significantly higher lipoprotein concentrations [11]. Lipid Profile one of the marker of cardiac risk factor include Cholesterol, Triglycerides, LDL & VLDL, while HDL has protective role in Heart diseases.[12] There is alteration in lipid profile ie dyslipidemia observed in thyroid dysfunction. Thyroid hormones regulate different body metabolism including lipid metabolism.[13] Thyroid hormones induce the 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase, which is the first step in cholesterol biosynthesis. So hyperthyroidism causes increase level of cholesterol. Triiodothyronine (T3) upregulates LDL receptors by controlling the LDL receptor gene activation. Thyroid hormones can influence HDL metabolism by increasing cholesteryl ester transfer protein (CETP) activity, which exchanges cholesteryl esters from HDL2 to the very low density lipoproteins (VLDL) and TGs to the opposite direction [15,16,17]. In addition, thyroid hormones stimulate the lipoprotein lipase (LPL), which catabolizes the TG-rich lipoproteins, and the hepatic lipase (HL),

which hydrolyzes HDL2 to HDL3 and contributes to the conversion of intermediate-density lipoproteins (IDL) to LDL and in turn LDL to small dense LDL (sdLDL) [9, 10,19,20].

So, hypothyroidism is one of the most common causes of secondary dyslipidemia. Therefore, before starting hypolipidemic therapy, the evaluation of thyroid function is needed in PCOS women.

Objectives Of The Study

The aim of this study is

- To study the correlation between Serum T3, T4 & TSH with lipid profile test ie Serum cholesterol, Triglycerides, HDL, LDL & VLDL level in PCOS women
- To study correlation of Thyroid hormone dysfunction and cardiovascular diseases in PCOS so that early screening and prevention of heart diseases in PCOS with thyroid dysfunction can be advised

Material & Method Of Collection Of Data & Selection Of Subjects:

50 patients attending the outpatient department of OBGY ward, IGGMC, and Nagpur were selected for study. 5 ml of Fasting blood sample was drawn from patients who were attending biochemistry OPD for thyroid hormonal tests ie estimation of T3, T4 & TSH. History was taken as per designed proforma and consent form was obtained. Their Blood samples was analysed for thyroid profile as well as Lipid Profile test ie Serum cholesterol, Triglycerides, HDL, LDL & VLDL in clinical Biochemistry Laboratory.

Thyroid profile would be done in all patients who fulfill the inclusion criteria. Informed consent was obtained from all patients. Detailed clinical history and clinical examination were undertaken with

preference to thyroid and cardiovascular diseases in PCOS women.

Thyroid tests was run on Elisa reader & washer while Serum cholesterol, Triglycerides, HDL, LDL & VLDL tests was done on Autoanalyser EM 460. Serum values of Thyroid stimulating hormone (TSH), thyroxine (T4) and tri-iodo thyronine (T3) were assayed by ELISA tests and compared with Serum cholesterol, Triglycerides, HDL, LDL & VLDL tests and values were compared & correlated. The data was analysed & Student's T-test was used for the calculation. P <0.05 was considered significant.

METHODS

For Serum Cholesterol Estimation - Kit based on CHOD-POD method (Autoanalyser)

(Normal range – 150 mg% to 200 mg %)

For Serum Triglycerides Estimation - Kit based on GPO method (Autoanalyser)

(Normal range – 130 to 200 mg %)

For Serum HDL-Chol Estimation – Kit based on Precipitation method (Autoanalyser)

(Normal range – 35 mg% to 60 mg %)

For Serum VLDL & LDL Estimation - Friedwald Formula

VLDL = TG/5 & LDL = CHO – (HDL+VLDL)

For T3, T4, TSH – Immunoassay Elisa kit method on Elisa Reader & Washer

Normal Range – T3 - 0.52 to 1.85 ng/ml, T4 - 5 to 15 ug/dl, TSH – 0.39 to 6.16 uIU/ml

Analysis was carried on Autoanalyser EM – 460 in clinical Biochemistry lab, IGGMC for Serum cholesterol, Triglycerides, HDL, LDL & VLDL while Elisa Reader for Thyroid tests. All estimations was done & their values were compared & correlated.

RESULT

Table 1: Mean Values Of Thyroid Test & Lipid Profile Test In PCOS Women

Total (50)	T3	T4	TSH	CHO	TG	HDL	VLDL	LDL
Mean	1.0427	6.611	4.634	182.01	163.97	48.57	32.794	100.646
S. D	0.340	2.605	4.706	56.79	55.793	9.748	11.158	56.356

Table 2: Positive & Negative Correlation Of Thyroid Test With Lipid Profile Test In PCOS WOMEN

Correlation	CHO	TG	HDL	VLDL	LDL
T3	-0.62075	-0.49634	0.511922	-0.49634	-0.61585
T4	-0.64154	-0.46037	0.49117	-0.46037	-0.63941
TSH	0.756942	0.547646	-0.51238	0.547646	0.743021

Table3: Student's T test (P <0.05 was considered significant.)

p values	CHO	TG	HDL	VLDL	LDL
T3	<0.0001	<0.0001	<0.00001	<0.00001	<0.00001
T4	<0.0001	<0.0001	<0.00001	<0.00001	<0.00001
TSH	<0.0001	<0.0001	<0.00001	<0.00001	<0.00001

DISCUSSION

Table No 1 shows mean values of Thyroid & lipid profile in PCOS women

Our results shows (Table 2) positive correlation between Serum TSH (Mean=4.63) and serum cholesterol level (Mean= 182) in PCOS women as well as serum Triglycerides, LDL and VLDL level (Mean 163,100,32) while TSH shows negative correlation with HDL (Mean=48) .This positive correlation was found to be statistically significant between TSH and Serum cholesterol, Triglycerides, LDL & VLDL.

Similarly vice versa with T3 and T4 in PCOS. Table 3 shows positive correlation was found to be statistically significant (P value less than 0.005) between TSH and Serum cholesterol, Triglycerides, LDL & VLDL. Similarly vice versa with T3 and T4. These findings similar to findings of Canaris GJ, Manowitz NR, Mayor G, et al showing hypothyroid PCOS women have increased levels of TC and LDL-C [3]. While in another study by Asvold BO, Vatten LJ, Nilsen TI, Bjoro T et al (4) found decrease in HDL-C levels in hypothyroidism in PCOS, again these findings also correlate with finding of our study. Ochs et al, found a possible association, while Singh et al found no significant association.[20] The data presented here clearly indicates how the biochemical markers of cardiovascular diseases may be

affected by alteration in the level of Thyroid hormones in the PCOS.[11,13,14,20] This study showed a significant increase in cholesterol, triglycerides, LDL in PCOS women with hypothyroidism.

CONCLUSION-

This study is done to simplify the importance of interactions between thyroid function and cardiovascular risk in PCOS women. Hypothyroidism (increased TSH) correlated with increase Serum cholesterol, Triglycerides, LDL & VLDL level which are sensitive markers of cardiac failure in PCOS, our study shows there is positive correlation which is statistically significant. So these results will help to increase clinical knowledge and enable clinicians to provide better management for their PCOS patients who have thyroid dysfunction in preventing further cardiovascular risk.

PCOS women who receive appropriate treatment for their thyroid disease have a decreased chance of developing cardiovascular diseases, so early screening of lipid profile test is advised in thyroid dysfunction in PCOS women.

REFERENCES

- Duntas LH. Thyroid disease and lipids. *Thyroid*. 2002;12:287–93.
- Friis T, Pedersen LR. Serum lipids in hyper- and hypothyroidism before and after treatment. *Clin Chim Acta*. 1987;162:155–63.
- Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. *Arch Intern Med*. 2000;160:526–34.
- Asuncion M, Calvo RM, San Millan JL, Sancho J, Avila S, Escobar-Morreale HF. A prospective study of the prevalence of the polycystic ovary syndrome in unselected Caucasian women from Spain. *J Clin Endocrinol Metab*. 2000;85:2434–2438.
- Legro RS, Kuneselman AR, Dunaf A. Prevalence and predictors of dyslipidemia in women with polycystic ovary syndrome. *Am J Med*. 2001;111:607–613.
- Sniderman AD, Scantlebury T, Cianflone K. Hypertriglyceridemic hyperapob: the unappreciated atherogenic dyslipoproteinemia in type 2 diabetes mellitus. *Ann Intern Med*. 2001;135:447–459.
- Barter PJ, Brewer HB, Jr, Chapman MJ, Hennekens CH, Rader DJ, Tall AR. Cholesteryl ester transfer protein: a novel target for raising HDL and inhibiting atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2003;23:160–167.
- Asvold BO, Vatten LJ, Nilsen TI, Bjoro T. The association between TSH within the reference range and serum lipid concentrations in a population-based study. *The HUNT Study*. *Eur J Endocrinol*. 2007;156:181–6.
- Bakker O, Hudig F, Meijssen S, Wiersinga WM. Effects of triiodothyronine and amiodarone on the promoter of the human LDL receptor gene. *Biochem Biophys Res Commun*. 1998;249:517–21.
- Shin DJ, Osborne TF. Thyroid hormone regulation and cholesterol metabolism are connected through Sterol Regulatory Element-Binding protein-2 (SREBP-2). *J Biol Chem*. 2003;278:34114–8.
- Faure P, Oziol L, Artur Y, Chomard P. Thyroid hormone (T3) and its acetic derivative (TA3) protect low-density lipoproteins from oxidation by different mechanisms. *Biochimie*. 2004;86:411–8.
- Pramfalk C, Pedrelli M, Parini P. Role of thyroid receptor B in lipid metabolism. *Biochim Biophys Acta* 2011; 1812:929-37.
- Liberopoulos EN, Elisaf MS. Dyslipidemia in Patients with thyroid disorders. *Hormones* 2002; 1:218-23.
- Hueston WJ, Pearson WS. Subclinical Hypothyroidism and the Risk of Hypercholesterolemia. *Ann Fam Med* 2004; 2:351-5.
- Xu C, Yang X, Liu W, Yuan H, Yu C, Gao L, Zhao J. Thyroid stimulating hormone, independent of thyroid hormone, can elevate the serum total cholesterol level in patients with coronary heart disease; a cross sectional design. *Nutr Metab* 2012; 9:9-44.
- Lagrost L. Regulation of cholesteryl ester transfer protein (CETP) activity: review of *in vitro* and *in vivo* studies. *Biochim Biophys Acta*. 1994;1215:209–36. [Pub Med] [Google Scholar]
- Kuusi T, Saarinen P, Nikkila EA. Evidence for the role of hepatic endothelial lipase in the metabolism of plasma high density lipoprotein2 in man. *Atherosclerosis*. 1980;36:589–93.
- Santamarina-Fojo S, Gonzalez-Navarro H, Freeman L, Wagner E, Nong Z. Hepatic lipase, lipoprotein metabolism, and atherogenesis. *Arterioscler Thromb Vasc Biol*. 2004;24:1750–4.
- Prieur X, Huby T, Coste H, Schapp FG, Chapman MJ, Rodriguez JC. Thyroid hormone regulates the hypotriglyceridemic gene APOA5. *J Biol Chem*. 2005; 280:27533–43.
- Knochenhauer ES, Key TJ, Kahsar-Miller M, Waggoner W, Boots LR, Azziz R. Prevalence of the polycystic ovary syndrome in unselected black and white women of the southeastern United States: a prospective study. *J Clin Endocrinol Metab*. 1998;83:3078–3082.