



A RETROSPECTIVE STUDY OF CLINICAL AND ETIOLOGICAL PROFILES OF PATIENTS PRESENTING WITH PANCYTOPENIA

Dr.kamlesh Ninama Assistant Professor, Department of Medicine, ZMCH, DAHOD, GUJRAT, 389151.

Dr.Brajendra Kumar* Professor And HOD, Department of Medicine, ZMCH, DAHOD, GUJRAT, 389151.
*Corresponding Author

ABSTRACT Early diagnosis and treatment are crucial in the management of pancytopenia. A retrospective cross-sectional study was undertaken at ZMCH,DAHOD. All eligible participants more than 13 years of age were included. A thorough history taking along with clinical examination and laboratory investigations were performed among the study population. Data was analyzed using SPSS version 13. The most common finding suggestive of etiology was megaloblastic anemia at 38.4%, which was more common among females. Aplastic anemia was the second most common finding at 24.6%. Other diagnoses included hypersplenism, myelodysplastic syndrome and acute leukemia. Fever and pallor were the most common presenting symptoms. Blood examinations demonstrated anisopoikilocytosis, hypersegmented neutrophils, erythroblasts, macrocytes and reticulocytes, which were found in differing proportions in the various etiological diagnoses. It is noteworthy that megaloblastic anemia, a reversible condition, is common in India compared to the higher occurrence of aplastic anemia and myelodysplastic syndrome in developed nations.

KEYWORDS : Pancytopenia, Anemia, Megaloblastic, Aplastic.

INTRODUCTION

Pancytopenia is characterized by a decrease in the three blood elements (red blood cells, white blood cells and platelets) below the normal. Pancytopenia in adults is characterized by a triad of findings – a) hemoglobin less than 13.5 g/dL in males and less than 11.5 g/dL in females, b) total leucocyte count less than 4000/mm³, and c) platelet count less than 150000/mm³. It is a relatively less explored, but important clinic-hematological entity and not a disease which may occur due to a number of pathogenetic mechanisms. It may occur due to damage or suppression of pluripotent stem cells or committed progenitor cells. Pluripotent cell damage is less common while most cytotoxic drugs used in malignancies exert their action on the committed progenitor cells. Pancytopenia may be associated with a hypocellular bone marrow, hypercellular marrow or bone marrow infiltration. Pancytopenia with a hypocellular bone marrow may be caused by aplastic anemia, drugs, viruses, radiation, toxins, autoimmune disease, malignancies, PNH etcetera. In cases of pancytopenia with hypercellular bone marrow, the causative factors may be hypersplenism, megaloblastic anemia, myelodysplastic syndrome, infections such as HIV and TB etcetera. Malignancies, granulomas and fibrosis are causes of pancytopenia with bone marrow infiltration. The etiology of pancytopenia differs from one geographical area to another. The common causes identified in developing countries are megaloblastic anemia, infection, drugs, hypersplenism and aplastic anemia. Megaloblastic anemia has been reported as the cause of pancytopenia at a rate between 38% and 72% in local studies in the country, suggesting that it should be promptly evaluated for. The severity of the condition and the underlying pathogenesis decide the management and prognosis of the patients. Usually patients with pancytopenia present symptoms attributable to anemia and thrombocytopenia. The most common clinical manifestations are fever, fatigue, dizziness, weight loss, anorexia, night sweats, pallor, bleeding, splenomegaly, hepatomegaly and lymphadenopathy. The present study attempts to add to the limited literature available on the clinical and etiological profile of patients presenting with pancytopenia.

MATERIAL AND METHODS

The present study was a cross-sectional study conducted among patients admitted in the general medical wards at the ZMCH,DAHOD. All patients aged more than 13 years, with hemoglobin less than 12 g/dL in women and less than 13 g/dL in men, WBC count less than 4000 cells/dL and platelet count less than 150000/dL were included in the study. Patients with a known hematological condition, those on cancer chemotherapy and who had received blood transfusion were excluded from the study. After acquiring informed consent from all eligible subjects, a detailed relevant history was obtained including the dietary history, treatment history, intake of alcohol and drugs and radiation exposure. This was followed by a clinical examination of the patients for pallor, fever, bleeding tendencies, jaundice, hepatomegaly, splenomegaly, sternal tenderness and lymphadenopathy. Basic investigations were performed for each subject. The panel consisted of Hemoglobin, total leucocyte count, platelet count, reticulocyte count and

liver function tests. Values such as packed cell volume (PCV), Mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) were calculated for every patient. Viral markers (HBsAg, Anti HCV, HIV) were done in all patients. Chest radiograph and abdominal ultrasonography were done for all patients. Peripheral smear examination and Bone marrow examination were done in all patients and wherever required, a trephine biopsy was also performed. Vit. B12 levels, serum folate levels, upper and lower gastrointestinal endoscopy were done in selected patients. Also thyroid profile is done in selected patients.

RESULTS

Among the study population, 35 (53.85%) participants were male remaining 30 (46.15%) participants were female. Among male participants, 6 (17.14%) participants were aged below 21 years, 10 (28.57%) participants were aged between 21 to 30 years, 9 (25.61%) participants were aged between 31 to 40 years, 5 (14.28%) participants were aged between 41 to 50 years and above 50 years age group for each. Among female participants, 2 (6.67%) participants were aged below 21 years, 5 (16.67) participants were aged between 21 to 30 years, 7 (23.33%) participants were aged between 31 to 40 years, 9 (30%) participants were aged between 41 to 50 years and 7 (23.33%) participants were aged above 50 years age. Among the male participants, majority of 31.42% participants had megaloblastic anemia, followed by Aplastic anemia, hypersplenism and myelodysplastic syndrome was 28.57%, 17.14% and 8.57% respectively. Among the female participants, majority of 46.67% participants had megaloblastic anemia, followed by Hypersplenism and Acute leukemia was 20% and 13.33% respectively. Among the people with fever, majority of 35% participants had aplastic anemia, followed by, megaloblastic anemia and acute leukemia was 35% and 15% respectively. Among the people with pallor, majority of 39.06% participants had megaloblastic anemia, followed by Aplastic anemia, hypersplenism and myelodysplastic syndrome 25%, 12.5% and 9.38% respectively. Among the people with Anisopoikilocytosis, majority of 79.31% participants had megaloblastic anemia and 13.79% participants had aplastic anemia. Among the people with Erythroblasts all of them 100% had megaloblastic anemia. Among the people with hyper segmented neutrophils, all of them 100% participants had megaloblastic anemia. Among the people with Immature WBC, all of them 100% participants had acute leukemia. Among the people with Lymphocytosis was 100% participants had megaloblastic anemia. Among the people with Reticulocytosis, majority of 60% participants had megaloblastic anemia and 40% participants had hypersplenism. Among the people with macrocytes, majority of 93.75% participants had megaloblastic anemia and 6.25% participants had Aplastic anemia.

DISCUSSION

A total of 65 subjects were included in the study. The proportion of males (53.85%) was slightly higher than the proportion of females (46.15%). This is similar to the study by Makheja, et al. were 58% of those with pancytopenia were males and 42% were females. The

highest incidence of pancytopenia was found among those 21 to 30 years old and 31 to 40 years old with regards to men. While among women, pancytopenia was most common among 41 to 50 years followed by 21 to 30 years and above 50 years of age. The most common diagnosis made as to the etiology of pancytopenia was megaloblastic anemia among both genders (31.42% among males and 46.67% among females). The combined prevalence was 38.4%. This is similar to the study by Doshi, et al. where the most common cause of pancytopenia was megaloblastic anemia found among 45% of the study population. Gayathri and Rao, et al. report a much higher prevalence of megaloblastic anemia (74.04%). Makheja, et al. report that 41.9% of pancytopenia cases were due to megaloblastic anemia. Similarly, Khodke, et al. Tilak and Jain, et al. also report megaloblastic anemia to be the most common cause of pancytopenia. This increased prevalence of megaloblastic anemia is more commonly reported among developing countries due to the increased incidence of malnutrition and vitamin deficiencies. Hence in countries such as India, megaloblastic anemia is one of the reversible etiologies of pancytopenia. In our study, the higher prevalence of megaloblastic anemia among females also is indicative of the fact that females are at an increased risk of micronutrient deficiencies. Aplastic anemia was the second most common etiology (28.57% among males and 20% among females). The combined prevalence was 24.6%. In the study by Doshi, et al. aplastic anemia was less common (3%). Other causes were hypersplenism, myelodysplastic syndrome and acute leukemia. When peripheral smear examination was performed among the patients, anisopoikilocytosis, hypersegmented neutrophils, erythroblasts, macrocytes and reticulocytes were the most common findings among those with megaloblastic anemia. This is similar to the study by Doshi, et al. where macrocytes and hypersegmented neutrophils were the common peripheral smear findings in pancytopenia. They also report that the presence of megaloblasts indicate the early stages of megaloblastosis. Those with aplastic anemia had anisopoikilocytosis, macrocytes and microcytic, hypochromic RBCs.

CONCLUSION

The pathological and etiological profile of pancytopenia is variable among different countries. Conditions such as aplastic anemia, myelofibrosis and myelodysplastic syndrome are more common among the developed nations. In contrast, among developing countries such as India, reversible causes such as megaloblastic anemia are more common, especially among females. A complete history taking, assessing the clinical features such as fever, pallor along with performing a complete hemogram including a peripheral smear is essential for the diagnosis of pancytopenia. The rare causes such as Myelodysplastic syndrome, connective tissue disorders, HIV infection and aplastic anemia also have to be kept in mind after excluding the most common causes.

REFERENCES

1. Gayathri BN, Rao KS. Pancytopenia: A Clinico Hematological Study. *J Lab Physicians*, 2011;3(1): 15-20.
2. Ishtiaq O, Baqai HZ, Anwer F, et al. Patterns of pancytopenia patients in a general medical ward and a proposed diagnostic approach. *J Ayub Med Coll Abbottabad*, 2004; 16(1): 8-13.
3. Imbert M, Scoazec JY, Mary JY, Jouzult H, Rochant H, Sultan C. Adult patients presenting with pancytopenia: a reappraisal of underlying pathology and diagnostic procedures in 213 cases. *Hematol Pathol.*, 1989; 3(4): 159-67.
4. Gayathri BN, Rao KS. Pancytopenia: A Clinico Hematological Study. *J Lab Physicians*, 2011;3(1): 15-20.
5. Khodke K, Marwah S, Buxi G, Yadav R, Chaturvedi N. Bone Marrow Examination in Cases of Pancytopenia. *Journal of Indian Academy of Clinical Medicine*, 2001;2(1): 5.
6. Tilak V, Jain R. Pancytopenia-a clinico-hematologic analysis of 77 cases. *Indian J Pathol Microbiol.*, 1999 Oct; 42(4): 399-404.