



## A STUDY ON ELECTROLYTE CHANGES IN NEONATES RECEIVING PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINAEMIA

<b>Dr. Nitish Kumar</b>	M.D. (Pediatrics), Senior Resident, Department of Pediatrics, All India Institute of Medical Sciences (AIIMS), Patna, Bihar.
<b>Dr. Aparna Kumari*</b>	M.D. (Skin & VD), Senior Resident, Department of Dermatology, Employee State Insurance Corporation Medical College and Hospital (ESICMCH), Bihta, Patna, Bihar. *Corresponding Author
<b>Dr. Kripa Nath Mishra</b>	Professor and Head, Department of Pediatrics, Darbhanga Medical College and Hospital (DMCH), Laheriasarai, Bihar.

**ABSTRACT** **BACKGROUND:** Hyperbilirubinemia is the most common ailment requiring medical attention in newborns. Neonatal hyperbilirubinemia nearly affects 60% of term & 80% of preterm neonates during the first week of life. Neonatal jaundice has much higher incidence in premature babies and often requires therapeutic intervention. Neonatal physiological jaundice could result either due to increased breakdown of foetal erythrocytes or low concentrations of hepatic glucuronyl transferase. Hypocalcaemia is one of the known adverse effects of phototherapy. 90% of the preterm and 75% of the full-term neonates develop hypocalcaemia post phototherapy. We intended to study the effects of photo therapy on serum electrolytes (Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>++</sup>).

**METHODS:** It is a prospective interventional study. Neonates who were born or admitted to Department of Pediatrics, Darbhanga Medical College and Hospital, Laheriasarai, Bihar, from March 1<sup>st</sup>, 2016 to February 28<sup>th</sup>, 2018 receiving phototherapy for unconjugated hyperbilirubinemia after 24 hrs. of life without any co-morbidities were studied.

**RESULTS:** Serum calcium and serum sodium levels decrease with increasing duration of phototherapy. The comparative correlation between the preterm and term neonates considering the change in calcium level before and after phototherapy shows more change in values in term neonates.

**CONCLUSIONS:** Electrolyte changes are seen more commonly in pre-term neonates. The incidence of hypocalcemia was 22%, it was 51.6% at >48 hrs. duration of phototherapy (PT) and 52.9% in preterm neonates. The incidence of hyponatremia was 10%, it was 19.3% at >48 hrs. duration of PT and 29.4% in preterm neonates. The mean serum calcium levels before and after PT were 1.08±0.135 mmol/L and 1.008±0.161 mmol/L respectively.

**KEYWORDS :** Electrolytes, Phototherapy, Neonatal Hyperbilirubinemia

### INTRODUCTION

Hyperbilirubinemia is the most common ailment requiring medical attention in newborns. Neonatal hyperbilirubinemia nearly affects 60% of term and 80% of preterm neonates during first week of life.<sup>1</sup> This is a normal physiologic phenomenon in most infants. However excessively elevated serum bilirubin can be detrimental as unconjugated bilirubin is neurotoxic<sup>2</sup> and can cause residual neurologic sequelae or even death in some and hence is a cause for concern in pediatricians and family.<sup>3</sup>

Neonatal jaundice has much higher incidence in premature babies and often require therapeutic intervention.<sup>4</sup> Neonatal physiologic jaundice could result either due to increased breakdown of foetal erythrocytes or low concentrations of hepatic glucuronyl transferase.<sup>5,6</sup> When additional factors like immune or non-immune hemolytic anemia, polycythemia, several metabolic and endocrine disorders accompany the basic mechanisms pathologic neonatal jaundice ensues.

Incidence of neonatal jaundice is higher in eastern world. Various genetic influences are known to increase the incidence, especially mutations/polymorphisms in the genes that code for enzymes and proteins involved in bilirubin metabolism.<sup>7</sup> Recent data suggests a correlation between higher levels of epidermal growth factors both in breast milk and infants with neonatal jaundice.<sup>8</sup> Incidence is also higher in premature infants, infants with low birth weight, infants of mothers with diabetes and congenital infections.

Main treatment modalities in neonatal jaundice are Phototherapy, exchange transfusion and intravenous immune globulin. Phototherapy is the primary treatment in neonates with unconjugated hyperbilirubinemia.<sup>9</sup> Phototherapy isomerizes the bilirubin into water-soluble forms by photo oxidation, configurational isomerization, and structural isomerization. These photo isomers of bilirubin are excreted in bile and urine thereby reducing the serum bilirubin levels.

Adverse effects of phototherapy include feed intolerance, loose stools, bronze baby syndrome, hypocalcaemia etc.<sup>10</sup>

Although many studies were carried out, to depict the other side effects those relating to dyselectrolytemia have been few, especially those

relating to serum sodium and potassium. Hypocalcaemia is one of the known adverse effects. 90% of the preterm and 75% of the full term neonates develop hypocalcaemia post phototherapy.<sup>11</sup> We intended to study the effects of photo therapy on serum electrolytes (Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>++</sup>).

### AIMS AND OBJECTIVES

1. To study the electrolyte changes in neonates receiving phototherapy for neonatal jaundice at Department of Pediatrics, DMCH, Laheriasarai, Bihar.
2. To assess the effect of Phototherapy on Serum Sodium, Serum Potassium and Serum Calcium levels.

### MATERIAL AND METHODS

It is a prospective interventional study. Neonates who were born or admitted to Department of Pediatrics, Darbhanga Medical College and Hospital, Laheriasarai, Bihar from March 1<sup>st</sup> 2016 to February 28<sup>th</sup> 2018, receiving Light Emitting Diode(LED) phototherapy for unconjugated hyperbilirubinemia after 24 hrs. of life without any co-morbidities were studied. All the investigation was done in the laboratory of DMCH, Laheriasarai, Bihar.

### Inclusion Criteria

Neonates receiving phototherapy for unconjugated hyperbilirubinemia after 24 hrs. of life without any co-morbidities were included in the study.

### Exclusion Criteria

1. Neonates with evidence of Hemolysis.
2. Onset of jaundice < 24 hrs. & received exchange transfusion.
3. Neonates with Conjugated Hyperbilirubinemia.
4. Neonates who receive I.V. fluid and intensive care.
5. Neonates with formula feeding
6. Neonates with co-morbidities like birth asphyxia, septicaemia, acute renal failure and others.
7. Abnormal electrolyte status detected Pre phototherapy.

### Study Protocol

Patients were randomly selected following the inclusion and exclusion criteria. The Phototherapy was administered as per standard American

Association of Pediatrics (AAP) guidelines. Electrolytes were checked at 0 hour (First sample) and at 48 hours of phototherapy or at discontinuation of phototherapy (Second sample) whichever is earlier. The first sample was considered as control. Comparative study was made between these two samples groups to determine the changes in electrolytes. The blood tests were run on the collected samples and data tabulated. Serum electrolytes (Na<sup>+</sup>, K<sup>+</sup> and Ca<sup>2+</sup>) were estimated using auto analyzer which employs 'ion selective electrode' for estimation. By Diazo method Serum bilirubin was estimated. A total of 100 neonates (including 56 males and 44 females) including both term and preterm were studied. The data obtained were tabulated and analyzed by proper statistical method.

## RESULTS

**Table 1 : Incidence of Neonatal Hyperbilirubinemia with respect to Birth Weight**

Weight (Kg)	Gender		Total (%)
	Male	Female	
LBW (<2.5 kg)	11(19.6%)	12(27.2%)	23(23%)
Normal	45(80.4%)	32(72.8%)	77(77%)
Total	56	44	100

**Table 2 : Incidence of Neonatal Hyperbilirubinemia with respect to Gestational age**

Gestational age (weeks)	Gender		Total (%)
	Male	Female	
<37	8(14.3%)	9(20.5%)	17(17%)
37-40	46(82.1%)	34(77.3%)	80(80%)
>40	2(3.6%)	1(2.2%)	3(3%)
Total	56(100%)	44(100%)	100(100%)

Mean gestational age is 37.22 ± 2.01 wks. contingency coefficient 0.0870, p=0.6828

**Table 3 : Comparison between pre-phototherapy and post-phototherapy serum calcium with respect to duration of phototherapy**

Calcium	Duration of PT			Total	p-value
	<24 hrs.	24-48 hrs.	>48 hrs.		
<1mmol/L	0(0%)	5(7.7%)	17(51.6%)	22(22%)	<0.0001
1-1.23mmol/L	4(100%)	60(92.3%)	15(48.4%)	78(78%)	
>1.23mmol/L	0	0	0	0	

**Table 4 : Comparison between pre-phototherapy and post-phototherapy serum calcium with respect to Gestational Age**

Serum Calcium (Ionic)	Gestational Age			Total	p-value
	<37 (17)	37-40 (80)	>40 (3)		
<1mmol/L	9(52.9%)	12(15%)	1(33.3%)	22(22%)	<0.0113
1-1.23mmol/L	8(47.1%)	60(85%)	2(66.7%)	78(78%)	
>1.23mmol/L	0	0	0	0	

**Table 5 : Comparison between pre-phototherapy and post-phototherapy serum sodium with respect to duration of phototherapy**

Serum Sodium	Duration of PT			Total (100)	p-value
	<24 hrs. (4)	24-48 hrs. (65)	>48 hrs. (31)		
<135mmol/L	0	4(6.2%)	6(19.3%)	10(10%)	0.0167
135-145mmol/L	4(100%)	61(93.8%)	22(70.9%)	87(87%)	
>145mmol/L	0	0	3(9.6%)	3(3%)	

**Table 6 : Comparison between pre-phototherapy and post-phototherapy serum sodium with respect to Gestational Age**

Serum Sodium	Gestational Age			Total	p-value
	<37 wks (17)	37-40 wks (80)	>40 wks (3)		
<135mmol/L	5(29.4%)	4(5%)	1(33.3%)	10(10%)	0.0204
135-145mmol/L	12(70.6%)	73(91.2%)	2(66.7%)	87(87%)	
>145mmol/L	0	3(3.8%)	0	3(3%)	

**Table 7 : Comparison between pre-phototherapy and post-phototherapy serum potassium with respect to duration of phototherapy**

Serum Potassium	Duration of PT			Total	p-value
	<24 hrs.	24-48 hrs.	>48 hrs.		
<3.5mmol/L	0(0%)	0(0%)	1(3.2%)	1(1%)	0.3375
3.5-5.5mmol/L	4(100%)	65(65%)	29(93.6%)	98(97%)	
>5.5mmol/L	0(0%)	0(0%)	1(3.2%)	21(2%)	

**Table 8 : Comparison between pre-phototherapy and post-phototherapy serum Potassium with respect to Gestational Age**

Serum Potassium	Gestational Age			Total	p-value
	<37 wks (17)	37-40 wks (80)	>40 wks (3)		
<3.5mmol/L	1(5.9%)	0(0%)	0(0%)	1(1%)	0.248
3.5-5.5mmol/L	16(94.1%)	78(97.5%)	3(100%)	97(97%)	
>5.5mmol/L	0(0%)	2(2.5%)	0(0%)	2(2%)	

**Table 9 : Conclusive comparative evaluation of pre and post-phototherapy serum electrolytes**

	Pre.	Post.	Difference	t-value	p-value
Ionic Calcium	1.08±0.14	1.007±0.16	0.07	4.536	<0.0001
Sodium	139.1±2.7	137.6±2.8	1.5	5.56	<0.0001
Potassium	4.4±0.5	4.5±0.5	-0.1	2.00	0.05

## Principles of Biochemical Tests

SERUM Na<sup>+</sup>, K<sup>+</sup> and Ca<sup>2+</sup> (86) – Ion Selective Electrode: An ion-selective electrode (ISE), also known as a specific ion electrode (SIE), is a transducer (or sensor) that converts the activity of a specific ion dissolved in a solution into an electrical potential. The voltage is theoretically dependent on the logarithm of the ionic activity, according to the Nernst equation. Ion-selective electrodes are used in analytical chemistry and biochemical/biophysical research, where measurements of ionic concentration in an aqueous solution are required. Serum Bilirubin – Diazo.

Ehrlich's diazo reagent reacts with the direct bilirubin in the serum to form a pink to reddish-purple colored compound (Azobilirubin) read at one minute.

## Data Analysis

All the statistical analysis was done using Windows Excel, IBM SPSS 24 software, Med Calc 9.0.1 and Systat 12.0. To assess the association of different parameters Pearson's Chi-Square test was used. Differences were considered statistically significant if p<0.05, very significant if p<0.01 and very highly significant if p<0.001.

In our study incidence of low birth weight babies was 23%, mean birth weight was 2.61 ± 0.38 Kgs. In our study we found out that among low birth weight (LBW) neonates, incidence of females was more than males. However, it was not statistically significant.

In our study group, the incidence of preterm babies was 17% compared to 80% in 37-40 wks. gestational age and 3% in >40 wks gestational age group. The mean gestational age in the study group was 37.22±2.01 weeks.

The Incidence of hypocalcaemia was 51.6% when duration of phototherapy was >48 hrs. as compared to 7.7% when the duration was between 24-48 hrs. Thus, it infers that babies were at higher risk of hypocalcaemia if kept under phototherapy for more than 48 hrs. chi-square test revealed the p value obtained was <0.0001. Thus, it is considered to be statistically significant.

The Incidence of hypocalcaemia following phototherapy was more in preterm neonates (52.9%) than in term neonates (15%). Thus, it infers that preterm babies were at more risk of hypocalcaemia following phototherapy than term babies. By chi-square test, using test for paired sample means, the p value obtained was <0.05. Thus, it is considered to be statistically significant.

The Incidence of hyponatremia was 19.3% when duration of phototherapy was >48 hrs. as compared to 6.2% when the duration was between 24-48 hrs. Thus, it infers that babies were at higher risk of hyponatremia if kept under phototherapy for more than 48 hrs. Chi-square test revealed the p value obtained was 0.0167. Thus, it is considered to be statistically significant.

The Incidence of hyponatremia following phototherapy was more in preterm neonates (29.4%) than in term neonates (5%). Thus, it infers that preterm babies were at more risk of hyponatremia following phototherapy than term babies. By chi-square test the p value obtained was <0.05. Thus, it is considered to be statistically significant.

The Incidence of hypokalaemia was 3.2% when duration of phototherapy was >48 hrs. Thus, it infers that babies were at higher risk of hypokalaemia if kept under phototherapy for more than 48 hrs.

However, Chi-square test revealed that the p value was 0.3375. Thus, it is considered to be statistically insignificant.

The Incidence of hypokalaemia following phototherapy was more in preterm neonates (5.9%) than in term neonates. Thus, it infers that preterm babies were at more risk of hypokalaemia following phototherapy than term babies. By chi-square test the p value obtained was 0.226. Thus, it is considered to be statistically insignificant.

Overall, there was a significant decline in serum calcium and serum sodium values post phototherapy.

## DISCUSSION

Decreasing serum electrolytes is one of the known complications of phototherapy. While hypocalcaemia has been well documented to be associated with phototherapy effects of phototherapy on other major serum electrolytes has been meagrely studied. Our study studied the effects of phototherapy on major serum electrolytes in neonates being treated for hyperbilirubinemia and noted statistically significant results. The present study included a total of 100 neonates 35 weeks. Term gestation was considered to be more than and equal to 37 weeks' gestation and those <37 weeks' gestation age were considered to be pre term. Karamifar et al<sup>12</sup> 2002 selected pre terms more than 31 weeks whereas Eghbalian et al<sup>13</sup> (2002) included only term neonates. Arora et al<sup>14</sup> (2014) included 46 preterm and 54 term neonates. All the term and preterm neonates in the present study were appropriate for gestational age. Same was the case in the studies done by Romagnoli et al<sup>15</sup> (1979), Sethi et al<sup>16</sup> (1993), and Arora et al<sup>14</sup> (2014).

### Comparison of Mean Birth Weight with Other Studies

Mean birth weight of our study group members is 1.94 Kgs  $\pm$  140 grams in pre-term vs 2.73 Kg  $\pm$  275 grams in term neonates. These results match Jain et al (18) study in which mean birth weight in preterm vs term were 2150  $\pm$  150 grams vs 2800  $\pm$  220 grams and Karamifar et al<sup>12</sup> 2077  $\pm$  316 grams Vs 2889  $\pm$  474 grams respectively.

### Comparison of Duration of Phototherapy

In our study the mean duration of phototherapy was 40  $\pm$  9.7 hrs. Mean duration of phototherapy in a similar study conducted by Eghbalian et al<sup>13</sup> was 3 days. This could be because of larger sample size and more sample of preterm neonates in their study.

### Comparison of Hypocalcaemia Incidence with Other Studies

In our study we found out that 9 preterm and 13 term/post term cases were found to have hypocalcaemia. They amount to 52.9% and 15.6% of incidence respectively. The relation was found to be statistically significant with a p value of 0.011. the significant fall in serum calcium level in preterm and term babies after phototherapy was in correlation with study by Karamifer et al.<sup>12</sup> The trends and significance levels in our study are in correlation with similar studies by Yadav RK et al,<sup>17</sup> Arora et al<sup>14</sup> and Reddy et al.<sup>18</sup>

Sethi et al<sup>16</sup> study has shown that 90% of Preterm neonates and 75% of full-term neonates treated with phototherapy developed hypocalcaemia.

Jain et al<sup>19</sup> have found that 55% of preterm neonates and 30% of full-term neonates developed hypocalcaemia after phototherapy. This study is in correlation with our study. Jain et al have found that 55% of preterm neonates and 30% of full-term neonates developed hypocalcaemia after phototherapy. This study is in correlation with our study.

### Comparison of Mean Serum Calcium Levels Before and After Phototherapy in Preterm Neonates

In our study the mean serum ionized calcium level before and after phototherapy in preterm neonates was 0.89  $\pm$  0.2 mmol/L and 0.92  $\pm$  0.2 mmol/dl respectively and it was statistically significant. This was similar to study done by Karamifar et al.<sup>12</sup>

Comparison of mean serum calcium levels before and after phototherapy in term neonates.

In our study the mean serum calcium level before and after phototherapy in preterm neonates was 1.12  $\pm$  0.06 mmol/L and 1.02  $\pm$  0.14 mmol/L respectively and it was statistically significant. This was similar to study done by Karamifar et al<sup>12</sup> and Reddy AK et al.<sup>18</sup>

## Hyponatremia

There are very few studies regarding phototherapy induced other electrolyte changes. The differential effect of other electrolytes with phototherapy has not been studied by other workers except that for Curtis MD et al<sup>20</sup> (1981) study which stated that absorption of water, sodium chloride, and potassium was significantly impaired in the patients receiving phototherapy. Incidence of hyponatremia in our study group was 10% and found to be higher in <37 wks. group and LBW babies than in >37 wks. group and normal weight babies. Mean serum sodium levels were significantly decreased after phototherapy. As the P value <0.05, this difference is considered to be statistically significant.

## Hypokalaemia

There were no significant potassium changes in our study. Tan KL et al<sup>21</sup> (1981) study in healthy full-term neonates demonstrated a transient raise in potassium levels after phototherapy which was in contrast to our study. It is evident that in the present study phototherapy induced hypocalcaemia and hyponatremia was more in preterm and LBW babies but the actual relationship in these babies with phototherapy has to be evaluated with larger sample studies for estimation of the incidence as the metabolic side effects are more common in preterm and LBW babies.

## CONCLUSIONS

Serum calcium and serum sodium levels decrease with increasing duration of phototherapy. These electrolyte changes are seen more frequently in pre-term neonates. The comparative correlation between the preterm and term neonates considering the change in calcium level before and after phototherapy shows more change in values in term neonates. A total of 100 neonates (including 56 males and 44 females) including both term and preterm were studied. The mean birth weight was 2.61 $\pm$ 0.38 Kgs. mean gestational age was 37.22 $\pm$ 2.01 weeks. The incidence of hypocalcaemia was 22%, it was 51.6% at >48 hrs. duration of PT and 52.9% in preterm neonates. The incidence of hyponatremia was 10%, it was 19.3% at >48 hrs. duration of PT and 29.4% in preterm neonates. The mean serum calcium levels before and after PT were 1.08  $\pm$  0.135 mmol/L and 1.008  $\pm$  0.161 mmol/L respectively. The mean serum sodium levels before and after PT were 139.1  $\pm$  2.7 and 137.6  $\pm$  2.8.

## REFERENCES

1. Watchko JF. Hyperbilirubinemia in African American neonates: clinical issues and current challenges. *Semin Foetal Neonatal Med* 2010;15(3):176-82.
2. Shapiro SM. Bilirubin toxicity in the developing nervous system. *Pediatr Neurol* 2003;29(5):410-21.
3. Maisels MJ, Kring E. Length of stay, jaundice and hospital readmission. *Pediatrics* 1998;101(6):995-8.
4. Kaini NR, Chaudhary D, Adhikary V, et al. Overview of cases and prevalence of jaundice in neonatal intensive care unit. *Nepal Med Coll J* 2006;8(2):133-5.
5. Huang MJ, Kua KE, Teng HC, et al. Risk factors for severe hyperbilirubinemia in neonates. *Pediatr Res* 2004;56(5):682-9.
6. Christensen RD, Yaish HM, Lemons RS. Neonatal hemolytic jaundice: morphologic features of erythrocytes that will help you diagnose the underlying condition. *Neonatology* 2014;105(4):243-9.
7. Hua L, Shi D, Bishop PR, et al. The role of UGT1A1\*28 mutation in jaundiced infants with hypertrophic pyloric stenosis. *Pediatr Res* 2005;58(5):881-4.
8. Kumral A, Ozkan H, Duman N, et al. Breast milk jaundice correlates with high levels of epidermal growth factor. *Pediatr Res* 2009;66(2):218-21.
9. Woodgate P, Jardine LA. Neonatal jaundice. *BMJ Clin Evid* 2011;2011. pii: 0319.
10. Maisels MJ. Neonatal Jaundice. *Pediatr Rev* 2006;27(12):443-54.
11. Narayan S, Aggarwal R, Upadhyay A, et al. Survival and morbidity in extremely low birth weight (ELBW) infants. *Indian Pediatr* 2003;40(2):130-5.
12. Karamifar H, Pishva N, Amirhakimi GH. Prevalence of phototherapy induced hypocalcaemia. *Iran J Med Sci* 2002;27(4):166-8.
13. Eghbalian F, Monsef A. Phototherapy-induced hypocalcemia in icteric newborn. *Iran J Med Sci* 2002;27(4):169-71.
14. Arora S, Narang GS, Singh G. Serum calcium levels in preterm and term neonates on phototherapy. *J Nepal Paediatr Soc* 2014;34(1):24-8.
15. Romagnoli C, Polidori G, Cataid L, et al. Phototherapy induced hypocalcemia. *J Pediatr* 1979;94(5):815-16.
16. Sethi H, Salli A, Dutta AK. Phototherapy induced hypocalcemia. *Indian Pediatr* 1993;30(12):1403-6.
17. Yadav RK, Sethi RS, Sethi AS, et al. The evaluation of effect of phototherapy on serum calcium level. *People's J Sci Res* 2012;5(2):1-4.
18. Reddy AT, Bai KV, Shankar SU. Electrolyte changes following phototherapy in neonatal hyperbilirubinemia. *Int J Sci Res* 2015;4(7):752-8.
19. Jain BK, Singh H, Sigh D, et al. Phototherapy induced hypocalcemia. *Indian Pediatr* 1998;35(6):566-7.
20. Curtis-Cohen M, Stahl GE, Costarino AT, et al. Randomized trial of prophylactic phototherapy in the infant with very low birth weight. *J Pediatr* 1985;107(1):121-4.
21. Tan KL. Efficacy of bidirectional fiber-optic phototherapy for neonatal hyperbilirubinemia. *Pediatrics* 1997;99(5):E13.