



STUDY OF ER PR RECEPTORS IN BREAST CARCINOMA

Dr. Manish Shah

M.B.B.S, M.S. (Gen. Surg.), Senior Resident, Department of General Surgery, Indira Gandhi Institute of Medical Sciences (IGIMS), Patna, Bihar.

Dr. [Prof.] Manish Mandal*

Professor and Head of Department, Department of GI Surgery, Indira Gandhi Institute of Medical Sciences (IGIMS), Patna, Bihar. *Corresponding Author

ABSTRACT

Background: Estrogen and Progesterone receptors are found in breast cancer cells that depend on estrogen and related hormones to grow. These are steroidal nuclear type of receptors. The most common method currently used to test a tumor for estrogen and progesterone receptors is immunohistochemistry or IHC. Patients with positive hormone receptor cancer have better survival. They are candidates of anti-hormonal therapy.

Methods: This study was carried out in the Department of Surgery of a IGIMS, Patna, Bihar from May 2019 to February 2020. 30 randomly selected case records of breast cancer patients, who presented in this period, and who had undergone surgery, were taken and studied in terms of history, examination, investigations, treatment given, histopathology report and development of recurrence, metastasis and survival.

Results: In this study out of 30 patients, 25 patients received adjuvant chemotherapy. 2 patients developed recurrence at local site. 1 patient was ER/PR both negative and 1 patient was ER +/PR-. 3 patients developed metastasis. Among these, 2 patients were ER/PR both negative and 1 patient was having ER/PR both positive.

Conclusions: Hormone receptors play a significant role in breast carcinoma. Breast cancer is more common in postmenopausal group as compared to premenopausal group. Mean age at diagnosis of breast cancer in western countries is 61 years and in India it is 50 years.

KEYWORDS : Carcinoma breast, Estrogen receptor, Progesterone receptor**INTRODUCTION**

Breast cancer is the most common female cancer in the world with an estimated 1.67 million new cancer cases diagnosed in 2012. With an annual incidence of approximately 1,44,000 new cases of breast cancers in India, it has now become the most common female cancer in urban India. In India, the incidence of breast cancer is significantly lower than western countries. Breast cancer in India varies from as low as 5 per 100,000 female populations per year in rural areas to 30 per 100,000 female populations per year in urban areas.¹ A cancer is called estrogen-receptor-positive (or ER+) if it has receptors for estrogen. This suggests that the cancer cells, like normal breast cells, may receive signals from estrogen that could promote their growth. The cancer is progesterone-receptor-positive (PR+) if it has progesterone receptors. Again, this means that the cancer cells may receive signals from progesterone that could promote their growth. Roughly two out of every three breast cancers test positive for hormone receptors.²

MATERIAL AND METHODS

This study was carried out in the Department of Surgery of Indira Gandhi Institute of Medical Sciences, Patna, Bihar from May 2019 to February 2020. 30 randomly selected case records of breast cancer patients, who presented in the period, and who had undergone surgery, were taken and studied in terms of history, examination, investigations, treatment given, histopathology report and development of recurrence, metastasis and survival. These 30 patients were divided according to ER/PR status in 4 groups that is ER+PR+ / ER+PR- / ER-PR+ / ER-PR-.

These 4 groups were compared in terms of breast cancer epidemiology, clinical features, stage, histopathology report, treatment given, and development of recurrence, metastasis and survival duration.

INCLUSION CRITERIA

All females having carcinoma breast who underwent surgery at least 1 year before, with known hormonal receptor status and who had taken chemotherapy/radiotherapy / hormonal therapy or combined therapy.

EXCLUSION CRITERIA

- Females age less than 20 years.
- Females with breast cancer but unknown hormonal receptor status.
- Females with breast cancer with metastasis.

RESULTS

In this study on ER PR receptor in breast carcinoma, 30 cases records of breast cancer patients were taken. In this study 6 patients were < 40-

year age, 14 patient in between 41 to 50 year and 10 patient having more than 51 years, so that the maximum number of patients of breast cancer were in age group of 41-49 years and next, more than 49 years in this study.

Table 1 : Stage of breast cancer and number of patients

Stage of carcinoma	Number of patient	Percentage
Early invasive	24	80%
Loco-regionally advanced	6	20%
Metastasis	0	0
Total	30	100%

11 patients were pre-menopausal, and 19 patients were post-menopausal. In this study breast carcinoma was more in Postmenopausal female as compared to Premenopausal females.

Out of 30 patients 25 patients having breast lump size 2-5 cm, 1 patient < 2 cm in size and 4 patients had > 5 cm size. Among the studied patient of breast carcinoma, majority of the patient were having breast lump of size in between 2-5cm.

In this study, 25 patients were having breast lump located in upper outer quadrant of the breast followed by 4 patients having lower inner quadrant. Among the patient presenting with breast lump, majority of the patients were diagnosed to have early invasive breast carcinoma, followed by loco regionally advanced breast carcinoma in this study (Table 1).

Table 2 : Type of breast carcinoma and number of patients

Type of carcinoma	Number of patient	Percentage
Infiltrating ductal carcinoma	30	100%
Other type	0	0
Total	30	100%

Table 2 shows all the patients of breast cancer in this study were having infiltrating ductal carcinoma.

Table 3 : Histological grade of breast cancer and number of patients

Histological grade	Number of patient	Percentage
1	2	7%
2	26	86%
3	2	7%
Total	30	100%

This Table 3 reveals that among the studied group of breast carcinoma, majority of patients were having histological grade 2 carcinoma.

Table 4 : Hormonal status and number of patients of breast cancer

Hormonal status	Number of patient	Percentage
ER/PR (+/+)	15	50%
ER/PR (+/-)	1	3%
ER/PR (-/+)	1	3%
ER/PR (-/-)	13	44%
Total	30	100%

In this study most patients were having either hormone receptor positive or both negative (Table 4).

Table 5 : Herceptin-2neu and number of patients of breast cancer

Herceptin-2neu	Number of patient	Percentage
Her-2neu (+)	14	53%
Her-2neu (-)	16	47%
Total	30	100%

Table 5 shows Herceptin-2neu receptor positivity was slightly more in this study.

Table 6 : Type of treatment given and number of patients of breast cancer

Treatment	Number of patient
Chemotherapy	29
Radiotherapy	30
Hormonal therapy	16
Neoadjuvant	5

In this study almost all patients received chemo and radiotherapy after surgery and some patients received hormonal therapy according to ER/PR status and few patients also received neoadjuvant therapy for down staging (Table 6).

Table 7 : Metastasis and number of patients of breast cancer

Metastasis	Number of patient	Percentage
Present	3	10%
Absent	27	90%
Total	30	100%

Table 7 shows most of the patients did not develop metastasis after treatment in this study.

Table 8 : Survival duration and number of patients of breast cancer

Survival duration	Number of patient	Percentage
Survival >12 months	27	90%
Survival <12 months	3	10%
Total	30	100%

Table 8 shows that most patients were having survival more than 12 months in this study. At the time of completion of study all remaining 27 patients were surviving and the duration ranged from 1 year to 5 years depending on the date of enrolment of study.

DISCUSSION

Breast cancer is one of the most common malignancies among women in most developed and developing countries of the world with nearly a million new cases reported each year. It accounts for nearly 21% of all cancers among women worldwide. As in the rest of the world, India too has been witnessing a surge in the incidence of female breast cancer, and it is now the most common cancer of women in many parts of India³.

In this study all the data of 30 patients of breast cancer has been tabulated and compared with similar type of study done in other institutions. Various studies which have been compared with this study are as follows:

The mean age at presentation of female breast carcinoma in the present study is 46.8 years which is a decade earlier compared to patients from West. The average age at diagnosis of breast carcinoma in the US is 61 years.^{4,5}

Most of patients in the study population are post-menopausal age group and they are also more, hormone receptor positive. Most patients in this study presented with painless breast lump in upper outer quadrant followed by painful lump followed by nipple discharge. In various studies conducted on carcinoma breast, cases were divided according to size of the lesion as cases with lesion size less than 2 cm,

cases with lesion size between 2 to 5 cm and those with lesion size more than 5 cm. In the present study most, patients were having lump size between 2 to 5cm. Study shows upper outer quadrant was most common quadrant involved in breast cancer. According to Bailey and Loves, Short practice of surgery, Upper outer quadrant is involved in 60%, upper inner 12%, lower outer 10%, lower inner 6% and central 12%.⁶

Most patients presented in this study were stage II cancer followed by stage III. The reason for late presentation may possibly be due to low socioeconomic status and lack of awareness regarding the disease. There was no significant difference between patients having any stage of cancer according to hormone receptor status. Most of the patients in various studies were having stage 2 and 3 breast cancer.

In this study all patients were having invasive ductal carcinoma-not otherwise specified (NOS) type. Various studies like Alsaeed, Daniele, Dhafir Al-Azawi, Gion also showed that majority of cases of carcinoma breast were Invasive ductal carcinoma-NOS type.

These various studies show that maximum number of patients were having grade 2 and they were also hormone receptor positive.

In this study receptor status was available in 30 cases of which 56.66% were ER and PR receptor positive and 43.33% were ER and PR negative. These studies show most breast cancers were ER receptor positive.

In present study PR positivity was more than all the studies for comparison. In present study Herceptin-2 neu positivity was more than almost all the studies for comparison.

Chemotherapy

In this study out of 30 patients, 5 patients received neoadjuvant chemotherapy, 25 patients received adjuvant chemotherapy. Most patients received either CAF or AC followed by TAXANES (paclitaxel/docitaxel). Chemotherapy was not related with hormone receptor status.

Metastasis

In this study of 30 patients, 3 patients developed metastasis. Among these, 2 patients were ER/PR both negative and 1 patient was having ER/PR both positive. There is no relation among hormone receptor status and development of metastasis.

In present study PR positivity was more than all the studies for comparison. In present study Herceptin-2 neu positivity was more than almost all the studies for comparison.

Hormone receptor and survival

Survival was less in hormonal receptor negative patients as compared to hormone receptor positive patient in present study.

Similar was seen in study of 'Hormone receptor status, tumor characteristics, and prognosis: a prospective cohort of breast cancer patients' done by Department of Epidemiology, University of Washington, USA and Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, USA, in which women with ER+/PR-, ER-/PR+, or ER-/PR- tumors experienced higher risks of mortality as compared to women with ER+/PR+ tumors, which were largely independent of the various demographic and clinical tumor characteristics.²⁰

In one more study by Williams MR, et al, the contribution of ER status to survival has been examined in those patients responding to therapy and compared to those progressing despite treatment. It was seen that in the latter group no significant difference existed between those patients with ER-positive and those with ER-negative tumours.

However, in patients assessed as responding to therapy a significant difference did occur between these two groups. Patients assessed as responding to therapy with ER-positive tumours had a significant survival advantage over the small number of responding patients with ER-negative Primary tumours.²¹

CONCLUSION

Most of the patients of breast cancer in India present a decade earlier as compared to western country. Mean age at diagnosis of breast cancer in

western countries is 61 years and in India it is 50 years. Breast cancer is more common in postmenopausal group as compared to premenopausal group. Most patients of breast cancer come with complaint of painless breast lump followed by breast pain and nipple discharge. Most of patients in India are diagnosed when patients have early invasive cancer followed by locally advanced cancer.

Most common type of breast carcinoma is infiltrating ductal carcinoma - NOS. Most patients have lymph node positivity and histological grade 2 in histopathological report. About 60-65% all breast cancer are hormone receptor positive and remaining are hormone receptor negative. Postmenopausal women have higher hormone receptor positivity. Proportion of lymph node positivity and histological grade 2 is high in hormone receptor positive group. Most patients receive chemotherapy and radiotherapy after surgery followed by hormonal therapy according to hormone receptor status. Neoadjuvant chemotherapy can be given for down staging of locally- advanced breast cancer. Most common chemotherapy regime is Cyclophosphamide + Adriamycin +5- Flurouracil or Adriamycin+ Cyclophosphamide followed by Paclitaxel/Docetaxel as chemotherapy. Hormone receptor negative patients have high risk of recurrence.

REFERENCES

- Hunt KK, Robertson JFR, Bland KI. The Breast, In: Brunicaardi FC, Anderson DK, Billiar TR, Dunn DL, Hunter JG, Matthews JB, et al, eds. Schwartz's Principles of Surgery, 10th Edition. McGraw-Hill Education; 2015;17:467-564.
- Breast cancer. Hormone receptor status, Breastcancer. Available at https://www.breastcancer.org/symptoms/diagnosis/hormone_status
- ICMR. Consensus Document for management of breast cancer. ICMR. 2016. Available at <https://www.icmr.nic.in/sites/default/files/guidelines/Buccal%20Mucosa%20Cancer%20final%20pdf%209.6.14.pdf>
- Lester SC. Breast, In: Kumar V, Abbas AK, Fausto N, Aster JC. Robbins and Cotran pathologic basis of disease. 8th edition. Saunders, an imprint of Elsevier Inc; 2010;23:2058-2078.
- Leong SPL, Shen ZZ, Liu TJ, Agarwal G, Tajima T, Paik N et al. Is breast cancer the same disease in Asian and western countries? *World J Surg.* 2010;34:2308-24.
- Sainsbury R, Breast, In: Williams NS, Christopher JK, Ronan BP, O'Connell. Bailey and Love's short practice of surgery. 25th edition. Edward Arnold (Publishers) Ltd; 2008;50:827-850.
- Alsaed EF, Abdulkarim H, Tunio MA. Elevated preoperative serum cancer antigen 15.3 levels are associated with reduced disease-free survival: a single-institution experience. *Breast Cancer: Targets and Therapy.* 2013;5:53.
- Al-Azawi D, Kelly G, Myers E, McDermott EW, Hill AD, Duffy MJ, et al. CA 15-3 is predictive of response and disease recurrence following treatment in locally advanced breast cancer. *BMC Cancer.* 2006;6(1):220.
- Ali HQ, Mahdi NK, Al-Jowher MH. The value of CA15-3 in diagnosis, prognosis and treatment response in women with breast cancer. *J Pak Med Assoc.* 2013;63(9):1138-41.
- Kopans DB. Diagnostic breast imaging: textbook of breast cancer. LWW; 2006;9:116-120.
- Gion M, Mione R, Nascimben O, Valsecchi M, Gatti C, Leon A, et al. The tumour associated antigen CA15.3 in primary breast cancer: evaluation of 667 cases. *British J Cancer.* 1991;63(5):809.
- Dutta V, Chopra GS, Sahai K, Nema SK. Hormone receptors, HER-2/neu and chromosomal aberrations in breast cancer. *MJAFI.* 2008;64:11-5.
- Vasdhia BM, Bharti JM, Prashant PM. Correlation of hormone receptor and HER-2/neu expression in breast cancer: a study at tertiary care centre in south Gujarat. *Nat J Med Res.* 2012;2:295-8.
- Azizun-Nisa, Bhurgri Y, Raza F, Kayani N. Comparison of ER, PR and HER-2/neu (C-erb B 2) reactivity pattern with histologic grade, tumor size and lymph node status in breast cancer. *Asian Pac J Cancer Prev.* 2008;9:553-6.
- Sofi GN. Estrogen receptor and progesterone receptor status in breast cancer in relation to age, histological grade, size of lesion and lymph node involvement. *Asian Pacific J Cancer Prev.* 2012;13(10):5047-52.
- Lal P, Tan LK, Chen B. Correlation of HER-2 status with estrogen and progesterone receptors and histologic features in 3,655 invasive breast carcinomas. *Am J Clin Pathol.* 2005;123:541-6.
- Shet T, Agrawal A, Nadkarni M. Hormone receptors over the last 8 years in a cancer referral center in India: what was and what is? *Indian J Pathol Microbiol.* 2009;52:171.
- Allemani C, Weir HK, Helena. Global surveillance of cancer survival 1995-2009: analysis of individual data for 25 676 887 patients from 279 population-based registries in 67 countries (CONCORD-2). *The Lancet.* 2015;385(9972):977-1010.
- Naeem M, Nasir A, Aman Z, Ahmad T, Samad A. Frequency of HER-2/neu receptor positivity and its association with other features of breast cancer. *J Ayub Med Coll Abbottabad.* 2008;20(3).
- Patey DH, Dyson WH. The prognosis of carcinoma of the breast in relation to the type of operation performed. *Br J Cancer.* 1948;2:7-13.
- Williams MR. Survival patterns in hormone treated advanced breast cancer. *British J Surg.* 1986;73:752-5.