



## A CLINICAL STUDY OF ECTOPIC PREGNANCIES IN A TERTIARY CARE HOSPITAL IN NORTH EASTERN PART OF HARYANA.

<b>Dr. Tanisha*</b>	Second Year Post Graduate Student, Department of Obstetrics & Gynaecology, MMIMSR, Mullana, Ambala, Haryana. *Corresponding Author
<b>Dr. Tajinder Kaur</b>	Professor, Department of Obstetrics & Gynaecology, MMIMSR, Mullana, Ambala, Haryana.
<b>Dr. Reena Bisht</b>	Third Year Post Graduate Student, Department of Obstetrics & Gynaecology, MMIMSR, Mullana, Ambala, Haryana.

**ABSTRACT** **Background:** Ectopic pregnancy is an abnormal condition in which implantation of the blastocyst occurs outside the endometrium of the uterus. It is gynecological important, particularly in the developing world, because of its association with enormous rate of high morbidity, during the first trimester of pregnancy. A better understanding of its risk factors can help to prevent its prevalence. Diagnosis is frequently missed and should be considered in any woman in the reproductive age group presenting with abdominal pain or vaginal bleeding.

**Aims and Objectives:** The aim of this study is to determine the incidence, clinical presentation, risk factors, treatment, morbidity and mortality associated with ectopic pregnancy.

**Methods:** The present retrospective study was conducted in the Department of Obstetrics & Gynaecology, Maharishi Markendeshwar Institute of Medical Sciences and Research (MMIMSR), Mullana, Ambala, Haryana, India during the period from September 2020 to November 2021. Retrospective analysis of case histories of patients admitted with ectopic pregnancy at our college was done. The following parameters: age, parity, gestational age, risk factors, clinical presentation, need for blood transfusion and findings on ultra-sonogram and at surgery and morbidity associated with ectopic pregnancy were noted.

**Results:** In this study, a total of 31 cases of ectopic pregnancy were operated giving the incidence of ectopic pregnancy of 5.6/1000 deliveries. The commonest risk factors present were history of abortion (29%), history of tubal surgery (9.6%), infertility (3.2%) and pelvic inflammatory diseases (3.2%). The commonest symptoms were abdominal pain (80.6%), amenorrhea (77.4%) and abnormal vaginal bleeding (61.3%); and commonest signs were abdominal tenderness (64.5%), cervical excitation (51.6%) and adnexal tenderness (48.4%). Surgery by open method in the form of salpingectomy (90.3%), salpingo-oophorectomy (6.5%) and salpingostomy (3.2%) were the mainstay of management. Morbidity included anemia (41.9%), blood transfusion (54.8%) and wound infection (32.2%). No maternal mortality were noted in the present study.

**Conclusions:** Common risk factors for EP must be identified. Early diagnosis and timely intervention in the form of conservative or surgical treatment will help in reducing the morbidity and mortality associated with ectopic pregnancy. Since ectopic pregnancy remains a gynecological catastrophe in developing country like India, it should be considered a relevant public health issue. With its rising incidence, which is likely to continue increasing because of the various risk factors, it is necessary to devise means of early detection and treatment.

**KEYWORDS :** Ectopic pregnancy, Amenorrhoea, Tubal Pregnancy, Risk Factors, Vaginal bleeding, Salpingectomy, Hemoperitoneum.

### INTRODUCTION:

Ectopic pregnancy (EP) is an abnormal condition in which implantation of the blastocyst occurs outside the endometrium of the uterus. These abnormal sites of implantation in decreasing order of frequency include uterine tube (tubal pregnancy), abdominal cavity or on the mesentery (abdominal pregnancy), and in the ovaries (ovarian pregnancy) [1,2]. Blastocysts that do not implant in the uterine wall are generally unable to develop normally because; the space is incapable for developing blastocyst. Ectopic pregnancy can cause ruptures of fallopian tube, cervix and abdomen on which they are implanted. Rupture of ectopic pregnancy result in severe bleeding, many organ damage, and maternal mortality [3,4,5].

An ectopic pregnancy occurs when a fertilized ovum implants outside the normal uterine cavity [6]. It is the most important cause of maternal mortality and morbidity in the first trimester [7]. In developing countries, a majority of hospital based studies have reported ectopic pregnancy case-fatality rates of around 1%–3%, 10 times higher than those reported in developed countries [8]. Although women with ectopic pregnancy frequently have no identifiable risk factors, a prospective case-controlled study has shown that increased awareness of ectopic pregnancy and a knowledge of the associated risk factors helps identify women at higher risk in order to facilitate early and more accurate diagnosis [9]. Most risk factors are associated with risks of prior damage to the Fallopian tube. These factors include any previous pelvic or abdominal surgery, and pelvic infection [9].

Chlamydia trachomatis has been linked to 30-50% of all ectopic pregnancies [10]. Patients with an ectopic pregnancy commonly present with pain and vaginal bleeding between 6 and 10 weeks' gestation [6]. The diagnosis of ectopic pregnancy has become more frequent during the last decades but the incidence of ectopic pregnancy rupture has declined. This decline is due to quantitative hCG measurements, minimally invasive surgery and transvaginal

ultrasonography [11]. Early diagnosis reduces the risk of tubal rupture and allows more conservative medical treatments to be employed [12]. This retrospective analysis was done to determine the incidence, clinical features, risk factors, treatment and morbidity and mortality associated with ectopic pregnancy in a tertiary care hospital.

### METHODS:

The present retrospective study of ectopic pregnancies was conducted in the Department of Obstetrics & Gynaecology, Maharishi Markendeshwar Institute of Medical Sciences and Research (MMIMSR), Mullana, Ambala, Haryana, India during the period from September 2020 to November 2021. The case sheets of the patients with ectopic pregnancy were traced through the labour ward registers and operation theatre registers. Information regarding the total number of deliveries in the study period, details of demographic characteristics, clinical symptoms and signs, diagnostic tools used, treatment, risk factors for the ectopic pregnancy as well as associated morbidity and mortality were obtained. All the surgeries were partial/total salpingectomy done by open laparotomy and spinal/general anesthesia was used in all the cases in the present study. Data was entered in Microsoft Excel spreadsheet and analysed using SPSS software version 19.0. For categorical variables, data was compiled as frequency and percent. For continuous variables, data was calculated as mean  $\pm$  SD.

### RESULTS:

During the study period, there were a total of 5497 deliveries in our hospital and 31 cases of ectopic pregnancies were operated giving the incidence of ectopic pregnancies of 5.6/1000 deliveries. A majority of the patients (74.2%) belonged to the age group of 25-30 years. 16.1% were primiparas and 83.9% were multiparas.

Risk factors were previous history of tubectomy (3.2%), spontaneous and induced abortion (29%), 3.2% had a history of infertility. Copper-

T was inserted in 6.4% cases. 12.9% had a history of D and C; 9.7% gave a history of taking MTP pills. A history of previous ectopic in (3.2%) where partial salpingectomy was done. 12.9% had a history of previous LSCS and history of PID was found in 3.2% of the case (Table 1).

**Table 1: Risk Factors For ECTOPIC Pregnancy.**

Risk Factors	Numbers	%
Previous Abortion	9	29
Intrauterine Contraceptive Device	2	6.4
Previous Ectopic	1	3.2
Pelvic Inflammatory Disease	1	3.2
Infertility Treatment	1	3.2
Tubal Surgery	1	3.2

The commonest presenting complaints were abdominal pain (80.6%), amenorrhea (77.4%) and abnormal vaginal bleeding (61.3%). Abdominal tenderness was present in 64.5%, cervical excitation in 51.6% cases and adnexal tenderness in 48.4%. The urinary pregnancy test was positive in 87.1% of the cases and Ultrasound revealed ectopic pregnancy in 77% cases. Seasonal variation was noted in the ectopic pregnancies. A majority of the ectopic pregnancies (61.3%) were found between July and December. The commonest site of location of the ectopic pregnancy was in the ampulla of the fallopian tube. Other sites were fimbria, cornu, isthmus, heterotopic pregnancy and ovarian pregnancy as mentioned in (Table 2).

**Table 2: Site Of ECTOPIC Pregnancies On Laparotomy.**

Site Of Ectopic Pregnancy	Number	%
Ampulla Of Fallopian Tube	14	45.2
Fimbria	7	22.6
Cornu	6	19.4
Isthmus	2	6.5
Adhesions to Bowel Serosa	4	12.9
Ovarian	2	6.5
Heterotopic	2	6.5

Right sided tubal pregnancy was present in 18(62.1%) cases and left tubal involvement in 11(37.9%) cases. Ruptured ectopic pregnancy was present in 61.3% cases on laparotomy, 22.5% had unruptured ectopic and tubal abortion in 12.9% cases. 56% of the cases showed a hemoperitoneum on laparotomy. The most common surgery done was unilateral salpingectomy in 28 (90.3%), salpingoophorectomy in 2 (6.5%) and salpingostomy in 1 (3.2%). Morbidity included anemia (41.9%), blood transfusion (54.8%) and wound infection (32.2%). No maternal mortality was noted in the present study.

## DISCUSSION:

The present study was a hospital based retrospective study, which was aimed to identify determinants of ectopic pregnancy among pregnant women attending a tertiary hospital in Ambala, in the North-Eastern edge of Haryana. The prevalence of ectopic pregnancy among women who go to an emergency department with first trimester bleeding, pain, or both ranges from six to 16 percent [13]. In the present study, incidence of ectopic pregnancy was 5.6/1000 deliveries. In a study conducted by Rashmi Gaddagi and AP Chandrashekhar, the incidence was 1: 399 pregnancies [14]. In Porwal Sanjay *et al* study, the incidence was 2.46 per thousands of deliveries [15].

A majority of the patients (74.2%) belonged to the age group of 25-30 years in our study. Similar results were found in Khaleeqe *et al* study [16]. Hoover KW and colleagues reported that the ectopic pregnancy rate increases with age; it was 0.3% among girls and women aged 15-19 years and 1.0% among women aged 35-44 years [17]. 83.9% were multiparas and 16.1% were Primiparas. Multiparous women were found to be more prone to have ectopic pregnancy (61%) in Laxmi Karki *et al* study [18].

The commonest predisposing factors were tubectomy, pelvic inflammatory disease, spontaneous and induced abortion, and history of infertility, prior history of Copper-T insertion, D and C, previous ectopic pregnancies and previous LSCS. Similar risk factors were noted in various other studies [15]. Roussos D *et al* in their study observed that rupture of the tube is more often observed in women with a history of ectopic pregnancy and in women with full-term pregnancy [19].

The commonest presenting complaints were abdominal pain,

amenorrhea and abnormal vaginal bleeding. Clinical signs included abdominal tenderness, cervical excitation and adnexal tenderness. In Porwal Sanjay *et al* study, 87.5% reported with pain abdomen, bleeding per vagina encountered in 67.5% and 90% of cases had history of amenorrhea ranging from 6 weeks to 4months. These features help in early diagnosis of ectopic pregnancies [15]. The urinary pregnancy test, Serum  $\beta$ -hCG and ultrasound were the diagnostic tools used for diagnosis of ectopic pregnancy. Studies have shown that Ultrasonography should be the initial investigation for symptomatic women in their first trimester; when the results are indeterminate, the serum  $\beta$  human chorionic gonadotropin concentration should be measured. Serial measurement of  $\beta$ -hCG and progesterone concentrations may be useful when the diagnosis remains unclear [20].

In the present study, seasonal variation was noted in the ectopic pregnancies. A majority of the ectopic pregnancies were found between July and December. Mamdoh Eskandar and colleagues in a study conducted in Abha Maternity Hospital, Saudi Arabia reported that there was a seasonal variation in the incidence of ectopic pregnancy with highest mean incidence in the winter season [21]. Studies have shown that as a result of the influence of season on ovarian activity, it may be plausible to anticipate a seasonal variation in the incidence of ectopic pregnancy [22].

The commonest site of location of the ectopic pregnancy was in the ampulla of the fallopian tube. Ampulla part of the tube was commonly involved in most of the ectopic pregnancies in other studies [23] also. Heterotopic pregnancy was present in 6.5% of the ectopic pregnancies. Studies have shown that in a natural conception cycles, heterotopic pregnancy is a rare event, occurring in <1/30,000 pregnancies [24]. The high index of suspicion is to ensure for early and timely diagnosis and management, a timely intervention can result in a successful outcome of intrauterine pregnancy and prevent tubal rupture and hemorrhagic shock which can be fatal [25]. Right sided tubal pregnancy was present in 18(62.1%) cases and left tubal involvement in 11(37.9%) cases, consistent with other studies [26]. Ruptured ectopic pregnancy was present in 61.3% cases, 22.5% had unruptured ectopic and tubal abortion in 12.9% cases. In Latchaw G *et al* study, tubal rupture was present in 59% cases and 41% had unruptured ectopic pregnancies. They concluded that the patients with a history of a previous ectopic pregnancy are significantly more likely to experience a tubal rupture [27]. Studies have shown that low hemoglobin and hematocrit values, together with higher gravidity at the time of admission, may indicate an increased risk of tubal rupture [28]. In the present study, 83.9% were multiparas and 41.9% women were anaemic at the time of admission.

The most common surgeries performed were partial/total salpingectomy, salpingoophorectomy and salpingostomy. Conservative surgery is superior to radical surgery at preserving fertility. Conservative surgery is not followed by an increased risk of repeat ectopic pregnancy, but by the risk of persistent ectopic pregnancy, which should be taken into account when deciding on the operative procedure [29]. Canis M *et al* in their study concluded that the surgical treatment should be performed if the patient is hemodynamically unstable,  $\beta$ -hCG is >10000 mIU/mL, the ectopic pregnancy is 4cm in diameter, if there is a medical contraindication to methotrexate, and if the patient may not be followed adequately after treatment [30].

Mahboob reported a success rate of 80% by treating 12 out of 15 women with single dose MTX with initial  $\beta$ -hCG levels equal to 5000mIU/ml [7]. As medical management needs extremely close follow up & hospitalization, surgical management is still the method of choice in our country [31]. Laparoscopy and medical therapy have now emerged as the widely used therapeutic modalities with great success terms of reduced morbidity, shorter hospital stay and conservation of fertility [32].

However, choice depends upon early identification of ectopic pregnancy and stable condition of patients [33]. Morbidity included anaemia, blood transfusion and wound infection. By reducing and identifying the risk factors and 'catching' the patients at the earliest it is possible to improve the prognosis so far as morbidity, mortality and fertility are concerned [34].

In our study no maternal mortality was found, which was consistent with A. Abbas and H. Akram study [35]. Studies suggest that around

60% of women affected by an ectopic pregnancy go on to have a viable IUP. There is thought to be a 5-20% risk of a recurrence of ectopic pregnancy with one previous ectopic pregnancy and a risk of 32% or more following more than one previous ectopic [35].

Prevention and treatment of PID and encouraging women to undergo an early transvaginal ultrasonography to confirm the location of pregnancy is likely to prevent late diagnosis. This will also allow medical management or fertility sparing conservative surgical management. Setting up Early Pregnancy Assessment Units has been shown to result in higher quality and cost-effective care, and to have a positive effect on early pregnancy care in the UK [36]. Future studies are required to evaluate usefulness of such EPAUs and feasibility of setting them up in India.

Ultrasonography being the mainstay for evaluating Ectopic Pregnancy (EP), its availability at the point of care will also help majority of patients by allowing safe and timely discharge of patients presenting to emergency departments with clinical suspicion of an EP [37]. Future research may be directed at assessing the impact of training doctors at primary and secondary levels of healthcare with two-week intensive ultrasonography courses, on the mortality and morbidity associated with Ectopic Pregnancy.

### CONCLUSION:

It was concluded from our study that early identification of underlying risk factors, diagnosis with the essential aids like transvaginal ultrasound and  $\beta$ -hCG and timely intervention in the form of medical or surgical treatment will definitely help in reducing the morbidity and mortality associated with ectopic pregnancy and will also improve the future reproductive outcome. Hospitals should give emphasis on prevention and early detection of risks of ectopic pregnancy and create awareness in order to reduce the burden of ectopic pregnancy in a resource poor country like India.

### REFERENCES:

- Bhuria V, Nanda S, Chauhan M, Malhotra V. A retrospective analysis of ectopic pregnancy at a tertiary care Centre: one year study. *Int J Reproduct Contracept Obstetr Gynecol.* 2016;5:2224-7.
- Gaym A. Maternal mortality studies in Ethiopia--magnitude, causes and trends. *Ethiop Med J.* 2009;47(2):95-108.
- Yeasmin MS, Uddin MJ, Hasan E. A clinical study of ectopic pregnancies in a tertiary care hospital of Chittagong, Bangladesh. *Chattagram Maa-O-Shishu Hosp Med Coll J.* 2014;13(3):1-4.
- Gharoro EP, Igbafe AA. Ectopic pregnancy revisited in Benin City, Nigeria: analysis of 152 cases. *Acta Obstet Gynecol Scand.* 2002;81(12):1139-43.
- Basnet R, Pradhan N, Bharati L, Bhattarai N, Bb B, Sharma B. To determine the risk factors associated with ectopic pregnancy. *Asian J Pharm Clin Res.* 2015;8(2):93-7.
- Walker JJ. Ectopic pregnancy. *Clin Obstet Gynecol.* 2007; 50: 89-99.
- Mahboob U, Mazhar SB. Management of ectopic pregnancy: a two-year study. *J Ayub Med Coll Abbottabad.* 2006 Oct-Dec; 18(4):34-7.
- Goyaux N, Leke R, Keita N, Thonneau P. Ectopic pregnancy in African developing countries. *Acta Obstet Gynecol Scand* 2003; 82:305-12.
- Karaer A, Avsar FA, Batioglu S. Risk factors for ectopic pregnancy: a case-control study. *Aust N Z J Obstet Gynaecol.* 2006; 46:521-527.
- Turner C, Horner P, et al. British Fertility Society Impact of Chlamydia trachomatis in the reproductive setting; British Fertility Society Guidelines for practice. *Hum Fertil (Camb)* 2010; 13:115-125.
- Timmerman D. Predictive models for the early diagnosis of ectopic pregnancy. *Verh K AcadGeneesk Belg.* 2004; 66(2):155-71.
- Barnhart KT. Clinical practice. Ectopic pregnancy. *N Engl J Med.* 2009; 361:379-387.
- Murray H, Baakdah H, Bardell T, Tulandi T. Diagnosis and treatment of ectopic pregnancy. *CMAJ.* 2005; 173(8):905.
- Rashmi AGaddagi, AP Chandrashekhara. A Clinical Study of Ectopic Pregnancy. *JCDR* 2012;6:867-869.
- Gupta R, Porwal S, Swarnkar M, Sharma N, Maheshwari P. Incidence, trends and risk factors for Ectopic Pregnancies in a tertiary care hospital of Rajasthan. *JPBMS* 2012; 16(07):1-3.
- Khaleeque F, Siddiqui RI, Jafarey SN. Ectopic pregnancies: A Three year study. *J Pak Med Assoc* 2001; 51:240-243.
- Hoover KW, Tao G, Kent CK. Trends in the diagnosis and treatment of ectopic pregnancy in the United States. *Obstet Gynecol.* 2010 Mar; 115(3):495-502.
- R. C (Karki) L, Pradhan B, Duwa S. Annual Analysis of Ectopic Pregnancy in Tertiary Care Hospital. *PMJN* 2011; 11:5-8.
- Roussos D, Panidis D, Matalliotakis I, Mavromatidis G, Neonaki M, Mamopoulos M, Koumantakis E. Factors that may predispose to rupture of tubal ectopic pregnancy. *Eur J Obstet Gynecol Reprod Biol.* 2000 Mar; 89(1):15-7.
- Murray H, Baakdah H, Bardell T, Tulandi T. Diagnosis and treatment of ectopic pregnancy. *CMAJ.* 2005 Oct 11; 173(8):905-12.
- Esikandar M, Archibong EI, Sadek AA, Sobande AA. *Bahrain Med Bull* 2002; 24(2):63-65.
- Warren CW, Gwinn ML, Rubin GL. Seasonal variation in conception and various pregnancy outcomes. *Soc Biol* 1986; 33:116-26.
- Swende TZ, Jogo AA. Ruptured tubal pregnancy in Makurdi, North Central Nigeria. *Niger J Med* 2008; 17(1):75-7.
- Ludwig M, Kaisi M, Bauer O, Diedrich K. Heterotopic pregnancy in a spontaneous cycle: do not forget about it! *Eur J Obstet Gynecol Reprod Biol.* 1999 Nov; 87(1):91-3.
- Espinosa PM, Alcantar Mendoza MA. Heterotopic pregnancy: Report of a case and review of literature. *Gynecol Obstet Mex.* 1997; 65:482-6.
- GO Udigwe, OS Umeononihu, I.I Mbachu. Ectopic pregnancy: A 5 year review of cases at Nnamdi Azikiwe University Teaching Hospital (NAUTH) Nnewi. *NMJ* 2010 ; 51: 160-163.
- Latchaw G, Takacs P, Gaitan L, Geren S, Burzawa J. Risk factors associated with the

- rupture of tubal ectopic pregnancy. *Gynecol Obstet Invest.* 2005; 60(3):177-80.
- Knafel A, Basta P, Skotniczny K, Pawel M, Krzysztof B, Rokita W, Obrzut B, Wicherek L. Ectopic pregnancy rupture--can it be prevented? *Ginekol Pol.* 2009 Oct; 80(10):734-9.
- Bangsgaard N, Lund CO, Ottesen B, Nilas L. Improved fertility following conservative surgical treatment of ectopic pregnancy. *BJOG* 2003 Aug; 110(8):765-70.
- Canis M, Savary D, Pouly JL, Wattiez A, Mage G. Ectopic pregnancy: criteria to decide between medical and conservative surgical treatment? *J Gynecol Obstet Biol Reprod (Paris).* 2003 Nov; 32(7 Suppl):S54-63.
- Chatterjee S, Dey S, Chowdhury R G, Ganguli D. Ectopic Pregnancy In Previously Infertile Women Subsequent Pregnancy Outcome After Laparoscopic Management. *Al Ameen J Med Sci* 2009; 2(1):67-72.
- Jurkovic D. Ectopic pregnancy. In: Edmonds DK, editor. *Dew Hurst's textbook of Obstetrics & Gynecology.* 7 ed. USA: Blackwell Publishers; 2007.
- Shah N, Khan NH. Ectopic pregnancy: Presentation and risk factors. *J Coll Physicians Surg Pak.* 2005; 15:535-8.
- Majhi AK, Roy N, Karmakar KS, Banerjee PK. Ectopic pregnancy--an analysis of 180 cases. *J Indian Med Assoc.* 2007 Jun; 105(6):308,310,31.
- Abbas A, Akram H. Ectopic Pregnancy; Audit at Maula Bakhsh Teaching Hospital Sargodha. *Prof Med J* 2011; 18(1): 24-27.
- Van den Berg MMJ, Goddijn M, Ankum WM, van Woerden EE, van der Veen F, van Wely M, et al. Early pregnancy care over time: should we promote an early pregnancy assessment unit? *Reprod Biomed Online.* 2015; 31(2):192-98. [PubMed] [Google Scholar].
- French S, Henry T, Williams E. Evaluation of Waiting Times and Sonographic Findings in Patients with First Trimester Vaginal Bleeding at the University Hospital of the West Indies. *Can Emergency Department Ultrasound Make a Difference? West Indian Med J [Internet]* 2014 Jul 21 [cited 2015 Nov 17] Available from: <http://myspot.mona.uwi.edu/fms/wimj/article/1854>. [PMC free article] [PubMed] [Google Scholar].